Management, blood and reproductive parameters of rabbits reared in Imo State

Iwuji, T. C., Nwapi, D. O., Ogbuewu, I. P., Kadurumba, E. O., Egenuka, F. C. and Okere, P. C.

Department of Animal Science and Technology, Federal University of Technology,

P. M. B. 1526, Owerri, Imo State, Nigeria.

Corresponding author: tiwuji@gmail.com

Abstract

The management, haematological, serum biochemical and reproductive parameters of rabbits reared in the 3 zones (Okigwe, Orlu and Owerri) of Imo State were investigated. Data for this study were collected through questionnaire, on-the-farm assessment and fluid sample (blood and semen) from the rabbits. The hutch/cage type of housing is the most prevalent in Imo State, constituting 73.58 %. Most rabbit farmers in Imo State (45.28 %) combine concentrate and forage in rabbit feeding, 47.17 % of the rabbit house/environment were fairly clean, 30.19 % dirty and 22.64 % were clean. Rabbits with ages ranging from 5 – 8 months recorded average live weight of 1.91 kg, while annual average kindling and litter size of rabbit does reared in Imo State were 5.67 and 7.43, respectively; with average litter survival rate of 60.16 % between parturition and weaning. Haematological analysis of the adult rabbits recorded 27.80 %, 9.56 g/dl and $3.40 \times 10^6/\text{ml}$ in packed cell volume, haemoglobin concentration and red blood cell count, respectively, which were lower than normal reference values. Total serum protein, globulin and urea values were lower than normal reference values recording 4.44 g/dl, 1.57 g/dl and 16.56 mg/dl, respectively. Semen parameters obtained from adult rabbit bucks were inferior to semen parameters of adult rabbit bucks reared in similar geographical locations and environment, but under standard management practices. This study was therefore designed to assess the level of management, blood and reproductive parameters of rabbits in Imo State.

Keywords: Management, rabbits, blood parameters, reproductive parameters, Imo State.

Introduction

There is need to fill the gap in animal protein need of developing countries and the rabbit (*Oryctolagus cuniculus*) which has been described as a micro livestock specie is likely the cheapest and sustainable means of producing high quality animal protein for expanding population of developing countries like Nigeria (Onifade *et al.*, 1999); no thanks to less attention given to rabbit production in developing countries, unlike other livestock species like poultry, sheep, goat and cattle (Assan, 2014). Rabbits have exceptional attributes which include short generational and gestation intervals, fecundity, rapid growth rate, genetic

diversity ability to utilize forage and agricultural by-products and adaptation over a wide range of ecological environment (Arijeniwa *et al.*, 2000). These attributes of rabbits are the reasons why rabbit production can provide the impoverished urban population and the resource-poor rural dwellers the opportunity to earn additional income on a sustainable basis (Onifade *et al.*, 1999). Furthermore, increased rabbit production, which is basically the increase in the number of rabbits kindled per doe per year and the survival of the kits (Lukefahr *et al* 1991), will increase the availability of research animals for researchers and students in higher institutions (Iwuji, 2016).

Management and Physiological parameters of rabbits in Imo State

Unfortunately, rabbit production is still at subsistent level or emergence in Nigeria (Onifade *et al.*, 1999). Traditionally, rabbits have not been included to the food of the community, however, some people keep rabbits for meat (as bush meat delicacy), as pet or for research purposes (khan *et al.*, 2014). Recently, especially in Imo State, sourcing rabbits for research works has turned into a nightmare and are very costly when seen, most times the needed number or types may not be available. Therefore, it becomes imperative to conduct this study on the assessment of rabbit production in Imo State, with emphasis on their management, blood and reproductive parameters, for the realization of possible solutions to rabbit scarcity.

Materials and methods

This study lasted for three months and covered the three zones (Okigwe, Orlu and Owerri) in Imo State (Figure 1). Imo State is situated in the Southern rain forest vegetation belt of Nigeria. It lies within latitudes 4°45'N and 7°15'N, and longitude 6°50'E and 7°25'E. The area is dominated by plains 200 m above sea level except for elevations associated with the Okigwe uplands. It has an annual rainfall of about 1700 to 2500 mm, which is concentrated almost entirely between March and October. Average relative humidity is about 80 % with up to 90 % occurring during the rainy season. The mean daily maximum air temperature ranges from 28 to 35 °C while the mean daily minimum ranges from 19 to 24 °C (Okoli *et al.*, 2007).

Data for the assessment of rabbits reared in Imo State were collected based on the parameters outlined below, through a pre-designed questionnaire, on-the-farm assessment and laboratory analyses of blood and semen of the rabbits;

- **Management:** Management practices were evaluated based on housing, feeding and cleanliness of the rabbit houses or environment. The rabbits in Imo State were mainly housed in hutches/cages, open room and burrows. It was observed that burrows were not intentionally provided by the farmers, but were dug by the rabbits kept on non-cemented floors or broken cemented floors, thereby exposing the earth.
- **Performance and production:** This was assessed in terms of live weight, number of kindling per annum, litter size and litter survival rate of the rabbits.
- **Purpose of rabbit farming:** Assessment of the reason(s) why each farmer engaged in rabbit farming was classified into meat production, source of revenue, research and entertainment.
- **Constraints to rabbit production:** This was assessed on the basis of fund, space, predators/theft and diseases.
- **Haematology:** Haematological parameters analyzed were packed cell volume, haemoglobin concentration, red blood cell count, white blood cell count and white blood cell differential counts.
- **Serum biochemistry:** Serum

biochemical parameters analyzed were total protein, albumin, globulin, urea, creatinine, glucose, cholesterol, alkaline phosphatase, alanine transaminase and aspartate transaminase.

- **Semen:** Semen volume, total sperm count, progressive sperm motility and live sperm proportion of the male rabbits were analyzed.

A total of 53 rabbit farms were identified in Imo State (Okigwe zone = 13, Orlu zone = 16, Owerri zone = 24) and were assessed physically and through interviews which was based on the questions in the questionnaire. Only live weight of adult male and female rabbits (5 months and above) were measured using a digital scale. The average number of kits the does kindle at a time was recorded as the litter size, while the average number of time the does kindle in a year was recorded as the annual kindling.

In each of the three zones, blood and semen samples were collected from six farms that were evenly distributed, for hematological, serum biochemical and semen analyses. In each farm selected, blood samples were collected from two adult male and female rabbits each, while semen were collected from three adult male rabbits in each farm selected. Blood samples were collected through the ear of the rabbits using needle and syringe. Blood samples for hematological analysis were discharged into Ethylene Diamine Tetra-acetic Acid (EDTA) bottles, while those for serum biochemical analysis were discharged into bottles without EDTA. Semen samples were collected with the help of a teaser doe, using artificial vagina (AV) as described by Herbert and Adejumo (1995). The blood and semen samples collected from the farms were analyzed at the Animal Science

Management and Physiological parameters of rabbits in Imo State

and Technology Laboratory, Federal University of Technology, Owerri.

Statistical analysis

The data base was prepared using Microsoft Excel and the mean values and percentage distribution of the various parameters computed using IBM SPSS software (2013).

Results and discussion

Management parameters of rabbit farms assessed in Imo State are presented in Table 1. The hutch/cage type of housing was the most predominant in Imo State, recording a total of 73.58 % than 20.75 and 5.66 % recorded for open room and burrow, respectively. Greater percentage (45.28 %)

of farmers in Imo State feed both concentrate and forage/household products to their rabbits, a slightly lesser percentage (43.40 %) feeds only forage/household products, while only a small proportion of the farmers (11.32 %) feed only concentrate to their rabbits; and are mainly those who cannot afford to go and look for forage or have difficulty in accessing them. In terms of cleanliness of the rabbit houses (clean, fairly clean or dirty), 47.17 % were fairly clean; 30.19 % were dirty, while 22.64 % were clean. It was observed that farmers who recorded clean rabbit environment were those who have knowledge (formal and informal) of rabbit production and they also keep some farm records.

Table 1: Distribution (%) of management parameters of rabbits reared in Imo State

Parameters	OKZ	OLZ	OWZ	Total
Housing				
Hutch/Cage	61.54	75.00	79.17	73.58
Open room	30.80	18.75	16.67	20.75
Burrow	7.69	6.25	4.17	5.66
Feeding				
Concentrate	15.38	6.25	12.50	11.32
Forage/Household products	46.15	81.25	16.67	43.40
Both	38.46	12.50	70.83	45.28
Cleanliness of rabbit house				
Clean	30.77	25.00	16.67	22.64
Fairly clean	30.77	37.50	62.50	47.17
Dirty	38.46	37.50	20.83	30.19

OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

At 5 – 8 months of age, the rabbits recorded an average live weight of 1.91 kg. Annual kindling and litter size of the does recorded 5.67 and 7.43, respectively, while average survival rate of the litters (Parturition – weaning) was 60.16 % (Table 2). Although these parameters are largely dependent on

the type of management, like nutrition, time of weaning and fattening period (Onifade *et al.*, 1999), they are generally low, compared to the values obtainable in temperate countries and in enhanced/adequate management situations (Onifade *et al.*, 1999; Oseni, 2012).

Table 2: Performance and production parameters of rabbits reared in Imo State

Parameters	OKZ	OLZ	OWZ	Total
Live weight (kg)	1.94	1.76	2.04	1.91
Annual kindling	5.8	5.00	6.2	5.67
Litter size	7.20	7.00	8.10	7.43
Litter survival rate (%)	58.33 (4.2)	68.57 (4.8)	54.32 (4.4)	60.16 (4.5)

OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

The survey result of this study on the purposes of rabbit farming in Imo State (Table 3) reveals that the farmers who rear rabbits for the purpose of revenue generation comprise 56.60 %, while farmers who rear rabbits for meat production, pet animal and research comprises 18.87, 15.09 and 9.43 %, respectively. It was observed that the purpose of the rabbit farming affected some

of the management practices carried out by the farmer. For instance, those who were rearing for research purpose were keeping some farm records and constitute major percentage of farm cleanliness recorded. Although during sourcing of rabbits for research, the student or researcher may buy from those whose purpose of rearing rabbits are not for research, which usually may result to non availability of the history or records of the rabbits bought.

Table 3: Distribution (%) of the purposes of rabbit farming in Imo State

Parameters	OKZ	OLZ	OWZ	Total
Meat production	15.38	37.50	8.33	18.87
Source of revenue	61.54	43.75	62.50	56.60
Research	7.69	6.25	12.50	9.43
Pet/Entertainment	15.38	12.50	16.67	15.09

OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

According to the farmers, 54.72 % of the constraints are fund-based, while space, predators/theft and diseases contribute 26.42, 7.55 and 7.55 %, respectively (Table 4). It is quite surprising that greater percentage of the farmers complained of

funds, when rabbit production venture has been adjudged to favour low investment, early benefits, subsistence on renewable resources for feeding, housing and general management (Onifade *et al.*, 1999; Oseni, 2012).

Table 4: Distribution (%) of the constraints to rabbit production in Imo State

Parameters	OKZ	OLZ	OWZ	Total
Fund	46.15	56.25	58.33	54.72
Space	23.08	25.00	29.17	26.42
Predators/Theft	7.69	12.50	4.17	7.55
Diseases	7.69	6.25	8.33	7.55

OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

Haematological analysis on the adult rabbits recorded average values of 27.80 %, 9.56 g/dl, $3.40 \times 10^6/\text{ml}$ and $11.67 \times 10^3/\text{ml}$, for packed cell volume (PCV), haemoglobin (Hb) concentration, red blood cell count (RBC) and white blood cell count (WBC), respectively (Table 5). The white blood cell differentials recorded 44.94, 49.77, 1.67, 2.33 and 1.00 % for lymphocytes, heterophils, monocytes and

basophils, respectively. The PCV, Hb and RBC of the rabbits were quite lower than the reference values documented for adult rabbits (Mitruka and Rawnsley, 1977; Merck, 1998) and could be attributed to poor nutrition of the rabbits, while the elevated WBC count and percent heterophils above the normal reference values may be due to morbidity resulting from poor management of the rabbits (Etim *et al.*, 2014).

Table 5: Haematological parameters of rabbits reared in Imo State

Parameters	OKZ	OLZ	OWZ	Total	Ref. values*
Packed cell volume (PCV) (%)	27.31	28.00	28.10	27.80	30 – 50
Haemoglobin (Hb) conc. (g/dl)	9.40	9.82	9.46	9.56	10 – 15
Red blood cells (x 10 ⁶ /ml)	3.28	3.81	3.10	3.40	3.6 – 7.5
White blood cells (x 10 ³ /ml)	12.31	12.28	10.41	11.67	4.5 – 11
Lymphocytes (%)	41.00	42.42	51.40	44.94	40 – 80
Heterophils (%)	53.30	51.80	44.21	49.77	35 – 43.2
Monocytes (%)	2.00	2.00	1.00	1.67	1 – 4
Basophils (%)	3.00	2.00	2.00	2.33	1 – 7
Eosinophils (%)	1.00	1.00	1.00	1.00	0 – 4

*(Mitruka and Rawnsley, 1977; Merck, 1998) . OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

Serum biochemical parameters of the adult rabbits are presented in Table 6 and recorded 4.44, 2.87 and 1.57 g/dl for average total protein (TP), albumin and globulin, respectively. Total protein and globulin values of the rabbits were below the recorded reference values (Mitruka and Rawnsley, 1977; Merck, 1998) which could be due to low amino acid profile of the feeds fed the rabbits. The low TP probably affected only the globulin values, since the albumin values were within the normal values. This may result in low production of immunoglobulin which helps

in the immune system of the rabbits (Frandsen *et al.*, 2009). The rabbits recorded average urea, creatinine, glucose and cholesterol values of 16.56, 2.33, 110.6 and 47.38 mg/dl, respectively, while the average values of alkaline phosphatase (ALP), alanine transaminase (ALT) and aspartate transaminase (AST) were 85.31, 53.26 and 83.88 IU/L, respectively. Urea is a product of protein metabolism (Frandsen *et al.*, 2009), therefore, it could be that the recorded average urea value of the rabbits, which is below the reference normal values may be as a result of the low TP of the rabbits.

Table 6: Serum biochemical parameters of rabbits reared in Imo State

Parameters	OKZ	OLZ	OWZ	Total	Ref. values*
Total Protein (g/dl)	4.42	4.81	4.10	4.44	5 – 7
Albumin (g/dl)	2.94	3.42	2.26	2.87	2.5 – 4
Globulin (g/dl)	1.48	1.39	1.84	1.57	2.5 – 4
Urea (mg/dl)	16.24	17.01	16.44	16.56	20 – 45
Creatinine (mg/dl)	2.41	2.32	2.26	2.33	0.5 – 2.6
Glucose (mg/dl)	101.2	110.4	120.2	110.6	75 – 155
Cholesterol (mg/dl)	43.45	51.44	47.24	47.38	10 – 80
Alkaline phosphatase (IU/L)	84.60	86.11	85.23	85.31	10 – 96
Alanine transaminase (IU/L)	53.21	54.14	52.44	53.26	55 – 260
Aspartate transaminase (IU/L)	81.43	84.12	86.08	83.88	10 – 98

*(Mitruka and Rawnsley, 1977; Merck, 1998) OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

Semen volume, sperm concentration, total sperm, progressive sperm motility and live sperm proportion of the adult rabbits recorded average values of 0.59 ml, 166.20 x 10⁶/ml, 98.30 x 10⁶, 58.54 % and 77.38 %, respectively (Table 7). Comparing the

semen parameters recorded in this study with semen parameters recorded in some empirical studies carried out in similar geographical locations and environment

Table 7: Semen parameters of rabbits reared in Imo State

Parameters	OKZ	OLZ	OWZ	Total	Ref. values*
Semen volume (ml)	0.51	0.65	0.62	0.59	0.3 – 0.9
Sperm conc. (x 10 ⁶ /ml)	188.49	151.20	158.90	166.20	250 – 600
Total sperm (x 10 ⁶)	96.41	100.10	98.40	98.30	-
Progressive sperm motility (%)	54.10	63.20	58.31	58.54	30 – 90
Live sperm proportion (%)	76.42	79.21	76.51	77.38	-

*(Merck, 1998) OKZ = Okigwe zone; OLZ = Orlu zone; OWZ = Owerri zone

with Imo State, but under standard management practices (Yousef *et al.*, 2003; Ahmed *et al.*, 2008; Iwuji and Herbert, 2012; Ogbuewu *et al.*, 2012), it was observed that the semen parameters of the rabbits reared in Imo State were inferior. Inferior semen parameters in adult rabbit bucks and some other animal models have been recorded with feeding of forage or administration of plant extracts without appropriate concentrate supplementation (Herbert *et al.*, 2005; Eze, 2012). Apart from high forage feeding of the rabbits (Table 1) without adequate concentrate supplementation, general poor management of the adult male rabbits may also be a contributing factor.

Conclusion

It was observed during the survey that rabbit farmers are very few in Imo state when compared with poultry farmers. Most of the rabbit farms were poorly managed in terms of nutrition, housing and hygiene, which may be the cause of low performance and production, poor reproductive characteristics and low survivability of the litters recorded in this study. The purposes of the majority (56.6 %) of rabbit farmers are to generate revenue, but unfortunately, they lack the adequate knowledge of rabbit production as evident in the performance of their farms and from the interactions with them.

Recommendations

Based on the findings of this study, we recommend the following;

- (i) That relevant agencies and institutions should embark on enlightenment campaign on the need and benefits of rearing rabbits.
- (ii) Provision of incentives and adequate knowledge of rabbit production to intending rabbit farmers.
- (iii) Increased research in rabbits so as to develop modern production techniques that will help increase rabbit performance and production.

References

- Ahmed, H. E., El-Sabei, M. E., Hamed, R. S. and Zeitouny, M. M. 2008. Effect of clomipramine hydrochloride on semen quality and biochemical parameters of rabbits. *Egypt. Poult. Sci.* Vol. 28 (3): 925 – 937.
- Arijiwa, A., Otaikhian, S. O. and Imaseum, J. A. 2000. Performance of weaner rabbits fed poultry grower mash supplemented with different grass legume. *Proceedings of 5th Annual Conference of Animal Science Association of Nigeria*, pp. 103 – 105.
- Assan, N. 2014. Micro-livestock farming and food security in Sub Saharan Africa. *J. Anim. Prod. Adv.* 4 (4): 374 – 387.

- Etim, N. N., Williams, M. E., Akpabio, U. and Offiong E. A. 2014.** Haematological parameters and factors affecting their values. *Agricultural Science* Vol. 2, (1): 37-47
- Eze, K. N. 2012.** Antifertility Effects of Ethanolic Extract of *Xylopi aethiopica* on Male Reproductive Organ of Wistar Rats. *American Journal of Medicine and Medical Sciences*, 2(1): 12-15.
- Fransson, R. D., Wilke, W. L. and Fails, A. D. 2009.** Anatomy and Physiology of Farm animals (7th edn). *Wiley-Blackwell Publishing*, USA.
- Herbert, U. and Adejumo, D. O. 1995.** Construction and evaluation of an artificial Vagina for collecting rabbit semen. *Delta Agric.* (2): 99-108.
- Herbert, U., Ozoje, M. O. and Adejumo, D. O. 2005.** Effect of leucaena and gliricidia leaf meals on the seminal characteristics, testis weights and seminiferous tubule diameters of rabbits. *Animal Research, EDP Sciences*, 54(3), pp.173-178.
- International Business Machine Corporation, 2013.** IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp.
- Iwuji, T. C. 2016.** Growth performance and physiological responses of rabbits administered *Panax ginseng* extracts. A PhD thesis submitted to the Department of Animal Breeding and Physiology, Michael Okpara University of Agriculture, Umudike.
- Iwuji, T. C. and Herbert, U. 2012.** Semen characteristics and libido of rabbit bucks fed diets containing *Garcinia kola* seed meal. *Rabbit Genetics* Vol. 2 (1):10-14.
- Khan, S., Khan, K., Shah, S. U. and Ahmad, N. 2014.** A preliminary assessment of rabbit farming and its scope in Kyber Pakhtunhwa Province of Pakistan. *Sarhad J. Agric.* 30 (30): 369-373.
- Lukefahr, S. D. and Cheeke, P. R. 1991.** Rabbit project development strategies in subsistence farming systems: 2. Research applications. *World Animal Review*, 69: 60-70.
- Merck Veterinary Manual. 1998.** A handbook of diagnosis and therapy for veterinarian. 8th edn. *Merck and Co. Rahway*, Ed. D. H. Siegmund.
- Mitruka, B. M. and Rawnsley, H. M. 1977.** Clinical, biochemical and haematological reference values in normal and experimental animals. *Masson Publishing, USA, Inc.*, 83: 134-135.
- Ogbuewu, I. P., Enumaibe, C., Kadurumba, O. E., Iwuji, T. C., Ogundu, U. E., Etuk, I. F., Opara, M. N., Okoli, I. C. and Illoeje, M. U. 2012.** Effects of dietary addition of toasted soybean seed meal on libido, semen quality and testis histology of rabbit bucks. *Scientific Journal of Medical Science*, Vol. 1(1) 7-12.
- Okoli, I. C., Alaehie, D. A., Akano, E. C., Ogundu, U. E., Okoli, C. G., Akujobi, C. T., Onyicha, I. D. and Chinweze, C. E. 2007.** Aerial Pollutant Gases Concentrations in Tropical Pig Pen Environment in Nigeria. *Research Journal of Environmental Sciences*, 1:110 - 115.
- Onifade, A. A., Abu, O. A., Obiyan, R. I. and Abanikannda, O. T. F. 1999.** Rabbit production in Nigeria: Some aspects of current status and

- promotional strategies. *World Rabbit Science*, Vol. 7 (2): 51 – 58.
- Oseni, S. O. 2012.** Rabbit production in low-input systems in Africa – Prospects, challenges and opportunities. World Rabbit Science Association. Proceedings 10th World Rabbit Congress – September 3 – 6 , Sharm El-Sheikh – Egypt, 719 – 731.
- Yousef, M. I., Abdallah, G. A. and Kamel, K. I. 2003.** Effect of ascorbic acid and Vitamin E supplementation on semen quality and biochemical parameters of male rabbits. *Anim. Reprod. Sci.* 76 (1-2): 99 - 111.

Received: 27th November, 2016

Accepted: 4th March, 2017