Correlation of milk intake and body weight with linear body measurements of white Fulani calves in semi-intensive system of management

Umoh, B. I., Okon, U. M. and Ekpo, J. S.

Department of Animal Science, Akwa Ibom State University, Obio Akpa Campus,
Akwa Ibom State, Nigeria.

Corresponding author: utibeokon93@gmail.com

Abstract

Body weight of animals in the rural areas has not been accurately ascertained due to lack of accurate measurement scales, hence this experiment. The study was conducted to determine the correlation between milk intake and body weight with linear body measurements in white Fulani calves raised in semi-intensive system of management. A total of seven (7) white Fulani calves managed semi-intensively was used for the study. The experiment lasted 12 weeks. The overall mean $\pm$ SEM value for milk intake, body weight, neck circumference, neck length, body length, body circumference, girth circumference, height at wither and tail length were 1.20 $\pm$ 0.40g, 5.18 $\pm$ 1.02kg, 21.54 $\pm$ 0.66cm, 13.23 $\pm$ 0.21cm, 28.36 $\pm$ 0.31cm, 36.41 $\pm$ 0.43cm, 35.84 $\pm$ 0.37cm, 35.87 $\pm$ 0.43cm and 23.43 $\pm$ 0.39cm respectively. There was significant ($P<0.05$) effect of lactation stage on all body measurements taken except for neck circumference and tail length. Correlation between all body pairs of measurements were high, positive and highly significant ($P<0.05$). The highest correlation was recorded for body length, body circumference, girth circumference and height at wither. Linear function provided the best fit for milk intake weight and linear body measurements indicating that body measurements can be used for selection and the estimation of body weight in the fields where accurate scales are not usually available. It is recommended that any of the body measurement parameters could be used in the estimation of the body weight. Also, an improvement of non-genetic factor such as nutrition and general husbandry of the animal as much as possible will bring about improvement in the body parameters as well as maintaining the animal's physiological balance.

Keywords: White Fulani, body weight, linear body measurements, correlation, milk intake

Introduction

The most populous and widely distributed breeds of cattle in Nigeria has been found to be the white Fulani cattle; many of which are reared under the transhumance farming system while a few belong to institutions and private farms (Dim *et al.*, 2012). Subjective pricing of meat animal is seen to be prevalent in developing countries against the objective marketing of live animals in developed countries. This is due to the ease of acquisition and setting up of weigh instruments (Laden *et al.*, 2009). Body weight is an important assessment index in animals but its measurement is usually not accurately ascertained in the rural areas due to lack of reasonable and accurate

measuring scales (Younas *et al.*, 2013). This therefore makes the rural farmers rely on systems and estimates that are open to doubt and ultimately leading to inaccurate decision making. Linear body measurements together with body weight measurements have been described by Salako (2006) to describe more completely an individual or population than the conventional methods of weighing and grading. Since young animals require milk as their first and only food owing to the fact that it contains important nutrients and is almost a complete food (Pandey and Voskuil, 2011), it therefore becomes pertinent to estimate the quantity of milk supplied to the young by the dam. This is basically for optimal development until

Correlation of milk intake and body weight with linear body measurements

weaning, for the perfection of new management techniques which would ultimately lead to flock productivity improvement. This study therefore was designed to determine the correlation of milk intake and body weight with linear body measurements in white Fulani calves at different stages of growth as well as investigate the prediction of live weight using some linear measurement parameters in white Fulani calves.

Materials and methods

Location of the study and management of the experimental animals

The experiment was conducted at the cattle unit of the Department of Animal Science Akwa Ibom State University, Obio Akpa, Akwa Ibom State. The area is situated between latitude 4°30'N and longitude 7°30'E (SLUK-AK, 1989). A total of seven (7) white Fulani calves were used for the study. The animals were managed under a semi-intensive system of animal husbandry at the cattle unit of the department of Animal Science, The animals were released for grazing from 9:00am to 4:00pm while the calves were restricted to their pens for the first 6 weeks and thereafter allowed to graze with the cows. The experimental animals were identified by ear tagging. The dam and the calves were put together and allowed to suckle for a period of 1 hour in the morning. The calves were weighed before suckling and after suckling using "Weigh Suckle Weigh (WSW)" method as adopted by Adewumi and Kazeem, 2010 and the difference between the weight before and after suckling gave the milk intake of each calf. The weight of each calf was taken with the aid of a spring balance. This procedure was repeated in the evening (10 hours later). The total of morning and evening weighing was the daily milk intake of each calf.

Milk Intake (kg) = weight after suckling (kg) – weight before suckling (kg).

Each of the calves selected for linear body measurements were confined during measurements. The body weight of the calves was taken in kilogram with the aid of a measuring scale. Seven metric traits were taken on individual calves with the aid of a flexible tailor's tape rule. Body measurement was taken in (cm) twice a week for 12 weeks as described by Younas *et al.* (2013). Reference points for the measurements were:

Neck Circumference (NC) was measured as the distance round the neck.

Neck Length (NL) was measured as the distance between the edges of the neck to the lower jaw region.

Body Circumference (BC) was the circumference around the body.

Body Length (BL) was measured from the joint of the scapular to the pin bone.

Girth Circumference (GC) was the circumferential measurement taken around the chest.

Height at Withers (HW) is the distance from the platform on which the animal stands to the point of its shoulders.

Tail Length (TL) was measured as the distance between the base of the tail close to the body of each calf and the tip of the tail.

Statistical analysis

The data collected were subjected to one way analysis of variance. Where F showed significant difference among the means, means were separated using Duncan's New Multiple Range Test (Duncan, 1995). Pearson Correlation was used to compare the relationship among the parameters. Regression analysis was used to obtain the best fit for body weight and linear body measurements.

Results and discussion

The overall mean±SEM value for MI, BW, NC, NL, BL, BC, GC, HW and TL were 1.20±0.40g, 5.18±1.02kg, 21.54±0.66cm, 13.23±0.21cm, 28.36±0.31cm,

36.41±0.43 cm, 35.84±0.37 cm, 35.87±0.43 cm and 23.43±0.39 cm respectively. There was no significant (P>0.05) effect of lactation on MI, TL and NC, however, there was significant (P<0.05) effect of lactation stage on NL, BL, BC, GC and HW. There was variability as the age of the animal increased most particularly in the weight gain. This result aligns with the report of Okon *et al.* (2018) and Orthenuata and Olutogun (1994) who reported that linear body measurement is closely related to live weight in cattle. MI increased between the early stages of the animal's growth and thereafter decreased as lactation progressed but all linear body measurements increased as the animal advanced in age. However, weight gain decreased as lactation progressed. This increment agrees with the work of

Adewumi and Kazeem (2010) in yankasa lambs, Singh *et al.* (1978) in grey and brown Bengal goats and Okon *et al.* (2018) in bunaji calves. The mean weight gain value (5.18) as observed in this study was slightly similar to the value of (5.15kg) reported by Okon *et al.* (2018) but lower than the value (3.25kg) reported by Afolayan *et al.* (2006) in a research with yankasa lambs. The low mean weight value of the calves could be attributed to the small number of animals used for the experiment as well as the inability of the calves at their later growth stage to utilize nutrients from forages eaten as they went along with the cows to feed on pasture due to inadequate fermentative rumen microbes to breakdown the feed for nutrient supply. This agrees with the report of Adewumi and Kazeem (2010) and Okon *et al.* (2018).

Table 1: Means±SEM of milk intake, body weight and linear body measurement as affected by lactation stages

Week	MI(g)	BW(kg)	NC(cm)	NL(cm)	BL(cm)	BC(cm)	GC(cm)	HW(cm)	TL(cm)
1	1.60±0.10 ^a	6.88±0.91 ^a	19.09±2.00	10.56±0.43 ^c	25.60±0.87 ^c	33.50±1.15 ^c	32.94±0.86 ^{cd}	32.80±1.32 ^c	21.03±1.22
2	1.53±0.18 ^{ab}	6.80±0.96 ^a	19.99±2.14	11.11±0.44 ^c	26.44±0.93 ^{bc}	34.04±1.15 ^{bc}	33.71±0.89 ^c	33.49±1.41 ^{bc}	21.67±1.22
3	1.44±0.14 ^{ab}	6.69±1.01 ^a	20.54±2.20	11.73±0.36 ^c	27.09±0.92 ^{abc}	34.89±1.06 ^{abc}	34.30±0.87 ^{bc}	34.10±1.43 ^{bc}	22.19±1.20
4	1.53±0.13 ^{ab}	6.53±1.03 ^a	20.97±2.26	12.36±0.39 ^{ab}	27.53±0.99 ^{abc}	35.53±1.16 ^{abc}	34.95±0.99 ^{bc}	34.76±1.47 ^{abc}	22.61±1.14
5	1.04±0.10 ^c	4.31±0.69 ^{ab}	21.14±2.38	12.79±0.46 ^{ab}	27.97±1.01 ^{ab}	36.20±1.32 ^{ab}	35.33±1.19 ^{abc}	35.36±1.50 ^{abc}	23.10±1.29
6	1.21±0.11 ^{bc}	4.26±0.93 ^{ab}	21.33±2.42	13.14±0.43 ^{ab}	28.44±1.05 ^{ab}	36.40±1.55 ^{ab}	35.56±1.24 ^{abc}	35.80±1.48 ^{abc}	23.44±1.24
7	1.20±0.10 ^{bc}	4.19±1.35 ^{ab}	21.53±2.45	13.57±0.40 ^{ab}	28.80±1.06 ^{ab}	36.91±1.65 ^{ab}	35.97±1.21 ^{ab}	36.19±1.49 ^{abc}	23.69±1.25
8	0.94±0.05 ^c	4.05±1.60 ^{ba}	21.91±2.47	13.99±0.44 ^{ab}	29.11±1.06 ^{ab}	37.20±1.64 ^{ab}	36.47±1.18 ^{ab}	36.64±1.48 ^{abc}	23.93±1.28
9	0.97±0.68 ^c	4.33±1.21 ^{ab}	22.34±2.51	14.36±0.48 ^{ab}	29.40±1.06 ^{ab}	37.49±1.63 ^{ab}	36.93±1.17 ^{ab}	37.30±1.36 ^{ab}	24.17±1.30
10	0.97±0.68 ^c	4.56±1.00 ^{ab}	22.79±2.57	14.71±0.50 ^{ab}	29.69±1.05 ^a	37.86±1.58 ^a	37.33±1.16 ^{ab}	37.67±1.40 ^{ab}	24.41±1.35
11	0.97±0.61 ^c	4.68±0.98 ^{ab}	23.26±2.66	15.11±0.55 ^a	29.99±1.08 ^a	38.27±1.56 ^a	38.53±1.87 ^a	38.00±1.43 ^a	26.10±2.08
12	0.96±0.08 ^c	4.89±0.59 ^{ab}	23.59±2.70	15.36±0.59 ^a	30.31±1.14 ^a	38.67±1.58 ^a	38.04±1.20 ^a	38.35±1.45 ^a	24.81±1.40
Overall mean	1.20±0.40	5.18±1.02	21.54±0.66	13.23±0.21	28.36±0.31	36.41±0.43	35.84±0.37	35.87±0.43	23.43±0.39

^{abcd} means with different superscript within columns are significantly different (P<0.05)

MI=milk intake, BW=body weight, NC=neck circumference, NL=neck length, BL=body length, BC=body circumference, GC=girth circumference, HW=height at wither, TL=tail length

Table 2: Pearson correlation of body weight and linear body measurements in white Fulani calves

	MI	BW	NC	NL	BL	BC	GC	HW	TL
MI	1	-0.203	0.285	-0.328	-0.170	-0.056	-0.112	-0.040	-0.023
BW		1	0.822**	0.877**	0.976**	0.938**	0.914**	0.952**	0.872**
NC			1	0.667**	0.835**	0.839**	0.805**	0.891**	0.813**
NL				1	0.803**	0.756**	0.755**	0.805**	0.748**
BL					1	0.928**	0.912**	0.954**	0.853**
BC						1	0.880**	0.892**	0.822**
GC							1	0.896**	0.888**
HW								1	0.926**
TL									1

**correlation coefficient is significantly high (P<0.05)

MI=milk intake, BW=body weight, NC=neck circumference, NL=neck length, BL=body length, BC=body circumference, GC=girth circumference, HW=height at wither, TL=tail length

Correlation of milk intake and body weight with linear body measurements

Table 2: Estimate of variables in linear function for milk intake, body weight and linear body measurement parameters of white Fulani calves

Independent	Dependent	Equation	R	R ²	SEM
Milk intake	BW	$Y=70.56-8.32x$	0.58**	0.47	6.62
Milk intake	NC	$Y=-15.83+4.77x$	0.83**	0.69	5.85
Milk intake	NL	$Y=-15.28+1.71x$	0.75**	0.56	1.79
Milk intake	BL	$Y=29.98-1.35x$	0.82**	0.66	2.86
Milk intake	BC	$Y=37.13-0.59x$	0.85**	0.82	3.92
Milk intake	GC	$Y=37.08-1.03x$	0.96**	0.92	3.35
Milk intake	HW	$Y=36.40-0.44x$	0.95**	0.90	3.97
Milk intake	TL	$Y=23.70-0.23x$	0.83**	0.75	3.62

**correlation coefficient is significantly high ($P<0.05$)

MI=milk intake, BW=body weight, NC=neck circumference, NL=neck length, BL=body length, BC=body circumference, GC=girth circumference, HW=height at wither, TL=tail length

Table 2 presents the coefficient of correlation between BW and LBM. Correlation between all body pairs of measurements were high, positive and highly significant ($P<0.05$). The highest correlation was recorded for BL, BC, GC and HW. This high correlation suggests that frame size and absolute height were complementary and that the total size of the animal is a function of length, height and circumference measurements. This result was in line with the findings of Adewumi and Kazeem (2010), Okon *et al.* (2018). Alade *et al.* (2008) and Cam *et al.* (2010) also reported high phenotypic correlation between BW and LBM in their research with goats and yarayaka sheep, respectively. There was however a negative and non-significant ($P>0.05$) correlation between MI and LBM parameters with the highest correlation recorded between MI and NC (0.285). This agreed with the findings of Adewumi *et al.* (2006) and Adewumi and Kazeem (2010) in their research on lactating ewes and yankasa lambs, respectively. This implied that as milk intake increases, LBM parameters tend to decrease though at a very slow rate. Milk intake was low but positively correlated with body weight (0.203).

Table 3 presents estimate of parameters in linear function fitted for milk intake, weight and linear body measurements. The

implication of the above is that the equation could be used to predict body weight effectively and accurately. The MI, NL and NC associated negatively but significantly ($P<0.05$) which implies that for every increment in milk intake, NC and NL tends to decrease though at a slow rate. This result agreed with the findings of Adewumi and Kazeem (2010) on a low NL and NC association with MI. Coefficient of determination R^2 values were high for all LBM indices assessed, with the highest R^2 value recorded for HW and GC. In this study, the equations show that BW as affected by MI is dependent on the growth of other component parts of white Fulani calves. This is in line with the findings of Alabi *et al.* (2012) that body weight is highly dependent on the growth of its component parts in three indigenous chicken breeds.

Conclusion and recommendation

From the findings, it can be concluded that milk intake increased between the early stages of the animal's growth and thereafter decreased as lactation progressed but all linear body measurements increased as the animal advanced in age. Correlation between all body pairs of measurements were high, positive and highly significant with the highest correlation recorded for BL, BC, GC and HW. Linear function

provided the best fit for milk intake, body weight and linear body measurements. It is recommended that any of the body measurement parameters could be used in the estimation of the body weight. Also, an improvement of non-genetic factor such as nutrition and general husbandry of the animal will bring about improvement in the body parameters.

References

- Adewumi, O. O. and Kazeem, G. 2011.** The relationship between milk intake, body weight and linear body measurement of Yanaksa lambs. Proceedings of the 15th Annual Conference of Animal Science Association of Nigeria, 15:163-166.
- Adewumi, O. O., Alokun, J. A. and Ologun, A. G. 2006.** Phenotypic variation and correlation among milk yield and physical body characteristics of yankasa and yankasa X WAD ewes. *African J. of science*. 5:39-43.
- Afolayan, R. A. and Lakpini, C. A. M. 2006.** Prediction of live weight from objective live-dimensional traits in Yankasa sheep. Proceedings of the 31st Annual conference of Nigerian society for Animal production. March 12th – 15th 2006, Bayero University, Kano, Nigeria
- Alabi, O. J., Ng'ambi, J. W., Norris, D. and Egena, S. S. A. 2012.** Comparative study of three indigenous chicken breeds of South Africa: Body weight and linear body measurements. *Agricultural Journal* 7(3):220-225.
- Alade, N. K., Rajiand, A. O. and Atiku, M. A. 2008.** Determination of appropriate model for the estimation of body weights in goats. *ARPJ. of Agric. Bio. Sci.* 3(4): 52-57.
- Cam, M. A., Olfaz, M. and Soydan, E. 2010.** Body measurements reflect body weights and carcass yields in Karayaka sheep. *Asian Journal Animal and Veterinary Advances*. 5: 120-127.
- Dim, N. I., Egahi, J. O. and Tanko, M. U. 2012.** Zoometrical measurements of body parts and their Relationship with live weight in bunaji cows in Zaria Nigeria. *I.J.A.B.R.*, VOL. 2(2):339-341
- Ladan, S. A., Egahi, J. O. and Dim, N. I. 2009.** Body biometry as a predictive index for body weights in Yankasa ewes under traditional husbandry system in North Central Nigeria. Proceedings of the 34th Annual Conference of the Nigerian Society for Annual Production, March 15-18, 2009, university of Uyo, Uyo, Nigeria.
- Okon, U. M., Umoh, B. I. and Ekpo, J. S. 2018.** The Relationship Between Body Weight and Linear Body Measurements in Bunaji Calves at Different Phases of Growth. *Aksu J. of Agric. and food sciences*. 2(1) 189-196.
- Ortheruata, A. M. and Olutogun, O. 1994.** Pre-and post-weaning phenotypic relationship between some N'dama cattle linear measurement in the tropics. *Niger. J. of Anim. Prod.*, 21: 76-82.
- Pandey, G. S. and Voskuil, G. C. S. 2011.** *Manual on Milk safety, quality and hygiene*. Golden Valley Agricultural Research Trust, Zambia. Pp 52.
- Salako, A. E. 2006.** Principal component factor Anaylsis of the morphological structure of immature uda sheep. *International Journal of morphology*, 24 (4): 571 – 774.
- Sluk, A. K. 1989.** Soil and land use studies. Government print office, Uyo, Akwa Ibom State. Soil survey staff, 1994. Keys to soil taxonomy. Soil management support services

Correlation of milk intake and body weight with linear body measurements

(SMSS), technical monogr. No.
19:306p.

**Younas, U., Abdullah, M., Bhatti, J. A.,
Pasha, T. N., Amhed, N., Nasir, M.
and Hussain, A. 2013.**
Interrelationship of body weight
with linear body measurements in
Hissardale sheep at different stages
of life. *Journal of Animal and Plant
Sciences*. 23(1):40-44.

Received: 5th November, 2019

Accepted: 19th February, 2020