Growth, nutritive value and dry matter degradability of three tephrosia species

O.J. Babayemi¹, M.A. Bamikoke², I.O. Daniel³, A.Ogungbesan¹, A. Babatunde¹

¹Department of Animal Science, University of Ibadan, Ibadan, Nigeria.

²Department of Animal Science, University of Benin, Benin City, Nigeria.

³Department of Plant Breeding and Seed Technology, ⁴University of Agriculture, Abeokuta, Nigeria

Abstract

Two trials were carried out to evaluate the potential of *Tephrosia bracteolata* (TB), *Tephrosia candida* (TC) and *Tephrosia linearis* (TL) for animal feed. In an 18 week growth experiment, *Tephrosia* species differed ($P<0.05$) in leaf number (LN) and stem height (SH). TB and TL had apparent ($P<0.05$) early and persistent growth than TC until weeks 12-18 when they retarded in growth at the commencement of flowering. TC however, increased geometrically in LN and SH. There were significant ($P<0.50$) differences among the species in dry matter (DM) and acid detergent fibre (ADF). The DM, CP, NDF, ADF, EE and ash range from 30-41.8, 24-26.5, 62.5-68.5, 45.8-48.4, 2.3-26 and 5.6-6.2% respectively. Calcium and sodium were highest ($P<0.05$) in TB and TL species were not significantly ($P>0.05$) different in degradation characteristics. Lower amount of crude protein (10-14.81%) was released in the 3 *Tephrosia* species. As a result of rapid growth, high nutritive values and easily degradable nutrients of *Tephrosia* shrubs, ruminants could benefit immensely from the plants.

Keywords: Nutrient, utilization, dry matter, degradation, *Tephrosia*, species

Introduction

Tephrosia plants are tropical shrub legumes with over 300 species distributed worldwide. They are notably dispersed throughout the tropics, sub-tropical Australia and North America (Dutta, 1979). One of the species, *Tephrosia candida* (TC) has been enlisted with suitable multipurpose trees (MPTS) and shrub species for alley farming systems in humid and sub-humid zones of Africa (Daniel, 1971, Bantsh and Paul, 1992).

In Nigeria, among the commonest species found in wild of the drier part of the west and

middle belt zones are *Tephrosia bracteolata* (TB), *Tephrosia linearis* (TL), *Tephrosia vogelli* (TV), *Tephrosia candida* (TC). However, TB and TL occur more frequently than other species. It was reported that some of these legume trees are used locally as firewood in Cameroon (Fontunyan and Mbeni, 1987).

The tephrosia species especially TB and TL are locally called "roro" in the western part of Nigeria and are highly valued by rural dwellers. The local farmers especially old women are preoccupied with it by adventuring into the bush to cut the shrubs for goats and sheep

feeding while excess of their need is marketed on daily basis. An unreported observation showed that the ubiquitous goat and sheep reared on extensive system go to mountaintops and valleys to browse on the legume.

Apparent digestibility has been reported (Fomunyan and Mbomi, 1987) for TV and suggested that its crude protein value can be utilized like cotton seed cake for protein supplementation. Some *Tephrosia* legumes were also established to be relishing to cattle (IBPGR, 1984) and goats (Ayoade *et al.*; 1998). There is limited documented information on *Tephrosia* species pertaining to growth, and nutritive value. Grazing animals will benefit little from a weak, too short or too tall legume trees/shrubs. The reports (Caren *et al.* 1987; Addlestone *et al.*, 1999) showed that successful stand establishment of forage is highly dependent on seedling vigour as measured by seedling height, stem diameter and herbage yield. The present study was undertaken to determine the growth and nutritive value of three *tephrosia* species.

Materials and methods

Site and land preparation

The growth experiment was conducted at International Institute for Tropical Agriculture (IITA), Ibadan, and rumen degradability study at International Livestock Research Institute (ILRI), Ibadan, (7°30'N, 3°54'E), 200m above sea level. The study was carried out late dry season (March) by irrigation and early rainy season (July), 2000. Existing shrubs were removed from the site for the nursery to disallow shading and competition with *Tephrosia* seedlings. Soil samples were collected using an auger to a depth of 15cm from the earmarked area of 50 by 50m. The sampled soils being a composite of five cores were spread in the laboratory for air-drying. The soil pH and nitrogen were determined while exchangeable K, Na, Ca and Mg were

extracted as described (Black, 1965). A polythene sheet was spread out upon which the soil filled polythene pots were placed. Polythene pots were watered to saturation for three consecutive days until water was observed to seep out from the drainage holes of the pots.

Seed treatment and planting

The seeds of the *tephrosia* species used were obtained from IITA, Ibadan and were exposed to pre-treatment culture for 30 seconds in boiled water as established (Babatunde, 2000). The treated seeds were counted and divided in a completely randomized design into five batches of 50 seeds per species. Each batch of 50 seeds was further sub-divided into five of 10 seeds sown into the pots at the depth of 2mm.

Plant height and leaf number

The plant height (PH) and leaf number (LN) of the shrubs at 20% minimum survival were taken at 4 weeks post planting. The subsequent measurements and counting were effected at 2 weeks interval for upward of 18 weeks.

Sample collection and dry matter determination

Samples were collected at 12 weeks old. Leaves were separated from the stem and the former were freshly weighed and dried in a drought oven at 65°C for 48 hours to a constant weight for dry matter (DM) determination. The dried samples were then milled in a micro-miller using a 1-mm diameter screen and 2.5mm sieve for rumen degradation.

Rumen degradability

In order to evaluate DM disappearance, the samples that were previously oven dried were then incubated in the rumen of fistulated sheep using the nylon bag technique (Orskov and McDonald, 1979). Three grams of the milled (2mm screen) samples were weighed into nylon bags measuring 140 x 75mm with pore size of 4mm. The bags were incubated in the

Degradation characteristics and nutritive value of *Tephrosia* species

rumen of the sheep in duplicates for 24, 48, 72 and 96h. The evacuated bags from the rumen were rinsed immediately under a running tap to terminate fermentations. The residues in the bags were oven dried at 65°C for 48 Hr and the DM was as sample weight loss. The results for DM disappearance were fitted to the exponential equation $P = a + b(1 - e^{-ct})$ as reported (Qrskov and McDonald, 1979) where a = water-soluble fraction; b = insoluble but degradable fraction; c = rate of degradation and p = the degradation at time t .

Chemical analysis

The dry matter (DM), crude protein, ether extract and ash of the samples were determined according to AOAC (1990). The neutral detergent fibre (NDF) and acid detergent fibre (ADF) were analysed as reported (Goering and Van Soest, 1970). Mineral elements were determined by atomic absorption spectrophotometer (AAS) model 490 Gallenkamp, London.

Statistical analysis

Data generated were subjected to analysis of variance using the general linear model procedure of statistical analysis (SAS, 1995). Statistical differences among treatment means were declared at 5% level. Means were separated using Duncan (1955) multiple range test.

Results and discussion

Physical and chemical characteristics of soil for trial one are shown in Table 1. Although much have not been established but scanty information showed that tephrosia plants could do well in a slightly acidic soil (AFNETA, 1992). The N level of the soil (0.66%) obtained for the present study is much higher than 0.07 - 0.11% reported else where (Tening et al., 1994) for the establishment of *Stylosanthes*.

The average leaf number (LN) and stem height (SL) are presented in Tables 2 and 3 respectively. There was a rapid increase in both LN and SH from weeks 4-8 and then followed by an increase at a decreasing production in weeks 10-18 for the legumes TB and TL. It was observed that TC improved ($P < 0.05$) in LN and SH for weeks 10-18 than either TB or TL. *Tephrosia candida* is suspected to be a perennial plant as it outgrew other species in week 18 and remained green till the end of the dry season. Although leaf: stem ratio was not determined, optimum production of leaf falls between 12-16 week of growth. Higher leaf content is desirable for livestock production. It was reported that the nutrient composition in leaf materials of tropical legumes is higher than that the stem or mixture of it (ILCA, 1994). In a grazing trial, Hamilton (1970) stressed the role of green leaf content of a pasture demonstrated and discovered that milk, meat and wool production deteriorated significantly with low leaf content of the pasture. The multiple productions of leaflets exhibited by the three shrub legumes of our study are similar to some reports (Tarawali *et al.*, 1999; Addlestone *et al.*, 1999) for tropical legumes.

The nutritive values of the shrubs are in Table 4. *Tephrosia candida* had a higher DM and NDF than what TB and TL contained. However, the composition of ADF in TB and TL is more apparent ($P < 0.05$) than that of TC. Crude protein, ether extract and ash constituents for the three-tephrosia species are similar ($P > 0.05$). The level of crude protein (24-26.5%) in the present study is higher than 22.8% reported (Ayoade *et al.*, 1998) and 15.8% established elsewhere (Le Houerou, 1980). Crude protein for *Tephrosia purpurea* (15.5%), *Cajanus cajan* (18.6%) was much

Table 1 Physical and chemical properties of soil for *Tephrosia* species

Measurement	values
PH (H ₂ O) (1:1)	5.6
Nitrogen (%)	0.66
Phosphorus (µg/g)	28.22
Calcium (meg/100g)	2.52
Sodium (meg/100g)	0.61
Magnesium (meg/100g)	0.75
Potassium (meg/100g)	0.50
Sand (%)	83.60
Silt (%)	10.00
Clay (%)	6.40

lower than those of the present study but are favourably compared with 25% *Gliricidia sepium* and 25.3% *Leucaena leucocephala* (Little *et al.*, 1989). Higher protein could be obtained in tephrosia species if harvested at a much lower age as the age of harvest influence nutrient composition (Norton, 1994). Neutral detergent fibre was observed to be significantly higher in TC than values in TB and TL. The reason for the difference is not known but the textural leaves of TB and TL are smoother and tender than that of TC. The NDF values are however, within the range of 25-71% reported for some valuable multipurpose trees in Nigeria (Bayer, 1990; Larbi *et al.* 1993). High ADF content were obtained in the legumes, which is comparable to 49.2% for *Ficus capensis* but higher than 31.0% for *Alchornea cordifolia* as reported (Akinlade *et al.*, 2001).

Table 5 shows the calcium, phosphorus, magnesium, potassium and sodium composition of the forages. There were significant ($P < 0.05$) differences among the shrubs in Ca and Na. *Tephrosia* bract and TL are apparently higher in Ca and Na than TC. No significant variation was noticed in P, Mg and K among the legumes. The amount of minerals in the study is comparable to those of Indonesian forages (Little *et al.*, 1989). Without mineral fortification in the diets, the amounts present in the three forages are high enough to meet the requirement for ruminant animals (Akinsoyinu and Akinyele, 1979; Afolabi, 1995; Babayemi *et al.*, 1999).

Rumen degradability characteristics of the three-tephrosia species are in table 6. In all the plant species, no differences were observed in DM losses (48h). The degradability increased (figure 2) as the incubation period increases (Adama and Lapeyronie, 1997). The values (85.7 - 86.2%) in the present study are higher than reported for *Ficus capensis* and *Alchornea cordifolia* (Akinlade *et al.*, 2001) and of *Chromolaena odorata* (Apori *et al.*, 2000). The DM release is also higher than 25.44 - 62.93% reported (Keyserlingk *et al.*, 1996) for alfalfa. The rumen degradable protein in this study is low (Figure 1) when compared with CP degradation of 41% guinea grass, 60% ruzi,

Table 2 Mean leaf number of the *Tephrosia* species

Species	Stage of growth (Week)							
	4	6	8	10	12	14	16	18
<i>Tephrosia bracteolata</i>	16 ^a	27 ^a	48 ^a	66 ^a	83 ^a	88 ^a	90	90
<i>Tephrosia candida</i>	11 ^b	23 ^b	37 ^c	48 ^c	65 ^b	74 ^b	86	98
<i>Tephrosia linearis</i>	12 ^b	25 ^a	41 ^b	56 ^b	81 ^a	87 ^a	91	90
SEM	0.76	0.58	1.16	2.60	2.85	2.26	0.76	1.33

^{a,b,c} Means on the same column with similar superscripts are not significantly ($P > 0.05$) different

Degradation characteristics and nutritive value of *Tephrosia* species

Table 3 Mean stem height (cm) of the *tephrosia* species

Species	Stage of growth (Week)							
	4	6	8	10	12	14	16	18
<i>Tephrosia bracteolata</i>	22.5 ^a	32.2 ^a	53.9 ^a	70.6 ^a	88.9 ^a	90.2 ^a	93.0 ^a	94
<i>Tephrosia candida</i>	14.7 ^b	21.4 ^c	33.3 ^c	45.1 ^c	54.8 ^b	58.1 ^b	68.5 ^b	82
<i>Tephrosia linearis</i>	20.7 ^a	24.2 ^b	42.3 ^b	63.9 ^b	88.4 ^a	88.3 ^a	91.6 ^a	92
SEM	1.18	1.62	2.98	3.82	5.64	5.20	3.97	1.86

^{a,b,c}Means on the same column with similar superscripts are not significantly ($P>0.05$)

Table 4 Nutrient composition of the leaf of *Tephrosia* species

Species	Composition (%)					
	DM	CP	NDF	ADF	EE	ASH
<i>Tephrosia bracteolata</i>	31.9 ^b	24.3	63.2 ^b	48.4	2.4	5.8
<i>Tephrosia candida</i>	41.8 ^a	26.5	68.5 ^a	45.8	2.6	6.2
<i>Tephrosia linearis</i>	30.1 ^b	24.0	62.3 ^b	47.9	2.3	5.6
SEM	1.82	0.39	0.97	0.39	0.004	0.008

^{a,b,c}Means on the same column with different superscripts differ ($P<0.05$) significantly.

69% leucaena and 79% gliricidia (Ibrahim *et al.*, 1995). Such low CP release could indicate un-degradable intake protein (UIP). According to Bohnert *et al.*, (1998), greater N degradation per hour resulted in less ruminal escape N. *Tephrosia* species could then be used as UIP. As reported elsewhere (Zinn *et al.*, 1981; Titgemeyer *et al.*, 1989), that suppling of ruminants with UIP can increase the flow of N and amino-acids to the small intestine. It will also result in appreciable growth and efficiency of N utilization (Stock *et al.*, 1981; Goedeken *et al.*, 1990).

Conclusion

Tephrosia species are shrub legumes, early maturing and possess high nutritive values. Dry matter degradability is high with a low release of crude protein. The implication of this is that the use of *Tephrosia bracteolata*, *candida* and *linearis* would not only enrich the soil but also improve the performance of livestock when fed. Further work needs to be carried out on the factors that could limit their ingestion and utilization as various tropical browse plants contain some anti-nutritional factors.

Table 5 Some mineral composition of the leaf of *Tephrosia* species

Species	Ca	P	Mg	K	Na (ppm)
<i>Tephrosia bracteolata</i>	1.52 ^a	0.27	0.64	1.23	269.1 ^a
<i>Tephrosia candida</i>	1.26 ^b	0.29	0.58	1.31	140.9 ^b
<i>Tephrosia linearis</i>	1.54 ^a	0.25	0.61	1.24	260.3 ^a
SEM	0.05	0.006	0.009	0.012	20.67

^{a,b,c}Means on the same row with similar superscripts are not significantly ($P<0.05$) different.

Table 6 Dry matter degradability characteristics of the leaf of *Tephrosia* species in rumen of WAD sheep

Measurement	TB	TC	TL	SEM
a (%)	37.5	36.9	37.4	0.093
B	48.2	49.3	48.6	0.161
P	85.7	86.2	86.0	0.073
c (%h ⁻¹)	3.27	3.80	3.21	0.094

a = soluble fraction, b = degradable fraction, p = potentially degradable fraction, c = rate of degradation of b.

References

- Adllestone, B.J. J.P. Mueller and J.M. Luginbuhl 1999. The establishment and early growth of three, leguminous tree species for use in silvopastoral systems of the southern USA. *Agroforestry Systems* 44: 253-275.
- Afnet, 1992. Alley farming network for tropical Africa. Multipurpose Tree Screening and Evaluation. *Alley Farming Training manual* vol. 1:1-17.
- Afolabi, G.O. 1995. Calcium and Phosphorus utilization during pregnancy and lactation by West African dwarf goats. A Ph.D. Thesis, University of Ibadan.
- Akinlade J., J.A. Olanite and M.A. Bamikole 2001. Dry matter degradation characteristics of rice stover with different proportions of *Ficus capensis* or *Alchornea cordifolia* in rumen fistulated sheep, goats or cattle. *Niger. J. Anim. Prod.* 28(2): 174-181.
- Akinsoyinu, A.O. and I.O. Akinyele 1979: Major elements in milk of the West African dwarf goats as affected by stage of lactation *J. Dairy Res.* 46:427-431. Analytical Methods for feeds, animal, excrements and animal tissues (1997): ILRI Nutritional Laboratory, ILRI - Ethiopia Addis - Ababa pg. 15.
- AOAC (Association of Official Analytical Chemistry) 1990. *Official methods of analysis 15th Edition* AOAC Incorporation, Virginia 22201 USA.
- Apori, S.O., R.J. Long, F.B. Castro and E.R. Qrskov 2000. Chemical composition and nutritive value of leaves and stems of tropical weed *Chromolaena odorata*. *Grass and forage Science* 55, 77-81.
- Ayoade, J.A., P.O. Ogebe, A.J., Okwori and J.O. Ogbeide 1988. Nutritional potential of *Tephrosia bracteolata* leaves as a sole feed for goat. In: O.O. Oduguwa, A.O. Fanimo, and O.A. Osinowo (eds.). *Animal Agriculture in West Africa. The sustainability question. Proceedings of the Silver Anniversary Conference of the Nigeria Society Animal Production* 21-26 March 1998 pp. 53-54.
- Babatunde, O.O. 2000. Preliminary studies on seed germination, seedling vigour and rumen degradation of *Tephrosia* species. *M.Sc. Project, University of Ibadan, Nigeria*.
- Babayemi, O.J., A.O. Akinsoyinu, O.A. Isah, A.A. Adeloye, M.A. Bamikole and M.K. Adewunmi 1999. Effect of Magnesium supplement on performance characteristics of lactating West African dwarf goats. *Trop. Anim. Prod. Invest.* 2: 61-68.
- Bansh, R.T. and J.P. Paul 1992. The AFNETA Alley farming training manual vol. 1: Alley Farming Network for Africa Pg. MPTs 2-12.

Degradation characteristics and nutritive value of Tephrosia species

- Bayer, W.** 1990. Use of native browse by Fulani cattle in central Nigeria. *Agroforestry Systems* 12:217-228.
- Black, C.A.** 1965. Methods of soil analysis, Agronomy number 9(2), American Society of Agronomy, Madison, Wisconsin.
- Hohnert, D.W., B.T. Larson, M.L. Bauer, A.F. Branco, K.R. Mcleod, D.L. Harmon and G.E. Mitchell, Jr.** 1998, Nutritional Evaluation of Poultry By-Product Meal as a Protein Source for Ruminants: Effects on Performance and nutrient Flow and Disappearance in Steers. *J. Anim. Sci.* 76:2474-2484.
- Caren, C.J., A.M. Milson, R.L. Gang and G.L. Thor** 1987. II Seed emergency in marginal soil moisture *J. Range Manage* 40:212-216.
- Daniel, F.R.** 1871. Flora of Tropical Africa volume II legumensac to filicoideae by Reeve L and Co 5 Henrietta street, Convent Garden.
- Duncan, F.R.** 1871 Flora of Tropical Africa volume II legumensiosac to filicoideae by Reeve L and Co 5 Henrietta street, Convent Garden.
- Duncan, D.B.** 1955. Multiple range and multiple f- tests - *Biometrics* 11: 1-42.
- Dutta, A.C.** 1979. *Botany for degree student*. Calcutta, Oxford University Press. Delhi ed. Pp. 728.
- Fomunyan, R.T. and S.E. Mboni** 1989. Tephrosia species and cotton seed cake supplementation of rice and maize stalks fed to sheep and goats in the dry season. In: overcoming constraints to efficient utilization of agricultural by-products of animal feed. *A proceeding of the 4th annual workshop held at the institute of animal and research Marikon station, Bamenda, Cameroon* 20-27, October, 1987.
- Goedeken, F.K., T.J. Klopfenstein, R.A. Stock, R.A. Britton, and M.H. Sinat** 1990. Protein value of feather meal for ruminants as affected by blood additions. *J. Anim. Sci.* 68:2936-2944.
- Goering, K. and Van Soest P.J.** 1970: Forage fibre analysis. Apparatus, reagents, procedures and some applications. *Agricultural Handbook*, 379, 1-20.
- Hamilton, R.I. Lambourne, L.J. Roe R. and Minson, D.J.** 1970. Quality of Tropical Pastures for Milk production in: *Proceedings of the XI International Grassland Congress*. pp. 860-864.
- Ibrahim, M.N.S. Tamminga and G. Zemelink** 1995. Degradation of tropical roughages and concentrate feeds in the rumen. *Anim. Feed. Sci. Tech.* 54:81-92.
- ILCA** 1994. Improving livestock production in Africa. Evaluation of ILCA's programme 1974-1994 pp. 150.
- International Board for Plant Genetic Resources (IBPGR)** 1984. Forage and browse plants for Arid and Semi Arid African royal Botanical Garden Kew pp. 235.
- Keyserlingk, M.A., G. Swift, M.C. Puchala and R. Shelford** 1996. Degradation of dry matter and crude protein of forages in ruminants. *Anim. Feed Sci. Tech.* 57:291-311.
- Larbi, A., D. Thomas and J. Hanson** 1993. Forage potential of Erythrina Abyssinica: intake, digestibility and growth rates for stalled sheep and goats in southern Ethiopia. *Agroforestry Systems* 21:263-270
- Le Houerou, H.N.** 1980. Chemical Composition and nutritive value of browse in West Africa in Le Houerou H.N (eds.), *Browse in Africa* ILCA Addis Ababa Ethiopia pg. 329-338.

- Little, D.A., S. Kompiang, and Petheram, R.J. 1989. Mineral composition of Indonesian forages *Tropical Agriculture* (Trinidad) 66:33-37.
- National Academy of Science 1979. *Tropical Legumes: Resources for the future*. Washington DC.
- Norton, B.W. 1994. Tree Legumes as dietary supplements for Ruminants. In R.C. Cutteridge and H.M. Shelton (eds.) *Forage legume in tropical Agriculture* (CAB International University Press Cambridge UK pp. 193-202.
- Orskov E.R. and McDonald I. 1979. The estimation of protein degradability in the rumen from incubation measurements according to rate of passage *J. Agric. Sci. Cambridge* 92:449-503.
- Statistical Analysis System Institute 1995. *SAS User's Guide*. SAS Institute, Cary Stock, R.N. Merchen. T. Klopfenstein, and M. Poos (1981). Feeding values of slowly degraded proteins, *J. Anim. Sci.* 53:1109-1119.
- Tarawali, S.A., Peters, M. and Scheullze Kraft 1999 Forage legumes for sustainable agriculture and livestock in sub-humid W/Africa, ILRI Project report ILRI Nairabi Kenya pp. 25-30.
- Tening, A.S., G. Tarawali, K.B. Adeoye, M.A. Mohammed - Saleem and J.A.I. Omuetti 1994. Management and nutrient requirements of Stylosanthes in pasture and cropping systems in the humid zone of Nigeria: A review, P.N. deLeeuw, M.A. Mohamed-Saleem and A.M. Nyamu (eds.) *proceedings of the Regional Workshop on the Use of Stylosanthes in West Africa held in Kaduna, Nigeria, 26-31 October, 1992*, pp. 113-1222.
- Titgemeyer, E.C., N.R. Merchen, and L.L.Berger 1989. Evaluation of Soybean meal, corn gluten meal, blood meal and fish meal as sources of nitrogen and amino acids disappearing from the small intestine of steers. *J. Anim. Sci.* 67:262-275.
- Zinn, R.A., L.S. Bull, and R.W. Hemken 1981: Degradation of supplemental proteins in the rumen. *J. Anim. Sci.* 52:857-866.

(Received 19 April 2001; Accepted 16 October 2001)