Assessment of production objective and breeding practices of rural goat keepers and implications for a breeding programme in north central Nigeria

*Yakubu, A. and Achapu, M. M.

Department of Animal Science, Faculty of Agriculture, Nasarawa State University, Keffi, Shabu-Lafia Campus, P.M.B. 135, Lafia, Nasarawa State, Nigeria.

*Corresponding author's email address: abdulmojyak@gmail.com

Abstract

Goat farming is a veritable source of livelihood of many rural families in Africa. This study aimed at determining prevailing production systems and breeding objectives of rural goat producers in north central Nigeria. A total of 180 rural goat keepers corresponding to 60 per State (Nasarawa, Benue and Plateau) were randomly sampled. Primary data (socio-economics of respondents, reasons for keeping goats, flock structure, management system, productivity and breeding practices) were collected through individual structured questionnaire administration. Cross tabulations and Chi square (χ^2) statistics were used to compare categorical variables, while rank means, arithmetic means and standard deviations were calculated for within- and between-state comparisons of the continuous variables. While more goat producers were involved in crop farming in Benue State (43.6%), only 34.5 and 21.8% engaged in farming in Plateau and Nasarawa State, respectively. Goats were kept for income generation, milk, meat and cultural/religious functions by about 61.1, 12.8, 15.0 and 6.1% of the producers while the relative importance given by respondents to the different objectives varied significantly (Chi-square=6.62; $P < 0.05$) across the States. The average flock sizes of goats for Nasarawa (9.68 ± 5.63), Benue (8.25 ± 4.73) and Plateau (8.80 ± 1.98) were not significantly ($P > 0.05$) different. Semi-intensive system predominated ($P < 0.01$). Productivity indices showed that for age of parturition, number of kids of Sahel doe and lifespan of goats, there was no significant difference ($P > 0.05$). Among all the breeding traits across the three States, only disease resistance varied ($P < 0.01$). Disease resistance, survival, fertility, number of offspring and body size appeared similar ($P > 0.05$) as preference for production traits. However, growth ($83.52-97.68$ mean ranks) (Plateau State) and cultural importance ($75.28-104.70$ mean ranks) (Benue State) varied across the States ($P < 0.05$ and $P < 0.01$, respectively). The present information will be useful in understanding the farmers' production objectives, management and breeding practices as a first step in designing a sustainable breeding programme for rural farmers in the study areas.

Keywords: Goats, management, breeding, productivity, Nigeria

Introduction

Goat production over the years is one of the major means of improving the livelihoods of poor livestock keepers, reducing poverty and attaining sustainable agriculture and universal food security (Abdul Aziz, 2010; Yakubu *et al.*, 2010; 2011; Abegaz *et al.*, 2013). The population of goats in Nigeria is estimated at 53.8 million (FAOSTAT, 2011) of which traditionally reared stock contributes 99.97%, while 0.03% of the

stock is commercially managed.

However, goat production in north central of Nigeria is constrained by many biological, environmental and socio-economical factors. Among them, inadequacy/lack of systematic breeding programs is an important constraint. There is no systematic goat breeding program in place and goat is the most neglected livestock species in research and development endeavours (Tsegahun *et al.*,

Blood chemistry and carcass characteristics of finisher broiler chickens fed Malted Sorghum sprout (MSP) or Wheat-offal based diets supplemented with yeast culture and enzyme

¹*Oke, F. O., Oluwatosin, O. O., ² Obadire, S. O., ³ Jegede, V. A., ⁴ Fafiolu, A. O., Oso, A. O., ⁵ Adeyemi, A. O., Olorunisola, R. A., ⁶ Adeoye, S. A.

¹Department of Animal Science Federal University Dutse, Jigawa State.

²Department of Animal Nutrition Federal University of Agriculture, Abeokuta, Ogun state.

³Laboratory Department Federal Medical Centre Birnikudu Jigawa State.

⁴Faculty of Science, Ondo State University of Science and Technology, Ondo, Nigeria.

Corresponding author: florenceobadire@gmail.com

Abstract

This experiment was conducted to determine the blood chemistry and carcass characteristics of broilers fed malted sorghum sprouts (MSP) or wheat offal (W/O)-based diets supplemented with yeast culture and enzyme. A total of two hundred and forty 28-day old, Marshal Broiler chickens used for the experiment were randomly allotted to 8 dietary treatment groups of 30 birds each. Each treatment group was replicated thrice with 10 birds per replicate. Data on selected blood indices and carcass traits of the birds were collected during the course of the feeding trial that lasted for 4 weeks. The experiment was a 2X4 factorial consisting of two (2) test ingredients (MSP and wheat offal) at 4 inclusion levels of enzyme or yeast (0g kg⁻¹, +0.01gkg⁻¹ yeast and Roxazyme G2[™] enzyme, +0.01gkg⁻¹ yeast, + Roxazyme G2[™] enzyme. At the end of the trial, the effect of MSP and wheat offal inclusion showed a significant effect ($P < 0.05$) for uric acid, WBC, Total protein, albumin and globulin. However, uric acid, total protein, Albumin and Globulin were highest for wheat offal diet. Furthermore, the results show that values obtained for white blood cell, uric acid, total protein, albumin, globulin, creatinine, ALT & AST were significantly ($P < 0.05$) affected by the enzyme, yeast and their combination. The carcass results show that none of the parameters measured differed significantly. In conclusion, supplementation with yeast + enzyme improves total protein, albumin and globulin. Meanwhile, it reduced concentrations of uric acid, ALT and AST by broiler chickens. Therefore MSP diet should be supplemented with yeast + enzyme or enzyme singly to improve Livestock ration for better utilisation.

Keywords: Blood parameters, carcass, starter broiler, malted sorghum sprout, wheat offal

Introduction

The increase in cost of conventional feedstuffs for basal energy and protein in poultry feeding have encouraged inadequate concentrate feeding and low productivity in poultry production. However, it has also resulted in competition between man and livestock of the limited resources. This high competitive rate of demand for grains especially maize has

made it imperative to augment with other non- conventional feed ingredients. There has been an increase in the awareness of the usefulness of agro-industrial by-products as replacements for grains especially maize in livestock diets (Ukachukwu *et al.*, 2003). One of such by-products is malted Sorghum Sprouts (MSP). MSP a by-product of sorghum malting has been reported by Oduguwa *et al.* (2007) to have a

categorical variables within and between states, while rank means and arithmetic means and their standard deviations were calculated for within- and between-state comparisons of the continuous variables. Within each state, comparisons of mean ranks were performed using the Friedman test, which compares the distribution of preference ranks of each animal function and of each selection criterion. Post hoc analyses were then applied using the non-parametric Wilcoxon Signed-rank test with Bonferroni's adjustments. The non-parametric Kruskal-Wallis test followed by the Mann-Whitney U test for post hoc separation of group means and mean ranks was used for comparison between states. A statistically significant difference was reported if the p value is less than 0.05. SPSS (2010) statistical package was employed.

Results

Socio-economic characteristics of farmers

More males (37.4%) were found in Benue State while the female respondents were more in Nasarawa State (43.9%), although not significantly different ($P>0.05$) (Table 1). Majority of the households were married (92.22%) while more singles (61.5, 53.5 and 0.0%, respectively) were observed ($P<0.05$) in Benue and Nasarawa compared to Plateau State. The primary occupation of the households varied from one State to another ($P<0.001$). While more goat producers (43.6%) were involved in crop farming in Benue State, only 34.5 and 21.8% engaged in farming in Plateau and Nasarawa State, respectively. However, more households were involved in trading in Nasarawa (44.7%) compared to 15.8% in Benue State. Only 14, 9 and 16 individuals ($P>0.05$) had access to credit facilities in Nasarawa, Benue and Plateau States, respectively. The average age of respondents was significantly ($P<0.05$)

higher in Plateau (45.85 ± 6.11) than Benue and Nasarawa States (40.17 ± 9.38 and 38.28 ± 10.56 , respectively). There was no significant difference ($P>0.01$) in the average household size in the three States. However, households in Plateau State had significantly ($P<0.05$) more years of goat experience (9.06 ± 4.42) compared to their counterparts (7.22 ± 4.41 and 7.20 ± 4.53).

Reasons for goat keeping

119 (61.1), 23 (12.8), 27 (15.0) and 11 (6.1%) kept goats for income generation, milk, meat and cultural/religious functions, respectively in Nasarawa, Benue and Plateau States (Figure 1). The relative importance given by respondents to the different production objectives varied significantly (Chi-square=6.62; $P<0.05$) across the States.

Flock composition

The average flock size of goats for Nasarawa (9.68 ± 5.63), Benue (8.25 ± 4.73) and Plateau (8.80 ± 3.98) was not significantly ($P>0.05$) different from one another. Across the three States, West African dwarf goats were the most reared animals (mean flock size of 7.17 ± 4.76 , 6.93 ± 4.09 and 7.85 ± 3.95 for Nasarawa, Benue and Plateau States, respectively) while Sahel goats were the least kept (Table 2).

Foundation stock and systems of management of goats

Most of the foundation stock were purchased from the markets and neighbourhoods, and cut across ($P>0.05$) the three States. The management system varied across ($P<0.01$) the three States, where semi-intensive system predominated and was more practiced (39.3 and 34.8%, respectively) in Plateau and Nasarawa States. A good number (82.78%) of the households had access to veterinary personnel which varied ($P<0.01$) from location one to another (Table 3).

Table 1: Socio-economic characteristics of goat keepers in Nasarawa, Benue and Plateau States

Characteristics	Nasarawa No (%)	Benue No (%)	Plateau No (%)	Pearson Chi-square	Significance
Categorical variables					
Sex					
Male	35 (28.5)	46 (37.4)	42 (34.1)	4.78	0.09
Female	25 (43.9)	14 (24.6)	18 (31.6)		
Marital status					
Married	55 (31.1)	51 (30.7)	60 (36.1)	10.27	0.036*
Single	5 (35.5)	8 (61.5)	0 (0.0)		
Widowed	0 (0.0)	1 (100.0)	0 (0.00)		
Education					
None	26 (35.1)	22 (29.7)	26 (35.1)	2.93	0.82
Primary	23 (33.3)	24 (34.8)	22 (31.9)		
Secondary	10 (31.2)	13 (40.6)	9 (28.1)		
Tertiary	1 (20.0)	1 (20.0)	3 (60.0)		
Primary occupation					
Crop farming	12 (21.8)	24 (43.6)	19 (34.5)	28.49	0.000**
Trading	34 (44.7)	12 (15.8)	30 (39.5)		
Artisan	0 (0.0)	3 (100.0)	0 (0.0)		
Civil service	12 (31.6)	15 (39.5)	11 (28.9)		
Others	2 (25.0)	6 (75)	0 (0.0)		
Access to credit					
No	46 (32.6)	51 (36.2)	44 (31.2)	2.55	0.28
Yes	14 (35.9)	9 (23.1)	16 (41.0)		
Non categorical					
	Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD		
Age of respondent	38.28 $\pm$ 10.56 ^a	40.17 $\pm$ 9.39 ^b	45.85 $\pm$ 6.11 ^c		
Household size	7.08 $\pm$ 3.97 ^a	6.47 $\pm$ 8.58 ^a	6.98 $\pm$ 2.29 ^a		
No of wives	1.22 $\pm$ 0.78 ^a	0.85 $\pm$ 0.52 ^b	1.07 $\pm$ 0.61 ^b		
No of male children	2.68 $\pm$ 1.83 ^a	1.98 $\pm$ 1.55 ^b	2.27 $\pm$ 1.07 ^b		
No of female children	2.20 $\pm$ 1.78 ^a	1.35 $\pm$ 1.30 ^a	2.18 $\pm$ 1.44 ^a		
No of dependants	0.30 $\pm$ 0.74	0.67 $\pm$ 1.62	1.38 $\pm$ 1.15		
Experience (years)	7.20 $\pm$ 4.53 ^b	7.22 $\pm$ 4.41 ^b	9.06 $\pm$ 4.42 ^a		

SD: Standard deviation

Means followed by different uppercase letters in rows are different ($P < 0.05$)

Figure 1. Reasons for keeping goats in Nasarawa, Benue and Plateau States
1= for cash; 2= for milk; 3= for meat; 4= for cultural/religious obligations

Table 2: Flock structure of goats kept in the Nasarawa, Benue and Plateau States, north central Nigeria

Genetic group	Nasarawa Mean±SD	Benue Mean±SD	Plateau Mean±SD
Flock size	9.68±5.63 ^a	8.25±4.73 ^a	8.80±3.98 ^a
West African dwarf	7.17±4.76 ^a	6.93±4.09 ^a	7.85±3.95 ^a
Red Sokoto	1.37±3.11 ^a	0.13±1.03 ^b	0.77±2.38 ^b
Sahel	0.85±3.32 ^a	0.12±0.90 ^b	0.00±0.00 ^b
Mixed breed	0.30±1.33 ^a	0.80±2.41 ^a	0.18±1.03 ^a

SD=Standard deviation

Means followed by different uppercase letters in rows are different ($P<0.05$)

Table 3: Management systems of goats kept in the Nasarawa, Benue and Plateau States, north central

Characteristics	Nasarawa No (%)	Benue No (%)	Plateau No (%)	Pearson Chi-square	Significance
Source of stock					
Inherited	3 (30.0)	5 (50.0)	2 (20.0)		
Market	33 (39.8)	31 (37.3)	19 (22.9)		
Neighbourhood	20 (25.6)	23 (29.5)	35 (44.9)		
Borrowed	4 (4.4)	1 (1.1)	4 (4.4)	12.39	0.054
Management system					
Semi-intensive	47 (34.8)	35 (25.9)	53 (39.3)		
Intensive	2 (25.0)	4 (50.0)	2 (50.0)		
Extensive	11 (29.7)	21 (56.8)	3 (13.5)	15.33	0.004**
Access to vet					
No	16 (51.6)	3 (9.7)	12 (38.7)		
Yes	44 (29.5)	57 (38.3)	48 (32.2)	10.37	0.006**

** Significant at $P<0.01$ **Productivity parameters of goats**

Productivity indices of goats kept in the Nasarawa, Benue and Plateau States showed that for the age of parturition (1.01 ± 0.04 , 1.00 ± 0.00 and 1.00 ± 0.00 , respectively), number of kids of Sahel doe (3.32 ± 1.19 , 2.67 ± 1.51 and 3.90 ± 0.99 , respectively) and lifespan of goats in the study areas, there were no significant difference ($P>0.05$). There was also no

significant differences ($P>0.05$) in the number of kids per WAD for Benue and Plateau States, respectively (Table 4). However, more RS kids were produced in Plateau compared to the two other States. More MB kids (3.30 ± 0.67 versus 2.48 ± 0.92) were also found in Plateau than Nasarawa State. The lifespan of goat (7.37 ± 0.71 , 7.28 ± 0.64 and 7.33 ± 0.77 , respectively) appeared similar ($P<0.05$) in all the States.

Table 4: Productivity indices of goats kept in the Nasarawa, Benue and Plateau States, north central Nigeria

Indices	Nasarawa Mean±SD	Benue Mean±SD	Plateau Mean±SD
Age at first parturition	1.01 ± 0.04^a	1.00 ± 0.00^a	1.00 ± 0.00^a
Number of kids per WAD doe	4.05 ± 1.27^a	1.73 ± 1.24^{ab}	3.49 ± 1.17^a
Number of kids per RS doe	3.12 ± 1.24^b	2.68 ± 1.20^b	3.88 ± 0.89^a
Number of kids per SH doe	3.32 ± 1.19^a	2.67 ± 1.51^a	3.90 ± 0.99^a
Number of kids per MB doe	2.48 ± 0.92^b	3.11 ± 1.27^{ab}	3.30 ± 0.67^a
Lifespan	7.37 ± 0.71^a	7.28 ± 0.64^a	7.33 ± 0.77^a

WAD= West African dwarf; RS= Red Sokoto; SH= Sahel; MB=Mixed breed

SD=Standard deviation; Means bearing different uppercase letters in rows are significantly different ($P<0.05$)

Choice of breeding traits within States

Number of offspring (4.19), disease resistance (4.20), body size (4.28), survival (4.42), fertility (4.48), mothering ability (4.78) and body conformation (4.79) were highly ranked ($P < 0.05$) in Nasarawa State.

However, body size, no of offspring, fertility, survival and mothering ability were the most preferred traits for breeding stock in Benue State. In Plateau State, body size, no of offspring, disease resistance, survival and fertility were more highly rated (Table 5).

Table 5: Mean ranks of factors preferred in the choice of breeding stock of goats and their significant level according to Friedman test*

Traits	Nasarawa Mean rank	Benue Mean rank	Plateau Mean rank
Body size	4.28 ^a	4.39 ^a	4.34 ^a
Body conformation	4.79 ^a	5.05 ^b	5.22 ^{bcd}
Mothering ability	4.78 ^a	4.71 ^{abc}	5.29 ^{cd}
Survival	4.42 ^a	4.51 ^{ab}	4.47 ^a
Fertility	4.48 ^a	4.48 ^{ab}	4.48 ^a
Heat tolerance	7.38 ^b	6.73 ^c	6.87 ^c
Disease resistance	4.20 ^a	5.34 ^{bc}	4.46 ^{abc}
No of offspring	4.19 ^a	4.42 ^a	4.42 ^{ab}
Coat colour	6.48 ^b	5.57 ^c	5.45 ^c
Friedman test (chi-square)	192.64	88.48	107.08
Asymptotic significance	$P < 0.05$	$P < 0.05$	$P < 0.05$

Means in columns followed by different lower case letters are different at the Bonferroni adjusted significance level $p \leq 0.007$ (Friedman test followed by Wilcoxon signed-rank post hoc tests with Bonferroni's correction for multiple comparisons)

*The lower the mean rank, the more important the trait.

Choice of production traits within States

The choice of production traits revealed that cultural value was least ranked (5.12, 4.32

and 5.55, respectively) in Nasarawa, Benue and Plateau States (Table 6).

Table 6: Mean ranks of preferences for production traits and their significant level according to Friedman test*

Traits	Nasarawa Mean rank	Benue Mean rank	Plateau Mean rank
Body size	3.72 ^a	3.75 ^{ab}	3.69 ^a
Growth	3.88 ^a	4.14 ^{ab}	3.80 ^a
Cultural	5.12 ^b	4.32 ^b	5.55 ^b
Survival	3.98 ^a	3.98 ^{ab}	3.72 ^a
Fertility	3.78 ^a	3.85 ^{ab}	3.65 ^a
Disease resistance	3.92 ^a	4.26 ^b	3.88 ^a
No of offspring	3.60 ^a	3.70 ^a	3.72 ^a
Friedman test (chi-square)	69.35	13.601	128.57
Asymptotic significance	$P < 0.05$	$P < 0.05$	$P < 0.05$

Means in columns followed by different lower case letters are different at the Bonferroni adjusted significance level $p \leq 0.006$ (Friedman test followed by Wilcoxon signed-rank post hoc tests with Bonferroni's correction for multiple comparisons)

*The lower the mean rank, the more important the trait.

Preference for breeding traits across the three States

Among all the breeding traits across the three States, only disease resistance varied

($P < 0.01$); and this was highly rated in Nasarawa and Plateau compared to Benue State (Table 7).

Table 7: Mean ranks of factors preferred in the choice of breeding stock of goats and their significant level according to Kruskal-Wallis test¹

Traits	Nasarawa Mean rank	Benue Mean rank	Plateau Mean rank	Kruskal-Wallis test	Asymptotic significance
Body size	89.50	91.00	91.00	0.154	0.926 ^a
Body conformation	86.58	89.68	95.24	1.713	0.425 ^a
Monitoring ability	87.74	86.94	96.82	2.935	0.231 ^a
Survival	90.40	89.22	91.88	0.292	0.864 ^a
Fertility	90.11	89.95	91.44	0.105	0.949 ^a
Heat tolerance	90.97	81.88	98.66	3.623	0.163 ^a
Disease resistance	83.75 ^a	100.62 ^a	87.13 ^a	10.888	0.004 ^{**}
No of offspring	88.48	91.46	91.57	0.677	0.712 ^a
Coat colour	101.10	84.60	85.80	4.989	0.083 ^a

Means followed by different uppercase letters in rows are different (Kruskal-Wallis test followed by Mann-Whitney U tests ($P \leq 0.05$)).

¹ The lower the mean rank, the more important the trait

^a Significance at $P \leq 0.01$

^a Not significant

Preference for production traits across the three States

Across the three States, disease resistance, survival, fertility, number of offspring and body size appeared similar ($P > 0.05$) as

preference for production traits (Table 8). However, growth (Plateau State) and cultural importance (Benue State) varied across the States ($P < 0.05$ and $P < 0.01$, respectively).

Table 8: Mean ranks of preferences for production traits and their significant level according to Kruskal-Wallis test¹

Traits	Nasarawa Mean rank	Benue Mean rank	Plateau Mean rank	Kruskal-Wallis test	Asymptotic significance
Body size	90.00	93.00	88.50	1.154	0.561
Growth	90.30 ^a	97.68 ^a	83.52 ^a	6.371	0.041 [*]
Cultural	91.52 ^b	75.28 ^a	104.70 ^b	12.277	0.002 ^{**}
Survival	92.26	94.84	84.40	4.218	0.121
Fertility	90.58	94.93	85.98	3.861	0.145
Disease resistance	88.44	98.08	84.98	5.467	0.065
No of offspring	88.50	93.00	90.00	1.474	0.479

Means followed by different uppercase letters in rows are different (Kruskal-Wallis test followed by Mann-Whitney U tests). ¹ The lower the mean rank, the more important the trait

^{*}, ^{**} Significance at $P \leq 0.05$ and $P \leq 0.01$, respectively ^a Not significant

Discussion

The higher number of male respondents in this study is congruous to the submission of Chah *et al.* (2013). Similarly, Nampanzira *et al.* (2015) reported that the Ownership (65%) of goats was the responsibility of men. The present finding on marital status is

also consistent with the report of Nampanzira *et al.* (2015) that majority of the households keeping goats were married. The educational status revealed that the literacy level of majority of the farmers is low which could affect their production level. It has been argued that access to

education is related to the capacity of farmers to use appropriate information and skills for the development of their agricultural enterprise with concomitant effect on income generation (Adeleye *et al.*, 2016). The very low number of farmers with access to credit may not be unconnected with the high poverty rate at the rural level. This is not surprising because the National Bureau of Statistics (NBS) (2010) put at 68% the rural poverty rate in Nigeria. Most of these farmers were resource poor, hence relatively low farming inputs (savings inclusive). This could have also negatively affected the scale of production. In a related study, Okewu and Iheanacho (2015) reported inadequate credit facilities as a bane in goat business. Most of the goat keepers were within the active working age group. Household labour is an essential resource that influences management practices, enterprise combinations, labour hiring/sharing strategies and overall levels of technical and economic performance (ILCA, 1990). The present finding is consistent with the report of Dossa *et al.* (2008) where younger household members were more likely to own small ruminants. Adams and Ohene-Yankyer (2014) also reported farmers' average age of 47.29 ± 16.00 years. However, it is contrary to the submission of Adeschinwa *et al.* (2004) in south-west Nigeria where most of the farmers are within the age range of 51 to 60 years. It is possible that the rural farmers in the present study were facing the reality of non-availability of white collar jobs which could have probably reduced rural-urban migration of the active men and women. The average household size of this study agrees with the between 6 and 8 family members reported by Adewumi *et al.* (2015). This is an indication of availability of more labour resources that could be

drafted to goat production.

Most of the goat producers kept stock largely to generate additional money for livelihood. The variation across states may be due to socio-economic and cultural differences. This is consistent with the findings of Dossa *et al.* (2015). The flock structure revealed the preponderance of WAD goats in the three study areas. This is similar to the findings of Ajayi *et al.* (2007). The average flock sizes in the present study are comparable to the values of 8.8 and 7.8 heads for goats reported for Kano (Nigeria) and Bobo Dioulasso (Burkina Faso) (Dossa *et al.*, 2015). They are however lower than the value of 13.9 obtained in Ethiopian goats (Asefa *et al.*, 2015) and an average of 13.0 reported for goats kept by rural small holder farmers in the semi-arid regions of Uganda (Nampanzira *et al.*, 2015).

Most of the foundation stocks were purchased from the markets and neighbourhoods indicating these two sources as being important in establishing the breeding stock. The semi-intensive management system mostly adopted by the farmers is contrary to the intensive (Chah *et al.*, 2013) and extensive system reported by Oseni and Ajayi (2014). This may be due to ecological differences and prevailing resources in the study areas. The high contact of households in Benue and Plateau to health officials may be due to their willingness to pay for veterinary services. This is an indication that great potential exist in these areas to address health related challenges and probably adopt innovations. As farmers have more contacts with veterinarians, their likelihood of being medium and high level adopters increases (Boz, 2015). The productivity indices revealed the extent of twinning in goats kept by farmers in Nasarawa, Benue and Plateau States. With respect to West African Dwarf, Oseni and Ajayi (2014) reported that this

breed of goats is renowned for its adaptation to hot and humid environments, high fertility and prolificacy under backyard systems.

Body size, fertility and number of offspring were more consistent for selection purpose in the study areas. Duguma *et al.* (2010) reported that body traits including size are important criteria of selection under traditional livestock breeding practices while Akpa *et al.* (2006) submitted that parity and litter size of does were major determinants of the herd structure for breeding does. In their own findings, Nandalo *et al.* (2016) reported body size and twinning as important traits for within-breed selection as a tool for sustainable goat improvement in Malawi. In a related study, however, development and conformation were the first criteria used for choosing animals (Alexandre *et al.*, 2009). Selection of good quality breeding stock is an essential component for high productivity (Chah *et al.*, 2013). Therefore, the development of a breeding goal for improvement of indigenous stock should, according to Okeno *et al.* (2014), focus on the traits perceived important by stakeholders. This is because when breeding goats are developed without considering the needs of all the stakeholders, there are high tendencies of their being rejected by end users.

The farmers' preferences for production traits in the flock revealed the importance of all apart from cultural value. It illustrates that given a choice, farmers would prefer goat that produced more offspring (high reproductive performance), more fertile (for procreation), large body size (for meat production) and highly resistant to diseases. The present findings are consistent with the reports of Dossa *et al.* (2015). Favourable correlation suggests that if major importance is placed on performance traits

in stressful environments, adaptability traits would not be compromised and thus the most productive and adapted animals for each environment need to be identified for breeding purposes. At the micro level the definition of breeding objectives means that for the given production environment, the relative importance of improvement of different traits of the breed must be identified (Phillipson, 2011).

Conclusion

The study revealed that goats play multi-functional roles in all the study areas. The variation in the production objectives across states may be due to socio-economic and cultural differences. Farmers adopted different criteria for selection of breeding and production traits within and across the States. Breeding strategies and policies targeting goat keepers in the study areas will be more effective when incorporating indigenous knowledge of the traditional choice attributes of goats as a basis for selection criteria in their production systems.

Acknowledgement

The authors would like to express their appreciation to all the farmers and local enumerators that took part in the field study.

References

- Abdul-Aziz., M. 2010. Present status of the world goat populations and their productivity. *Lohman Information*, 45: 42–52.
- Abegaz, S., Sölkner, J., Gizaw, S., Dessie, T., Haile, A. and Wurzinger, M. 2013. Description of production systems and morphological characteristics of Abergelle and Western low-land goat breeds in Ethiopia: implications for community-based breeding

- programmes. *Animal Genetic Resources Information*, 53: 69-78.
- Adams, F. and Ohene-Yankyer, K. 2014. Socio-economic characteristics of subsistent small ruminant farmers in three regions of Northern Ghana. *Asian Journal of Applied Science and Engineering*, 3: 93-106.
- Adeleye, O., Alli-Balogun, J. K., Afima, O. G. and Bako, S. 2016. Effects of Goat Production on the Livelihood of Women in Igabi, Chikun and Kajuru Local Government Areas, Kaduna State, Nigeria. *Asian Journal of Agricultural Extension, Economics & Sociology*, 11: 1-8.
- Adeschiawa, A. O. K., Okunola, J. and Adewumi, M. K. 2004. Socio-economic characteristics of ruminant livestock farmers and their production constraints in some parts of South-western Nigeria. *Livestock Research for Rural Development*, Vol. 16, Art. #61. Retrieved October 22, 2016, from <http://www.lrrd.org/lrrd16/8/ades16061.htm>
- Adewumi, O. O., Lawal-Adebawale, O. A. and Adegbemile, D. A. 2015. Assessment of farm families' acceptability of small ruminants' milk for consumption in selected rural communities in Ogun State, Nigeria. *Journal of Agricultural Extension and Rural Extension*, 7 (4): 135-141.
- Ajayi, E. A., Okunsebor, S. A. and Yakubu, A. 2007. Pests and diseases of crops, livestock and fish. In: Akwa, V.L., Binbol, N.L., Samaila, K.I. and Marcus, N.D. (Eds). *Geographical perspective on Nasarawa State*, 1st edition, Onaivi Printing and Publishing Co. Ltd., Nasarawa State, Nigeria. Pp. 81-89.
- Akpa, G. N., Duru, S. and Bawa, G. S. 2006. The herd structure of breeding does of smallholder red Sokoto goat. *Nigerian Journal of Animal Production*, 28 (2): 119-122.
- Alexandre, G., Leimbacher, F., Maurice, O., Domarin, D., Naves, M. and Mandonnet, N. 2009. Goat farming systems in Martinique: management and breeding strategies. *Tropical Animal Health and Production*, 41: 635-644.
- Asefa, B., Kebede, K. and Effia, K. 2015. Breeding objectives, selection criteria and breeding system of Indigenous goat types in Bale zone, Oromia, Ethiopia. *Int. J. Agril. Res. Innov. & Tech.*, 5 (2): 7-15.
- Ayalew, W., Rischkowsky, B., King, J.M. and Bruns, E. 2003. Crossbreds did not create more net benefits than indigenous goats in Ethiopian smallholdings. *Agric. Sys.*, 76: 1137-1156.
- Boz, I. 2015. Adoption of innovations and best management practices by goat farmers in eastern Mediterranean Region of Turkey. *Journal of Agricultural Extension and Rural Development*, 7 (7): 229-239.
- Chah, J. M., Obi, U. P. and Ndofor-Foleng, H. M. 2013. Management practices and perceived training needs of small ruminant farmers in Anambra State, Nigeria. *African Journal of Agricultural Research*, 8 (22): 2713-2721.
- Dossa, L. H., Sangaré, M., Buerkert, A.

- and Schlecht, E. 2015. Production objectives and breeding practices of urban goat and sheep keepers in West Africa: regional analysis and implications for the development of supportive breeding programs. *SpringerPlus*, 4:281.
- Dossa, L. H., Rischkowsky, B., Birner, R. and Wolle, C. 2008. Socio-economic determinants of keeping goats and sheep by rural people in southern Benin. *Agric. Hum. Values*, 25: 581-592.
- Duguma, G., Mirkena, T., Haile, A., Iniguez, L., Okeyo, A. M., Tibbo M., Rischkowsky, B., Sölkner, J. and Wurzinger, M., 2010. Participatory approaches to investigate breeding objectives of livestock keepers. *Livestock Research for Rural Development*, 22(64).
- FAOSTAT. 2011. Food and Agricultural Organization of the United Nations. <http://faostat.fao.org/default.aspx> (19 Jul. 2011)
- Gizaw, S., Komen, H., Van Arendonk, J. A. M. 2009. Optimal village breeding schemes under smallholder sheep farming systems. *Livestock Science*, 124: 82-88.
- Haile, A., Wurzinger, M., Mueller, J., Mirkena, T., Duguma, D., Mwai, M., Sölkner, J. and Rischkowsky, R. 2011. Guidelines for Setting up Community-based Sheep Breeding Programs in Ethiopia. ICARDA - tools and guidelines No.1. Aleppo, Syria, ICARDA.
- ILCA (International Livestock Centre for Africa). 1990. Livestock systems research manual. Working Paper 1, Vol. 1. ILCA, AddisAbaba, Ethiopia. 287 pp.
- Lyam, A. (2000). Wasarawa State. In: [Mumman, A.B., Oyeboji, J.O. and Peters, S.W. (eds)], Nigeria: A People United, a Future Assured. Survey of States, Vol. 2(2), Federal Ministry of Information, Abuja.
- Nampanzira, D. K., Kabasa, J. D., Nahule, S.A., Nakalembe, I. and Tabuti, J. R. S. 2015. Characterization of the goat feeding system among rural small holder farmers in the semi-arid regions of Uganda. *SpringerPlus*, 4:188, DOI: 10.1186/s40064-015-0961-3.
- Nandolo, W., Wurzinger, M., Mészáros, G., Tassell, C.V., Gondwe, T., Mulindwa, H., Lamuno, D. and Sölkner, S. 2016. Community-based goat breeding programs in Malawi: set-up and first experiences. http://www.tropentag.de/2016/abstracts/links/Wurzinger_Z2NG_UHG.pdf
- National Bureau of Statistics. 2010. Nigerian Poverty Assessment. National Bureau of Statistics (NBS), Abuja.
- Okeno, T. O., Kahi, A. K. and Peters, K. J. 2011. Breed selection practices and traits of economic importance for indigenous chicken in Kenya. *Livestock Research for Rural Development Volume 23, Article #209*. Retrieved September 4, 2012, from <http://www.lrrd.org/lrrd23/10/oken23209.htm>.
- Okewu, J. and Iheanacho, A. C. 2015. Socio-economic characteristics

- of goat marketers in Benue State, Nigeria. *ARC Journal of Social Sciences and Humanities*, 1: 54-56.
- Oseni, S. O. and Ajayi, B. A. 2014. Phenotypic characterization and strategies for genetic improvement of WAD goats under backyard systems. *Open Journal of Animal Sciences*, 4: 253-262.
- Otuma, M. O. 2003. Genetics and Cytological Evaluation of Nigerian eco-breed goats for reproductive capabilities. A Ph.D thesis Department of Animal Science and Fisheries, Enugu State University of Science and Technology, Abakaliki Campus.
- Phillipson, J. Rege, J. E. O, Zonabend, E. and Okeyo, A. M. 2011. Sustainable breeding programs for tropical farming systems In: Animal Genetics Training resource, version 3, 2011. Ojange, J.M. Magmfors, B. and Okeyo, M.(Eds), International Livestock Research Institute, Nairobi, Kenya, and Swedish University of Agricultural Sciences, Uppsala, Sweden. <http://cgspage.cgiar.org/bitstream/handle/10568/3971/Module3.pdf>
- SPSS. 2010. *Statistical Package for Social Sciences*. SPSS Inc., 444 Michigan Avenue, Chicago, IL60611.
- Tsegahun, A., Lemma, S., Amcha, S., Abebe, M., Zinash, S. 2000. National goat research strategy in Ethiopia. In: Merkel, R. C., Abebe, G. & Goetsch, A. L. (eds) *The Opportunities and Challenges of Enhancing Goat Production in East Africa*. Proceedings of a conference held at Debub University, Awassa, Ethiopia, November 10-12, 2000. E (Kika) de la Garza Institute for Goat Research, Langston University, Langston, OK 1-5.
- Yakubu, A., Salako, A. E., Imumorin, I. G., Ige, A. O. and Akinyemi, M. O. 2010. Discriminant analysis of morphometric differentiation in the West African Dwarf and Red Sokoto goats. *South African Journal of Animal Science*, 40 (4): 381-387.
- Yakubu, A., Salako, A. E., and Imumorin, I. G. 2011. Comparative multivariate analysis of biometric units of West African Dwarf and Red Sokoto goats. *Tropical Animal Health and Production*, 43: 561-566.

Received: 3rd November, 2016

Accepted: 12th February, 2017