Fungal biodegradation of plantain peel for broiler finisher feeding: *In vitro* digestibility, effects on performance, haematological and serum parameters

*T.E Lawal, *O.M Alabi, *A.O Oguntunji, I. A Alagbe and O.A Adebiyi**

*Department of Animal Science and Fisheries Management, Bowen University, Iwo, Osun State, Nigeria

**Department of Animal Science, University of Ibadan, Ibadan, Nigeria

Corresponding author: +234(0)8034275368 zeklawal@yahoo.com

Abstract

The objective of this study was to investigate the changes in in-vitro dry matter digestibility (IVDMD) of plantain peel (PPL) after its biodegradation with *Aspergillus niger* and also to determine the effect of degraded PPL on the performance, nutrient digestibility, weights of internal organs, haematological and serum biochemical parameters of broiler finishers. A total of 165 unsexed broiler finishers that were 4 weeks old were used. *Aspergillus niger* was used for the biodegradation of PPL. There were five dietary treatments of 33 birds each. There were 0% inclusion level of PPL (control), 7% inclusion level of undegraded PPL (UPPL) then 3, 5 and 7% inclusion levels of degraded PPL (DPPL). Completely Randomized Design (CRD) was adopted. Weight gain was significantly ($P < 0.05$) higher with the birds fed degraded PPL. The highest value (46.39 gramme /bird) was found in treatment 5 (7% DPPL) and the lowest value (38.05 gramme /bird) was found in treatment 2 (7% UPPL). Feed conversion ratio ($P < 0.05$) was best (2.22) in birds placed on 7% DPPL and this is comparable to those on 0%PPL, 3%DPPL and 5%DPPL. There were significant ($P < 0.05$) differences in the digestibility of dry matter, crude protein and crude fibre. Crop, gizzard and abdominal fat were significantly ($P < 0.05$) affected. The values of packed cell volume (PCV), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC) total protein, cholesterol and glucose were significantly ($P < 0.05$) affected by the treatments. Fungal biodegradation of PPL using *A.niger* has the potential of enhancing feed intake, nutrient digestibility and the body weight gain of broiler finisher.

Key words: *Aspergillus niger*, biodegradation, nutrient enhancement and broilers.

Introduction

Agro-industrial by-products (AIBs) in most developing countries vary from primary processing of farm produce wastes to wastes from agro allied industries. Some of these wastes are left unutilized, promoting environmental pollution and hazard. Those that are utilized are not maximally harnessed. An example of agro-industrial by-products that can be of good use in feeding animals is plantain peel. Plantain peel is a lignocellulose by-product from

plantain (*Musa paradisiaca*). It is the left-over after edible part of plantain has been removed. Its inclusion in diets for monogastric animals is possibly limited by its fibrousness (Ezieshi and Olomu, 2004). Ofuya and Nwajiuba (1990) observed that the residues of agro-industrial by-products still contain considerable amount of energy and protein which are present intracellularly in the fibrillan complex. The residue represents potential valuable and renewable resources which find application

in various areas among which is its inclusion in animal feed. Faniyi (2006) reported that waste (agricultural or industrial) refers to a material which at production has low content of protein or energy, low digestibility or bioavailability, low acceptability or palatability and or toxic. It may contain chemical and microbial contaminants and generally has low feeding or nutritional value. The use of fungi in biodegradation of agro industrial by-products (AIBs) has the potential of increasing productivity, efficiency and quality output in agroindustrial processing operations in many developing countries and it has the potential of reducing the cholesterol level which may be present in them. Onilude and Oso (1999) observed that some of the commercial feeds in Nigeria lead to increase in lipids and cholesterol levels. White *et al.* (1973) opined that in humans, cholesterol is known as the cause for arteriosclerosis and it has the potential of leading to heart failure. However, Kay and Truswel (1977); Petterson and Aman (1989) reported that degraded dietary fibre fed to the birds reduced serum cholesterol concentration by maintaining a drain of the bile acid pool of the enterohepatic system. *Aspergillus niger* belongs to the family group generally described as black mould. It is capable of utilizing an enormous variety of substance for food because of the number of enzymes they produce. Iyayi and Losel (2000) stated that the use of Solid State Fermentation (SSF) for protein enrichment of lignocellulosic residues has been a focus of attention due to its direct application on the degraded product for animals as feed ingredients. The objective of this study was to investigate the ability of *A.niger* to biodegrade plantain peel and its subsequent utilization by finisher broilers.

Materials and Methods

Isolation of microorganisms

The isolation of *A.niger* was carried out on Potatoes Dextrose Agar (PDA) using the samples obtained from decaying PPL. Samples of the fungus were suspended in 10ml of sterile distilled water and shaken vigorously for 10 minutes. This was diluted with sterile water until a spore count of approximately 3×10^6 per ml was obtained. Spore count was monitored by using the Hawksley Haemocytometer. Later, 1.0ml of the resulting liquid was spread on the surface of PDA and incubated at 37°C for 7 days. The fungal isolates formed were subcultured until a pure culture was obtained.

Solid-State Fermentation (SSF) cultivation

The plantain peel used in this experiment was collected from plantain processing centre. The process of biodegradation, which lasted for 7 days, was carried out according to the method of Iyayi and Aderolu (2004) using the fungus *Aspergillus niger*. The PPL was autoclaved at 121°C for 15 minutes after which it was allowed to cool down and then inoculated with the fungus culture and moistened with distilled water at the rate of 300 ml per kg of PPL. After a period of 7 days, the action of the fungus was stopped by oven drying the substrates at 80°C for 24 hours. The dried material was then incorporated into the diets.

Data analysis

All data generated were subjected to One-Way analysis of variance and significantly different ($P < 0.05$) means were separated using Duncan Multiple Range Test as contained in Statistical Analysis System (SAS, 1999) software package.

Chemical analysis

Undegraded and degraded PPL, feed and faecal samples were subjected to proximate

analysis using the method of AOAC (1995).

Experimental Birds

One hundred and sixty five unsexed broiler finishers of Anak strain were used for this study. The chicks were reared together for 4 weeks and were placed on the same type of broiler diet. At the end of the 4th week, the birds were randomly distributed into five dietary treatments of 33 birds each which were 0 % inclusion level of PPL (control), 7% inclusion level of undegraded PPL (UPPL) then 3, 5 and 7% inclusion levels of degraded PPL (DPPL). The experimental design adopted was Completely Randomized Design. Each group was further subdivided into three replicates and each replicate was kept on litter. The birds were fed from 4-8 weeks with formulated rations. Feed and water were provided *ad libitum* and uniform light was provided. Data collected were recorded weekly. Feed consumption, weight gain and feed conversion ratio were used as measures of birds' performance. Vaccination programmes were strictly followed.

Metabolic Studies

The determination of nutrient digestibility involved the collection of faecal samples

from each of the replicate in a metabolic cage for four successive days during the 4th week of the experiment. Six birds per treatment (2 birds /replicate) with close average weight were used for this purpose. The birds were allowed three days of acclimatization before faecal collection. Feed intake was measured during the same period of 3 days at 24 hours interval. The faecal sample for each replicate was weighed and kept in aluminum foil and dried in oven to constant weight at 80 °C for 24 hours. The Metabolizable Energy (ME) of degraded and undegraded PPL feeds and faecal samples were determined by the use of Ponzenga (1985) method: $ME = 37 \times \%CP + 81.8 \times \%Fat + 35.5 \times \%NFE$

Blood Sample Collection

At the end of the 4th week, the birds were restricted from feed for 10hrs. Approximately 10 ml of fresh blood was collected. 5 ml for biochemical analysis while the other part was poured in bottle containing measured quantities of EDTA (anticoagulant for haematological analysis). Haematological parameters were determined as follows: Packed Cell

Table 1: Proximate and Detergent fibre analysis of test ingredients and diets (g/100gDM)

Parameters	UPPL	DPPL	DIET 1 0%PPL	DIET 2 7%UPPL	DIET 3 3%DPPL	DIET 4 5%DPPL	DIET 5 7%DPPL
%Dry matter	88.67	86.92	87.81	88.26	87.56	87.14	88.53
%Crude protein	8.92	13.86	18.11	18.23	18.38	18.27	18.14
%Crude fibre	14.81	10.27	3.81	3.41	3.85	3.53	3.16
%Ether extract	6.22	5.42	5.22	5.31	6.03	6.23	5.01
%Ash	6.47	7.95	17.65	17.73	18.20	17.89	18.22
(%)Nitrogen free extract	44.08	46.19	56.32	56.22	56.14	56.11	57.22
Gross energy (kcal/g)	4.28	5.25	3.68	3.51	3.22	3.61	3.76
Neutral detergent fibre	40.81	38.21	35.24	35.51	35.81	36.01	34.89
Cellulose	24.66	15.22	6.19	6.21	6.14	6.81	6.11
Hemicellulose	16.15	12.43	5.21	5.14	5.81	5.29	5.02
Acid detergent fibre	10.35	8.12	4.44	4.91	5.12	4.38	4.61
Neutral detergent lignin	14.31	11.61	8.18	8.71	8.22	8.42	8.43

UPPL=Undegraded plantain peel, DPPL=degraded plantain peel.

Fungal biodegradation of plantain peel for broiler finisher feeding

Table 2: Gross composition of experimental diets for broiler finisher using degraded and undegraded plantain peel

INGREDIENTS	Diet 1 (0% UPPL&DPPL)	Diet 2 (7% UPPL)	Diet 3 (3% DPPL)	Diet 4 (5% DPPL)	Diet 5 (7% DPPL)
Maize	50.00	49.00	49.00	49.00	56.00
Degraded plantain peel	-	-	3.00	5.00	7.00
Undegraded plantain peel	-	7.00	-	-	-
Corn offal	6.30	10.80	9.30	6.30	5.30
Groundnut cake	9.00	8.50	9.00	9.00	11.00
Soya bean meal	21.00	22.00	21.00	22.00	22.00
Fish meal	3.00	3.00	3.00	3.00	3.00
Bone meal	2.00	2.00	2.00	2.00	2.00
Oyster shell	1.00	1.00	1.00	1.00	1.00
Premix (broiler finisher)	0.25	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25	0.25
Lysine	0.10	0.10	0.10	0.10	0.10
Methionine	0.10	0.10	0.10	0.10	0.10
Total	100.00	100.00	100.00	100.00	100.00
Determined Crude Protein (%)	20.61	20.73	20.71	20.77	20.92
Determined ME (Kcal/kg)	3018.11	3012.64	3033.10	3020.74	3088.63

Provided per kg diet: 5000 IU vitamin A; 1000 IU vitamin D; 0.8 mg vitamin E; 0.4 mg Content of the Premix: menadione K3; 1.2 mg riboflavin; 1.0 mg pantothenic acid; 0.004 mg vitamin B12; 3 mg niacin; 4 mg vitamin C; 112 mg choline; 24 mg manganese; 8 mg iron and 0.048 mg selenium.

Volume (PCV), Red Blood Cell Count (RBC), White Blood Cell (WBC) and haemoglobin were determined using Wintrobe microhaematocrit. Mean corpuscular volume (MCV), Mean Corpuscular Haemoglobin Concentration (MCHC) and Mean Corpuscular Haemoglobin (MCH) were derived as outlined by Jain (1986). Where: $MCV (fl) = PCV \div RBC \times 10$, $MCH (pg) = Hb \div RBC \times 10$ and $MCHC (g/dl) = Hb \div PCV \times 100$. The serum metabolite protein, total albumin, globulin, cholesterol and glucose

were determined as described by Kaneko (1989).

Results

Table 1 shows the results of the proximate, neutral detergent fibre, acid detergent fibre and neutral detergent lignin analysis of plantain peel before and after biodegradation. *A. niger* improved the crude protein content of PPL from 8.92 to 13.86%, which is 35.64% increase. Besides, depolymerization of PPL led to reduction in crude fibre content. *A. Niger*

Table 3: Performance of broiler finishers fed undegraded and degraded plantain peel

Parameters	(0% PPL)	7% UPPL	3% DPPL	5% DPPL	7% DPPL	SEM
Initial weight(g/bird)	410.00 ^d	390.00 ^c	480.00 ^c	510.00 ^b	560.00 ^a	10.15
Final weight(g/bird)	1520.40 ^d	1455.44 ^c	1620.64 ^c	1712.76 ^b	1858.92 ^a	15.28
Weight gained at 4 th week	1110.40 ^c	1065.44 ^d	1140.64 ^b	1202.76 ^a	1298.92 ^a	12.14
Daily weight gain(g/bird)	39.66 ^d	38.05 ^d	40.74 ^c	42.96 ^b	46.39 ^a	4.23
Daily feed intake (g/bird)	115.77 ^b	128.74 ^a	110.41 ^c	106.53 ^c	102.99 ^d	5.16
Final feed intake (g/bird)	3241.56 ^b	3604.72 ^a	3091.48 ^c	2982.84 ^c	2883.72 ^d	19.12
Feed conversion ratio	2.92 ^{ab}	3.38 ^a	2.71 ^b	2.48 ^b	2.22 ^b	0.46
Nutrients digestibility (%)						
Dry matter	80.11 ^c	83.61 ^b	83.78 ^b	85.00 ^{ab}	88.72 ^a	3.23
Crude protein	68.45 ^b	68.13 ^b	72.36 ^c	75.49 ^a	76.80 ^a	2.95
Crude fibre	41.24 ^c	43.24 ^b	45.24 ^a	42.57 ^b	46.81 ^a	2.67
Ether extract	64.82	64.81	64.74	64.58	64.21	0.00
Ash	52.79	53.10	52.79	52.79	53.11	0.00

DPPL= Degraded plantain peel, UPPL=Undegraded plantain peel

a,b,c: means in the same row with different superscripts differ significantly ($P < 0.05$)

succeeded in reducing the crude fibre drastically from 14.81 to 10.27 which is 54.53% reduction. There was also improvement in Ash (6.47 to 7.95) which is 18.62% increase. The energy improved from 4.28 to 5.25kcal/kg. However, just like the crude fibre, the detergent fibre content also reduced after biodegradation of PPL with *A.niger*. The results for the performance and nutrient digestibility are shown in Table 3. Average Daily Weight Gain (ADWG) and Average Daily Feed Intake (ADFI) were significantly ($P<0.05$) affected by the treatment. The highest final

feed intake (3604.72g/bird) was found in treatment with 7% UPPL while the highest value for ADWG was 46.39g/bird and it was also found in treatment 5. There were significant ($P<0.05$) differences also in the feed conversion ratio (FCR) of the treatments and the poorest value (3.38) was found in treatment with 7% Undegraded PPL (UPPL). Significant ($P<0.05$) differences were obtained in the digestibilities of crude protein and crude fibre. Besides, crop, gizzard and abdominal fat expressed as percentage live-weight were significantly ($P<0.05$) affected as

Table 4: Relative organ proportions of broiler finishers fed degraded and undegraded plantain peel expressed as percentage live weight

Parameters (%)	0% PPL	7% UPPL	3% DPPL	5% DPPL	7% DPPL	SEM
Crop	0.81 ^b	0.98 ^a	0.78 ^c	0.76 ^c	0.76 ^c	0.065
Proventriculus	0.78	0.78	0.77	0.77	0.77	0.058
Gizzard	2.80 ^b	3.14 ^a	2.65 ^c	2.66 ^c	2.65 ^c	0.241
Small intestine weight	2.51	2.50	2.50	2.51	2.52	0.310
Large intestine weight	1.64	1.65	1.64	1.64	1.65	0.043
Caecum	0.57	0.56	0.56	0.57	0.56	0.041
Heart	0.48	0.48	0.47	0.47	0.47	0.033
Spleen	0.10	0.10	0.11	0.10	0.11	0.021
Abdominal fat	0.88 ^a	0.89 ^a	0.64 ^b	0.65 ^b	0.64 ^b	0.045

DPPL=Degraded plantain peel, UPPL=Undegraded plantain peel

a,b,c: means in the same row with different superscripts differ significantly ($P<0.05$).

shown in Table 4. Table 5 shows the results on some haematological and serum biochemical parameters investigated. The values of packed cell volume, mean corpuscular volume, mean corpuscular haemoglobin, mean corpuscular haemoglobin concentration, total protein, cholesterol and glucose were significantly ($P<0.05$) affected by the treatments.

Discussion

It was observed that the degraded PPL is higher in crude protein, ash, dry matter and nitrogen free extract but lower in crude fibre, NDF, cellulose, hemicellulose, ADF and NDL after degradation with *A. niger*. The nutritional improvement observed in degraded PPL may be explained on the

basis that the PPL served as the medium for metabolism and subsequent growth of the inoculated organism. Besides, the organism (*A.niger*) depolymerised the crude fibre and the detergent fibres therein and then converted the products to other useful components such as protein and other useful nutrients

(Okon and Ogunmodede, 1996; Liu *et al.* 2005; Sabu *et al.*, 2006). There was improvement in the gross energy content of DPPL due to hydrolysis of the crude fibre into component disaccharides and monosaccharide which can be digested, made bioavailable and utilizable for energy liberation (Faniyi, 2006). There were no

significant ($P>0.05$) differences in average initial weights but there were significant ($P<0.05$) differences in the average final weights, average final feed intake, feed conversion ratio and nutrient digestibility, except for ether extract and ash. The improvements observed in feed conversion by birds placed on DPPL can be explained by the effect of fungal enzymic degradation

conferred on target substrate. The birds fed degraded PPL had higher body weights probably because of the biodegradation of the PPL which has led to the digest of the long chain carbohydrate molecules which bacteria often use to colonize the gastro intestinal tract. This indirectly alters the bacterial population of the different regions

Table 5: Haematological and serum metabolites parameters for broiler finishers fed undegraded and degraded plantain peel

Parameters	0% PPL	7% UPPL	3% DPPL	5% DPPL	7% DPPL	SEM
PCV (%)	28.11 ^a	26.20 ^b	29.14 ^a	29.82 ^a	29.51 ^a	0.31
Haemoglobin (%)	20.44	20.50	20.32	21.54	21.60	0.043
RBC (%)	2.63	2.96	2.80	2.97	2.99	0.03
WBC($\times 10^3/\mu\text{l}$)	25200	25985	25223	25431	25511	9.07
MCV(fl)	106.88 ^a	88.51 ^c	104.07 ^a	104.40 ^b	68.59 ^d	7.41
MCH(Pg)	77.72 ^a	69.26 ^c	72.57 ^b	72.52 ^b	72.24 ^b	3.28
MCHC (g/dl)	72.71 ^b	78.24 ^a	69.73 ^c	72.23 ^b	73.19 ^b	3.33
Urea (ng/dl)	36.65 ^a	31.38 ^b	30.52 ^b	25.11 ^b	25.42 ^c	1.22
Total Protein(g/dl)	6.32 ^c	6.11 ^c	6.54 ^c	7.24 ^b	8.21 ^a	0.06
Albumin (g/dl)	2.81	2.85	2.80	2.82	2.80	0.09
Globulin(g/dl)	2.11 ^c	2.12 ^c	2.01 ^c	2.92 ^b	3.61 ^a	0.04
Arginine(g/dm)	0.78	0.81	0.82	0.81	0.79	0.003
Cholesterol (g/dl)	2.14 ^a	2.80 ^a	1.14 ^b	1.25 ^b	1.31 ^b	0.03
Triglyceride(mg/dl)	15.27 ^b	17.85 ^a	14.45 ^c	11.87 ^d	11.51 ^d	0.05
Glucose(mg/dl)	130.23 ^b	141.11 ^a	125.32 ^c	125.42 ^c	126.91 ^c	0.08

PCV=Packed cell volume, RBC=Red blood cell, WBC=White blood cell, MCV=Mean corpuscular volume, MCH=Mean corpuscular haemoglobin and MCHC=Mean corpuscular haemoglobin concentration.

DPPL=Degraded plantain peel, UPPL=Undegraded plantain peel

a,b,c: means in the same row with different superscripts differ significantly ($P<0.05$)

of the tract (Gunal and Yasar, 2004) and significantly reduces the microbial population. By implication, it increases the quantity of amino acids digested and available in the precaecal section of the tract. Besides, according to Iyayi and Davies (2005), the microfloral population in the viscous fibre-based diets do affect digestibility of amino acids in the ileum. The decreased viscosity of the digested materials and the eventual release of nutrients via the enzymatic disruptions of cell wall aided the better growth of the birds placed on degraded PPL. Increase in feed consumption by birds on treated PPL may be due to production of vitamin B complex and flavour compounds as a result of

fermentation (Sabu *et al.*, 2006). The crop is the receptacle of the feed after it has been ingested. According to Aderemi *et al.* (1999), the weight of the crop increases when bulky feed is consumed by birds. The results obtained here is in line with the work of this author because the weights of the crops of birds placed on UPPL is higher than those placed on DPPL. The weight of the gizzards in the birds fed undegraded PPL is also higher probably due to more grinding work that the gizzard had to do (Ofuya and Nwajuiba, 1990). Excess energy supplied to animal could be stored in form of fat. Feeding of UPPL like most of other fibrous agro-industrial by-products, can cause a dilution in feed energy content

and this may explain why birds on diet 2 had highest abdominal fat. The haematological values show that the PCV of birds fed UPPL were low when compared with birds on DPPL. However, the values for haemoglobin and PCV were within normal range for normal broiler as reported by Mitruska and Rawnsley (1977). This implies that the birds fed UPPL and DPPL were not anaemic. The highest value of total protein observed in the birds fed 7% DPPL probably explains the enhanced growth of the birds. This high value of 8.21g/dl suggests good quality protein of this test diet since the higher the value of the total protein, the better the quality of protein of the test feed stuff (Eggum, 1980). According to Abiola (2001), increase in urea value is an indication of poor protein quality, hence, from the results, treatment 5 (7%DPPL) gave the best value and the least was given by treatment 2 (7%UPPL). Babatunde and Oluyemi (2000) opined that, the higher the value of serum globulin, the better the ability to fight against diseases, hence the results show that treatment 5 (7%DPPL) has the highest ability to combat the body diseases. The increase in glucose level of birds fed UPPL could be due to inhibition of glycolysis by the presence of glycol-proteins and possibly other antinutritional factors which may have some adverse effect on regulation of insulin from pancreatic β -cells and on blood sugars. Cholesterol level was lower in the birds fed DPPL. This is in agreement with the findings of Onilude (1999) who reported that the cholesterol and Triglycerides consistently reduced in the blood of birds fed degraded fibrous feed. He adduced that the low cholesterol could be as a result of slight reduction in lipogenesis.

Conclusion

The use of Solid State Fermentation (SSF) to upgrade the nutritive value of PPL

enhanced the performance of the broiler finisher birds. This was reflected in improvement in crude protein, ash and gross energy of the PPL. In addition, the results from haematological and serum metabolites/biochemical parameters also showed that the performance of birds placed on DPPL is better than those placed on UPPL treatments.

References

- Abiola, S.S. 2001. Haematological indices and serum metabolites of cockerels fed hatchery waste meal diet. *Trop. Vet.*, 18: 79-84.
- Aderemi, F.A., Tewe O.O and Ogundola F.I. 1999. Enzymatic supplementation of cassava root chaff (CRC) and its utilization by pullet chicks. *Trop. Anim. Prod. Invest.* 2: 195-203.
- AOAC 1995. Official methods for Analysis of the Association of Official Analytical Chemist. 16th edn., Virginia: Arlington. ISBN 0-93558424-2.
- Babatunde, B.B and Oluyemi, J.A. 2000. Comparative digestibility of three commonly used fibrous ingredients in maize-soyabean meal-fish diet by broiler chicks. *Trop. J. Anim. Sci.*, 3: 105-111.
- Davice, J.U and Lewis S.M 1991. Practical Haematology. Eighth edition, pg. 22-68
- Eggum, B.O. 1980. Blood urea measurement as a technique for assessing protein quality *Br. J. Nutr.*, 24: 983-988.
- Ezieshi, E.V. and Olomu, J.M. 2004. Comparative performance of Broiler Chickens fed varying levels of Palm Kernel Cake and Maize Offal. *Pak. J. Nutr.*, 3: 254-257.
- Faniyi G.F. 2006. Optimizing the usefulness of poultry faeces and crop residues for meat, fish and egg

- production; Inaugural lecture at Oyo state College of Education, Oyo State. Pg 1-30.
- Gunal, M and Yasar, S. 2004.** Performance and some digesta parameters of Broiler Chickens Given low or High Viscosity Wheat Based Diets with or without Enzyme Supplementation. *Turk. J. Vet. Anim. Sci.*, 2004; 28: 323-327.
- Iyayi, E.A and Losel, D.M. 2000.** Changes in carbohydrates fractions of cassava peel following fungal solid state fermentation. *J. Food Tech.*, 6:101-103.
- Iyayi, E.A and Aderolu, A.Z. 2004.** Enhancement of the feeding value of some agro- Industrial by-products for laying hens after their solid state fermentation with *Trichoderma viridae*; *Afr. J. Biotec.*; 3:182-185.
- Iyayi, E.A., Davies B.I. 2005.** Effect of enzyme supplementation of Palm Kernel Meal and Brewer Dried Grain on the performance of Broilers. *Int. J. of Poult. Sci.*, 4: 76-80.
- Jain, N.C. 1986.** Schalm's Veterinary Haematology 4th Edn. Lea and febiger , Philadelphia. Pg 81-88.
- Kaneko, J.J. 1989.** Clinical Biochemistry of Domestic Animals 4th Edn. Academic Press Inc. N. York. Pg 102-112.
- Kay, R.M and Truwell, A.S 1977.** Effect of citrus pectin on blood lipids and faecal steroid excretion in man. *Am. J. Clin. Nutr.*, 30: 171-175.
- Liu Y.G and Baidoo, S.K 2005.** Exogenous enzymes for Pigs diets: An Overview. *Proc. of Asian conf. on Feed Biotec.* 67-70.
- Ofuya, C.O. and Nwajuiba, C.J. 1990.** Microbial degradation and utilization of cassava peel. *World J. Microbiol. Biotechnol.*, 6:144-146.
- Okon, B.I. and Ogunmodede, B.K. 1996.** Carcass characteristics of broilers fed pennloinkle flesh and palm kernel cake. *Nig. J. Anim. Prod.*, 23: 67-72.
- Onilude, A.A. 1999.** Fermented poultry litter and fungal enzyme supplementation of high-fibre broiler diets: Growth responses, carcass characteristics and blood lipids of fed chicks. *World J. Microbiol. Biotechnol.*, 43: 54-60.
- Onilude, A.A. and Oso B.A. 1999.** Effect of fungi enzyme mixture supplementation of various dietary fibres contains diets fed broiler chicks 1: performance and carcass characteristic. *World J. Microbiol. Biotechnol.*, 15:1719.
- Pauzenga U 1985** Feeding parent-stock; *Journal of Zoo technical International*, 19:22-23
- Petterson, D and Aman P. 1989.** Effects of enzyme supplementation of diets based on Wheats, rye, or triticate on their productive value for broiler chickens. *Anim. Feed Sci. Tech.*, 20: 313-323.
- Sabu, A., Augur, C., Swati, C and Pandey, A. 2006.** Tannase production by *Lactobacillus* sp. ASR SI under solid state fermentation. *J. Process Biochem.* 41:575-580.
- SAS 1999.** SAS/STAT. SAS Guide for Personal Computers Version 6th edition, Cary, NC :SAS Institute, USA.
- White, A., Handler, P and Smith, E.L. 1973.** Principles of Biochemistry, 7th Edn. Tokyo: McGraw Hill Kogakusha Ltd. ISBN 0-07-069758-2. Pg 123-134.