

GROWTH PERFORMMANCE OF BROILER CHICKENS FED VARIOUS LEVELS OF AFRICAN LOCUST BEAN (*Parkia biglobosa*) BASED DIETS

¹Shuaibu, U. S., ¹Abubakar, A., ¹Abdullahi, U. Mijinyawa, A. and ²Egbo, M. L.

¹Department of Animal Health and Production, Federal Polytechnic Bauchi, ²Department of Animal Production, Abubakar Tafawa Balewa University, Bauchi, Bauchi State, Nigeria.

Corresponding author email: umarlutiatbu@gmail.com + 2347065267274

ABSTRACT

This experiment was conducted to evaluate the growth performance of broiler chickens fed various level of African locust bean (*P.biglobosa*) based diets. One hundred and twenty day old broiler chicks (marshal strain) were allotted to four dietary treatment in which African locust bean seed replace soya beans as protein source at 0, 10, 20 and 30% levels coded as diets 1, 2, 3 and 4 respectively, The experiment was in a completely randomized design (CRD) with three replicates per treatment and ten birds per replicate. Feed and water were supplied ad libitum and the trial lasted for eight weeks. At the starter phase, daily feed intake (58.47-60.87g), daily weight gain (16.19-18.39g) and feed conversion ratio (3.22-4.02) were not significantly different. At finisher phase the daily feed intake (90.18-102.87g) indicated significant difference ($P<0.01$) while the values of the daily weight gain (34.92-40.49g) were significant in birds fed diet 4 and 3 respectively. However, feed conversion ratio did not show any significant treatment effect, the values ranged from (2.53-2.95). From these findings African locust bean meal can replaced soya bean at 20% without any adverse effect on the performance of birds with concomitant reduction in cost.

Keyword: Africa locust bean seed, Soya bean, Performance, Broiler chicken, Feed.

INTRODUCTION

Poultry production is gaining popularity in the developing countries due to its role in bridging the protein nutrition deficit, resulting to economic empowerment of the resource poor segment of society (Gebremedhn and Alemu, 2015). Such growth in the poultry industry is having a profound effect on the demand for feed and raw materials. However, it is also becoming clear that the requirements for the four traditional feed ingredients (maize, soybean meal, fish meal and meat meal) cannot be met (Adewale *et al.*, 2016). The gap between local supply and demand for these traditional ingredients are expected to be widening over the coming decades thus, providing a compelling reason for exploring the usefulness of locally available alternative feedstuffs in feed formulation (Adewale *et al.*, 2016). The utilization of locust bean in monogastric nutrition as replacement for a protein source i.e groundnut cake was reported to improved production performance, nitrogen retention and feed utilization, with varied results (Bridget *et al.*, 2004; Alabi *et al.*, 2005). Ayanwale and Ari (2002) and Dawodu (2009) found a positive attribute for fermented locust beans as against the unfermented locust bean which was said to inhibit broiler growth due to less protein quality and essential vitamins (Fetuga *et al.*, 1974). However, Anti-nutritional factors (ANFs) such as tannins oxalate and hydrogen cyanide had been reported to limit the utilization of locust bean as feed ingredient (Apata, 2003). Fermentation has been reported to destroy some natural toxins which may occur in beans, thereby improving the nutritive-value, digestibility and enhance growth of chickens (Bridget *et al.*, 2004).

This study was designed to evaluate the growth performance of broiler chickens fed various levels of African locust bean (*P.biglobosa*) meal.

MATERIALS AND METHODS

Experimental Site:

The experiment was conducted at the Poultry unit of Abubakar Tafawa Balewa University Teaching and Research Farm, Bauchi. Bauchi State occupies a total land area of 49,119 km² representing about 5.3% of Nigeria's total land mass and is located between latitudes 9° 3' and 12° 3' North and longitudes 8° 50' and 11° East.

Materials for Experiment

Maize, soya bean, locust bean, groundnut cake, fish meal, wheat offal, bone meal, limestone, lysine, methionine, salt and vitamin premix were bought from Muda Lawal market in Bauchi town. One hundred and twenty (120) day old chicks were obtained from Zatech Hatchery.

Management of the Birds

The broiler chickens were brooded on deep litter for a week fed commercial diet containing 23% crude protein and 2900kcal/kg ME. Heat was provided during the brooding with two charcoal stoves and 200watts electric bulbs as sources of light and heat. After the brooding, birds were randomly assigned to four dietary treatments. Each treatment was replicated three times, vaccine were administered as required, feed intake was determined by subtracting the left over from the supply of the previous day. The weights of the birds were measured in groups immediately after the brooding period and weekly thereafter.

Experimental Design

The experimental design used was Completely Randomized Design

Experimental Diets

Four experimental diets for both starter (23% CP) and finisher (20% CP) phase were formulated. Locust bean meal were included at 0, 10, 20 and 30% level as replacement for soya bean, designated as diets T₁, T₂, T₃ and T₄ respectively for both starter and finisher phase.

Data Collection**Birds Performance**

Records of feed intake and body weight changes were taken daily and weekly respectively, feed conversion ratio was calculated using the values while mortality was recorded as they occurred.

Statistical Analysis

The data obtained were subjected to analysis of variance (ANOVA) were significant difference occurred means, least significant difference (LSD) was used to separate them.

RESULTS AND DISCUSSIONS

The performance of broiler chickens fed various levels of African locust bean (*P.biglobosa*) based diets for starter (1-4 weeks) were presented in Table 3. There was no significant ($P>0.05$) difference in the feed intake, weight gain, and feed conversion ratio between birds fed experimental diets implying that the inclusion levels of African locust bean (*P.biglobosa*) did not have any effect on these parameters. This is in line with the finding of Obun (2007) who reported no significant difference in feed intake among chickens fed graded level of fermented African locust bean (*P.biglobosa*) meal. Also the performance of broiler chickens fed various levels of African locust bean (*P.biglobosa*) based diets for finisher (5-8 weeks) were presented in Table 4. The feed intake were significantly affected by the dietary treatments ($P<0.05$) were birds on treatment 1, 2 and 3 were statistically similar and significantly higher than treatment 4 (30%). The daily weight gains were not significantly influenced by different dietary treatments. The observed increased feed intake of birds fed African locust bean meal as shown in Table 4 can be attributed to the less odour of locust bean meal due to low inclusion level which is 20%. At the level of inclusion 100%, the smell of the ALBM masked the smell of other ingredients making the diets unattractive and possibly unpalatable to the birds, while at 50% ALBM inclusion, birds can still tolerate the aroma (Aderemi *et al.* 2017). This result is also similar to the findings of Odunsi (2003) who reported that feed intake by animal's increases if the aroma of their diet is acceptable. However, the better FCR may be due the low inclusion level of locust bean 20% which make the diet to be more palatable and acceptable.

CONCLUSION AND RECOMMENDATION

It can be concluded that the African locust bean seed can replace soya bean at 20% level as sources of protein in broiler diets at both starter and finishing phase without any detrimental effect on the performance of broiler chickens hence the use of African locust bean seed could be well advocated and recommended to served as an alternative protein source for soya bean at up to 30% level. Thereby reducing completion and improving consumer access to animal protein.

**Table 1: Percentage Composition of Experimental Diets fed to broiler chickens
At the starter phase (1-4weeks)**

Parameters	Diets			
	T ₁ (0%)	T ₂ (10%)	T ₃ (20%)	T ₄ (30%)
Maize	40.25	40.25	40.25	40.25
Soya bean	33.55	30.20	26.87	23.50
Locust bean	0.00	3.35	6.71	10.05
Groundnut cake	5.00	5.00	5.00	5.00
Fish meal	2.00	2.00	2.00	2.00
Wheat offal	15.00	15.00	15.00	15.00
Bone meal	2.00	2.00	2.00	2.00
Limestone	1.50	1.50	1.50	1.50
Salt	0.25	0.25	0.25	0.25
Lysine	0.10	0.10	0.10	0.10
Methionine	0.10	0.10	0.10	0.10
Premix	0.25	0.25	0.25	0.25
Total (%)	100	100	100	100
Calculated composition (%)				
Crude protein	23.00	23.81	23.77	23.69
crude fiber	4.09	4.06	4.04	4.01
ME (kcal/kg)	2860	2,834	2822	2812

**Table 2: Percentage Composition of Experimental Diets fed to broiler chickens
At the finisher phase (5-8weeks).**

Parameters	Diets			
	T ₁ (0%)	T ₂ (10%)	T ₃ (20%)	T ₄ (30%)
Maize	51.56	51.56	51.56	51.56
Soya bean	22.24	20.02	17.80	15.50
Locust bean	0.00	2.22	4.44	6.66
Groundnut cake	5.00	5.00	5.00	5.00
Fish meal	2.00	2.00	2.00	2.00
Wheat offal	15.00	15.00	15.00	15.00
Bone meal	2.00	2.00	2.00	2.00
Limestone	1.50	1.50	1.50	1.50
Salt	0.25	0.25	0.25	0.25
Lysine	0.10	0.10	0.10	0.10
Methionine	0.10	0.10	0.10	0.10
Premix	0.25	0.25	0.25	0.25
Total (%)	100	100	100	100
Calculated composition (%)				
Crude protein	20.00	20.81	20.77	20.69
crude fiber	5.01	5.05	5.07	5.09
ME (kcal/kg)	3,019	3,020	3,024	3,032

Table 3: Performance of broiler chickens fed various level of African locust bean at the starter phase (1-4weeks)

Parameters	Dietary treatment				SEM	LOS
	T ₁ (0%)	T ₂ (10%)	T ₃ (20%)	T ₄ (30%)		
Daily feed intake (g)	59.84	60.00	60.87	58.47	13.7	NS
Daily weight gain (g)	16.79	16.85	16.19	18.47	0.75	NS
Feed conversion ratio	3.66	3.82	4.02	3.22	0.15	NS
Mortality	0.00	0.00	0.00	0.00		

LOS: Level of significance

NS: Non significance difference (P<0.05)

Table 4: Performance of broiler chickens fed various level of African locust bean at the finisher phase (5-8weeks)

Parameters	Dietary treatment				SEM	LOS
	T ₁ (0%)	T ₂ (10%)	T ₃ (20%)	T ₄ (30%)		
Daily feed intake (g)	102.33 ^a	102.37 ^a	102.47 ^a	90.18 ^b	2.74	*
Daily weight gain (g)	36.05	40.46	38.97	34.92	3.62	NS
Feed conversion ratio	2.53	2.54	2.68	2.95	0.30	NS
Mortality(%)	3.33	3.33	6.6	3.33		

SEM: Standard Error of Mean

LOS: Level of significance

NS: Non significance difference (P<0.05)

REFERENCES

- Adewale, R. T., Adeyinka, O. A., Akinyinka, A. O., Damilola, O. O. and Joseph, A. O. (2016) Response of Broiler Chicken to Parboiled Mango Seed Kernel Meal (PMKM) Based Diet Fortified with Vitamins. *International Journal of Livestock Research*, **6** (11), 25-30
- Aderemi, F.A., Ayoola, M.O., Alabi, O.M. and Oyelami L.O. (2017). Evaluation of fermented Locust bean meal (*Parkia biglobosa*) as replacement to soybean meal in production performance, blood profile and gut morphology of broiler chicken, *Journal of Livestock Science* **8**: 28-34
- Alabi, D.A., Akinsulire, O.R and Sanyaolu, M.A (2005). Qualitative determination of chemical and nutritional composition of *Parkia biglobosa* (Jacq) Benth. *African. Journal of Biotechnology*, **4**: 812-815.
- Apata, D.F., (2003). Egg production and haematological profile of laying hens fed dietary raw or processed proposit African seeds. In *proceeding of the 28th Annual Conference of The Nigeria society for Animal production (NSAP)*, 16th-20th March 2003, 280 - 282.
- Ayanwale, B. A and M.M. Ari, (2002). Replacement value of *Parkia filicoidea* in broiler. *Journal of Agricultural. Technology*, **1**: 42-52
- Bridget, O.O., S.F. Olumuyiwa, O.A. Bolanie and S.A.A. Adewusi, (2004). Biochemical Changes in Africa locust beans (*Parkia biglobasa*) and Melon (*Citrullus vulgaris*) seeds during fermentation to condiment. *Pakistan Journal of Nutrition*, **3**: 140-145.
- Dawodu, F.A., (2009). Physico-chemical studies on oil extraction processes from some Nigerian grown plant seeds. *Journal of Environmental Agriculture and. Food Chemistry*, **8**:102-110
- Esono, B. O. And bamgbose, A. M. (2016): evaluation of the nutritive value of wild raw variegated cocoyam meal (calcium hortilinum). *Nigerian poultry of science journal*. **7** (1): 149
- Fetuga, B.L., G.M. Babatunde and V. A. Oyenuga, (1974). Protein quality of some unusual protein foodstuffs:Studies of African locust-bean seed (*Parkia filicoidea* Welw). *British Journal of Nutrition*, **32**: 27- 36
- Gebremedhn, A and Alemu, B (2015). Review of Mango (*Mangifera indica*) Seed-Kernel Waste as a Diet for Poultry. *Journal of Biology, Agriculture and Healthcare* **5**(11):2224- 3208.

- Obun, C. O. (2007). Performance and some Haematological response of Finisher Broilers Fed Graded level of fermented locust bean (*parkia biglobosa* seeds meal). *Asian journal of agricultural research*, **10**: 125-130.
- Odunsi A.A (2003). Blend of blood and rumen digesta as replacement for fish meal and groundnut cake in layers diet. *International Journal of poultry science* 2 (1):68-61