

APRW 17

Performance and Hematological Characteristics of Starter Broilers Fed Diet Supplemented with Turmeric (*Curcuma longa*) meal

A.D. Olabode¹, T.J. Ojuoloruntaye¹, C.I. Agu² and O.E. Okelola³

¹Federal College of Agriculture, Ishiagu, Ebonyi State; ²Enugu State Polytechnic, Iwollo, Ezeagu, Enugu State

³Federal College of Fisheries and Marine Technology, Victoria Island, Lagos State

Corresponding author: A.D. Olabode; E-mail: adeoladavid2005@yahoo.com; Tel: 08063379970

Abstract

The aim of the present study is to assess the growth performance and haematological assay of broiler chickens at starter phase given diets supplemented with processed turmeric (*Curcuma longa*). Ninety-six (96), two weeks old unsexed "Agritech" broiler birds were randomly allocated in a completely randomized design (CRD) into four treatments (having 24 birds each), with three replicates per treatment (having 8 birds per replicate). The treatment groups were denoted as T1, T2, T3 and T4 respectively. Feed and water were given without restriction. All dietary treatments had processed turmeric portion except the control which had none. Supplementation of processed turmeric in the diet of starter broiler significantly ($p < 0.05$) influenced the final live weight, daily feed intake, daily body weight gain and feed conversion ratio. The processed turmeric treated group had poor performance in all traits compared to the control. Mean values of all haematological parameters evaluated namely, packed cell volume (PCV), hemoglobin (Hb), red blood cell (RBC) and white blood cell (WBC) differed significantly ($p < 0.05$) between the control and the treated group. However, all the values were within the recommended range value. It was concluded that processed turmeric could be safely used as a phyto-additive in starter broiler chickens, preferably in the feed.

Keywords: Growth performance, supplemented, starter broiler, turmeric meal, hematology

Introduction

The low level of animal protein consumption by Africans in general and Nigerians specifically is a reflection of the state of deterioration of the Nigerian livestock sector, especially poultry which has continued to show a discouraging slow rate of growth (Martins, 2001). Thus a realistic step towards enhancing animal protein supply and intake in Nigeria is the development and expansion of the poultry industry (Onyimonyi *et al.*, 2009). Despite the privileged role accorded the poultry industry, its future development depends to a large extent on the supply of appropriate, safe and cost effective feeds. The interest in feed additives grew over the last decade as the usage continues to increase across the country. These feed additives have received a high attention as feed supplements for various purposes in poultry production during the recent years (Zhang *et al.*, 2009). Some of the beneficial effects of bioactive plant substance in animal nutrition may include the stimulation of appetite and feed intake.

Turmeric (*Curcuma longa*) is an extensively used spice, food preservative and coloring material that has biological actions and medicinal applications (Akpbarian *et al.*, 2012). The significant biological properties of turmeric make it a potential substitute for use as a feed additive in livestock diets. The active and main ingredient found in turmeric plants is the *curcumin*, which has been found to have antioxidant and antibacterial activities (Kerami *et al.*, 2011). Over the years turmeric has proved to have a protective effect as a feed additive on aflatoxin-induced mutagenicity and hepatocarcinogenicity in livestock production.

The aim of the present study is to assess the growth performance and hematological assay of broiler chickens at starter phase given diets supplemented with processed turmeric (*Curcuma longa*).

Materials and Method

The experiment was carried out at the poultry unit of the Federal College of Agriculture, Ishiagu, Ebonyi state. The experiment lasted for a period of five weeks. The test ingredient, which is turmeric was sourced at the Research Institute, Umudike and also locally at the Ogbete main market, Enugu state. The fresh turmeric was peeled, cleaned and washed, and later sundried before been crushed into powder form at the college feedmill.

A total of ninety-six (96) day old Agritech broiler chicks were used for the experiment. The birds were randomly distributed into four treatments namely, T1, T2, T3 and T4. Each treatment was replicated three times, with eight birds per replicate and twenty-four birds per treatment. The birds were raised on a deep litter system with wood shavings serving as a source of litter. All chicks were raised in a well ventilated and sanitized deep litter broiler poultry house using wood shavings as litter materials. Similar routine management practices were maintained for all the treatment groups. The birds were vaccinated on schedule against Newcastle, gumboro and fowlpox disease. Determination of daily feed intake was done by obtaining the difference between the quantity of feed offered and the left over the following morning. All chicks were weighed at the beginning and at the end of the study, the weight was taken weekly. The body weight gain was calculated by subtracting the initial body weight from the final body weight. The body weight change (daily body weight gain) was obtained by dividing the body weight gain by the number of

days the experiment lasted. Data generated was used to calculate the feed conversion ratio. The quantity of feed consumed compared with the unit gain as a measure of the efficiency of the feed given. This was expressed as the ratio of daily feed intake to daily weight gain.

At the end of the feeding trial, blood samples (2ml bird⁻¹) were randomly collected from one bird per replicate (three birds per treatment) from the wing vein with a sterile needle into well-labeled sterilized bottles containing EDTA as an anticoagulant. These were used for the hematological analysis to estimate the Packed Cell Volume (PCV), Hemoglobin count (Hb), Red Blood Cell (RBC) and White Blood Cell (WBC).

Table 1. Composition of starter broiler diet

Ingredients	Treatments			
	T1 (0%tm)	T2 (0.5%tm)	T3 (1.0%tm)	T4 (1.5%tm)
Maize	52.25	51.75	51.75	51.75
Wheat offal	9.50	9.50	9.00	9.00
Groundnut cake	22.15	22.15	22.15	21.65
Total	100	100	100	100

Other feed ingredient had same value across treatment group: Soybean meal-5.00; Fishmeal- 3.00; Blood meal- 3.00; Bone meal-2.50; Limestone-1.50; Salt-0.25; Starter premix-0.30; Lysine-0.25; Methionine-0.30.

Results and Discussion

The growth performance traits of birds given the experimental treatments are presented in table 2. Supplementation of turmeric powder in the diets of starter broiler birds significantly ($p<0.05$) influenced the final live weight, daily feed intake, daily body weight gain and feed conversion ratio. The result for the final body weight showed that birds in treatment 1 (control) had the highest value of 1385 g which was similar ($p>0.05$) to those of birds in treatment 2 (1357 g), but different ($p<0.05$) from those in treatment 3 (1244 g) and 4 (1159 g) respectively. The significant decrease in value for final body weight across the treatment group may be due to the inability of the birds' enzymatic system to stimulate protein synthesis and also optimized the antioxidant activities of turmeric. This present research work is in contrast with the work of Samarasinghe *et al.* (2003) who found out that inclusion of turmeric meal in the diet increased their body weight. Average daily feed intake showed significant ($p<0.05$) effect among treatment group. Values for birds in treatment 1 (98.37 g) was similar ($p>0.05$) to those on treatment 2 (97.20 g), but was significantly ($p<0.05$) different from those obtained for birds on treatment 3 (90.18 g) and 4 (88.66 g). The result was in agreement with the work carried out by Rajput *et al.*, (2013) that found a significant difference in feed consumption, but disagrees with the observation of Nouzarian *et al.* (2011). Thus, the low level of feed intake observed in treatments containing turmeric meal could be due to the inability of the birds gut to maximize the turmeric meal.

Table 2: Performance and haematological characteristics indices of starter broilers fed diet supplemented with turmeric meal.

Parameters	Treatments				SEM
	T1	T2	T3	T4	
Initial body weight (g)	259	253	259	257	-
Final body weight (g)	1385 ^a	1357 ^a	1244 ^b	1159 ^c	35.33
Average daily Wt. gain (g)	53.62 ^a	52.57 ^a	46.91 ^b	42.95 ^c	1.70
Average daily feed intake (g)	98.37 ^a	97.10 ^a	90.18 ^b	88.66 ^c	1.39
Feed conversion ratio	1.84 ^c	1.85 ^c	1.92 ^b	2.06 ^a	0.05
Packed Cell Volume (%)	22 ^a	22 ^a	15 ^b	21 ^a	0.87
Haemoglobin (g/dl)	7.3 ^a	7.4 ^a	5.0 ^b	7.0 ^a	0.30
Red blood cell ($\times 10^{12}/l$)	2.3 ^a	2.6 ^a	1.6 ^b	2.6 ^a	0.13
White blood cell ($\times 10^9/l$)	5.4 ^c	6.2 ^b	4.5 ^d	7.5 ^a	0.34

The result of the haematological indices of starter broilers fed diets supplemented with turmeric meal is presented in table 2. Haematological parameters are good indicators of the physiological status of the animal (Esonu *et al.*, 2006). The blood parameters for hematology were significantly ($p<0.05$) affected by the *Curcuma longa* across the treatment group, even though they did not follow a particular pattern. All values obtained for packed cell volume (PCV), hemoglobin (Hb), red blood cell (RBC) and white blood cell (WBC) studied falls within the normal ranges for broiler as reported by The Merck Veterinary Manual (2015). The results obtained in the present study contradicts the findings of Abou-Elkhair *et al.* (2014) who reported that inclusion of turmeric meal in the diet of starter broilers had no effect on PCV, Hb, BBC and WBC respectively. The improvement in the haematological constituents observed in the present study is an indication that the inclusion of turmeric meal did not show any pathological effect on the bird thus did not cause or create any haematological disorder.

Conclusion

Generally, the results of this study indicate that the use of turmeric meal in the diets of broiler chickens does not have any adverse effect on them. Thus, the addition of turmeric meal upto 1.5% can be tolerated by the broiler birds at their starter stage without any negative effect. In conclusion, turmeric meal could be safely used as a phyto-additive for broiler chickens preferably in the feed.

References

- Abou-Elkhair, R., Ahmed, H.A. and Selim, S. (2014). Effect of black pepper (*Piper nigrum*), turmeric powder (*Curcuma longa*) and coriander seeds (*Coriandrum sativum*) and their combinations as feed additives on growth performance, carcass traits, some blood parameters and humoral immune response of broiler chickens. *Asian-Australas. J. Anim. Sci.*, 27(6):847-854.
- Akbarian, A., Abolghasem, G., Ahmadi, S. and Hossein, M. (2012) Effect of ginger root (*Zingiber officinale*) on the egg yolk cholesterol, antioxidant status and performance of laying hens. *Journal of Applied Animal Science*, 39: 19-21.
- Esonu, B.O., Opara, M.N., Okoli, I.C., Obikaonu, H.O., Udedibie, C. and Iheshiolor, O.M. (2006). Physiological response of laying birds to neem leaf meal based diets; body weight, organ characteristics and hematology. *Online Journal of Health and Allied Science*, 5: 26-30.
- Kerami, M., Alimon, A.R., Sazili, A.Q., Goh, Y.M. and Ivan, M. (2011). Effect of dietary antioxidants on the quality, fatty acid profile and lipid oxidation of longissimus muscle in Kacang goat with aging time. *Meat Sci.*, 88: 102-108.
- Martins, A.M. (2001). The future of the world food system. *Outlook on Agriculture*, 30: 11-19.
- Nouzarian, R., Tabeidian, S.A., Toghyani, M., Ghalamkari, G. and Toghyani, M. (2011). Effect of turmeric powder on performance, carcass traits, humoral immune responses and serum metabolites in broiler chickens. *J. Anim. & Feed Sci.*, 20: 389-400.
- Onyimonyi, A.E., Olabode, A.D and Okeke, G.C (2009). Performance and economic characteristics of broilers fed varying levels of neem leaf meal. *International Journal of Poultry Science*, 8(3): 256-259.
- Rajput, N., Muhammed, N., Yaan, R., Zhang, X. and Wang, T. (2013). Effect of dietary supplementation of *curcumin* on growth performance, intestinal morphology and nutrients utilization of broiler chicks. *J. Poult. Sci.*, 50: 44-52.
- Samarasinghe, K., WenkSilvak, K. and Gunasekera, J. (2003). Turmeric (*Curcuma longa*) root powder and mannan-oligosaccharides as alternatives to antibiotics in broiler chicken diets. *Asian-Australas. J. Anim. Sci.*, 16(10): 1495-1500.
- The Merck Veterinary Manual (2015). Recommended nutrient level for pullet chicken. A subsidiary of Merck and Co. Inc., Kenilworth, N.J., U.S.A
- Zhang, G.F., Yang, Z.B., Wang, Y., Yang, W.R., Jiang, S.Z. and Gai, G.S. (2009). Effects of ginger root (*Zingiber officinale*) processed to different particle sizes on growth performance, antioxidant status and serum metabolites of broiler chickens. *Poultry Science*, 88:2159-2166.