

ESTIMATION OF GENETIC PARAMETERS OF BIOMETRIC AND ALLOMETRIC TRAITS IN FOUR STRAINS OF THE NIGERIAN HEAVY LOCAL CHICKEN ECOTYPE (NHLCE)

Ohagenyi I.J C.W Eze and Udeh V. C.

Department of Animal Science, Faculty of Agriculture, University of Nigeria, Nsukka, Enugu state.
e-mail: ifemma.ohagenyi@unn.edu.ng

Abstract

The study was conducted to determine some genetic parameters of biometric and allometric traits in the Nigerian heavy local chicken ecotype of Nigeria. This study was carried out at the Animal Science teaching and research, University of Nigeria, Nsukka. Data from 80 progenies produced by the artificial insemination of 4 male heavy ecotype local chickens with 12 mature female chickens was used for this study. The data analyzed consisted of body weight records from 0-8 weeks as well as conformation traits (body length, chest girth, shank length) at 1, 4, 6 and 8 weeks. The results show that there was no significant difference ($p>0.05$) on the body weight of progenies at weeks 1 and 4, body length of progenies at weeks 1, 4 and 6, girth of progenies at weeks 1, 4 and 8, shank length at weeks 1 and 4. However, there was significant difference ($p<0.05$) on body weight of progenies at weeks 6 and 8, body length at week 8, chest girth at week 6 and shank length at weeks 6 and 8. The heritability estimates for body weight, body length, chest girth and shank length ranged from low to high with an estimated value of -0.44 to 0.62 respectively. However, high heritability estimates were reported for body weight, body length and shank length (0.40, 0.40, 0.62 respectively). From this study, body weight, body length and shank length at 8 weeks could be used as selection indices for selection of parents of next generation. It's not advisable to carry out selection at young age because heritability at early ages are been influenced by maternal and environmental effects.

Key words: Additive; Genetic progress; Heritability; Selection; Variance;

Introduction

The population of chicken in Nigeria has been estimated at approximately 166 million (FAOSTAT, 2007). The local chicken constitutes about 80% of this number (Dana *et al.*, 2011). The Nigerian local chicken is characterized by poor growth, small body size, small egg size and egg number which is not desirable in a competitive economy (Ebangi and Ibe, 1994).

Despite these undesirable characteristics, the local chicken still plays important role in the rural economy of Nigeria by providing meat, egg and house hold income to the rural people. Studies have shown that the local chicken in Africa exhibit high genetic variability within their populations indicating their potential for genetic improvement through selective breeding (Osei-Amponsah *et al.*, 2010, Dana *et al.*, 2011). Recent works revealed that the different ecotypes can be grouped into two major categories on the basis of body size and body weight as heavy ecotype and light ecotype (Momoh *et al.*, 2007). The heavy ecotype (also referred to as Fulani ecotype) is found in the dry Savannahs (Guinea and Sahel Savannah), montane regions and cattle Kraals of the North and weigh about 0.9-2.5 kg at maturity. The light ecotype are those chicken types from the Swamp, Rainforest and Derived Savannah agro-ecological zones whose mature body weight ranges between 0.68-1.5 kg.

Knowledge of genetic parameters is necessary for designing an appropriate breeding plan for genetic improvement of NLC (Ajayi, 2009). Relationships exists between body weight and linear body measurements. This is to organize the breeding programme so as to achieve an optimum combination of body weight and good conformation for efficient production and maximum economic returns. Body weight and body dimensions have been used as parameters for selection by local sellers and for research (Ajayi, 2010). The aim of this work is to estimate the genetic parameters of biometric traits in four strains of the Nigerian heavy local chicken ecotype (NHLCE).

Material and Methods

Location and duration of study

The experiment was carried out at the Poultry Unit of the Department of Animal Science research farm, University of Nigeria, Nsukka. The study lasted for a period of 2 months.

Foundation stock, experimental birds and management

The foundation stocks were 16 mature Nigerian heavy ecotype local chickens, comprising of 4 cocks and 12 hens. Mating of likes (associative mating) strains which are the black, barred, white and brown was adopted to generate progenies. The experimental birds consisted of 80 progenies produced from artificial insemination involving four different strains of the heavy ecotype local chicken (white, barred, black and golden). The experimental birds were grouped into 4 consisting of 20 chicks of the golden strain, 20 chicks of the white strain, 20 chicks of the barred strain and 20 chicks of the black strain and the brooding house was divided into 4 sections to house them respectively from 0-8 weeks. Necessary vaccinations and medications were administered in addition to proper management practices.

Data collection and statistical analysis

Data obtained from body weight, body length, shank length and chest girth was subjected to analysis of variance (ANOVA) in a paternal half sib method, all data generated was analyzed using SAS (2004) statistical procedure.

Experimental design

The experimental design used was the paternal half sib analysis

Model: $Y_{ijk} = \mu + \alpha_i + e_{ijk}$,

Where: μ - overall mean

α_i - random effect of the sire

e_{ijk} = the uncontrolled environment and genetic deviation attributed to individual progeny

Results and Discussion

The photos of the progenies of the four genetic groups of the Nigerian heavy local chicken ecotypes are shown in figure 1 below.

FIGURE 1. PICTURE SHOWING THE PROGENIES OF THE FOUR STRAINS OF THE NHLCE AT 4 AND 8 WEEKS OF AGE.



4 Weeks
barred strain



4 Weeks
black strain



4 weeks golden
strain



4 weeks white
strain



8 weeks
barred
strain



8 weeks
black strain



8 weeks golden
strain



8 weeks white
strain

Table 1. Descriptive statistics for biometric and allometric traits of four genetic group of Nigerian Heavy Local Chicken Ecotype

Parameters	Week	White strain	Barred strain	Black strain	Golden/brown strain
BW(KG)	1	0.038±0.002 ^{N.S}	0.036±0.002 ^{N.S}	0.036±0.022 ^{N.S}	0.036 ±0.002 ^{N.S}
	4	0.1960±0.009 ^{N.S}	0.960±0.002 ^{N.S}	0.200±0.009 ^{N.S}	0.188±0.008 ^{N.S}
	6	0.312±0.0143 ^b	0.360±0.0157 ^{ab}	0.370±0.0179 ^a	0.346± 0.016 ^{ab}
	8	0.564± 0.226 ^{ab}	0.516± 0.256 ^b	0.524±0.0173 ^{ab}	0.584 ±0.010 ^a
BL(CM)	1	6.410±0.177 ^{N.S}	6.450±0.116 ^{N.S}	6.270±0.209 ^{N.S}	6.410 ±0.094 ^{N.S}
	4	11.160 ±0.82 ^{N.S}	10.910±0.167 ^{N.S}	10.920±0.19 ^{N.S}	11.140±0.139 ^{N.S}
	6	14.220±0.222 ^a	14.350±0.263 ^a	14.330±0.265 ^a	14.200 ±0.251 ^a
	8	15.600±0.385 ^b	15.600±0.385 ^b	16.500±0.433 ^{ab}	17.020 ±1.255 ^a
CG(CM)	1	6.390±0.198 ^{N.S}	5.910±0.200 ^{N.S}	6.170±0.752 ^{N.S}	5.970±0.586 ^{N.S}
	4	10.290±0.065 ^{N.S}	10.150±0.005 ^{N.S}	10.260±0.177 ^{N.S}	10.180±0.916 ^{N.S}
	8	16.010±0.545 ^{N.S}	16.280±0.396 ^{N.S}	16.200±0.667 ^{N.S}	16.200±0.667 ^{N.S}
	8	16.010±0.545 ^{N.S}	16.280±0.396 ^{N.S}	16.200±0.667 ^{N.S}	16.200±0.667 ^{N.S}
SL(CM)	1	2.230±0.085 ^{N.S}	2.310±0.202 ^{N.S}	2.300±0.051 ^{N.S}	2.290±0.096 ^{N.S}
	4	4.980±0.129 ^{N.S}	5.000±0.210 ^{N.S}	5.000±0.066 ^{N.S}	4.940±0.106 ^{N.S}
	8	6.880±0.196 ^{ab}	6.500±0.290 ^b	6.500±0.290 ^b	7.360±0.160 ^a

Means having different subscripts in the same row are significantly different (P<0.05)

Legend:

BW-body weight, BL: body length, CG-chest girth, SL- shank length, N.S-not significant

The table 1 above shows the means and standard error for biometric traits of 4 genetic groups of the Nigerian heavy local chicken ecotype for four weeks

The mean result of the study on body weight of white, barred, black and golden strain for week 1,4,6,8 ranges from 0.036 ± 0.002 to 0.038 ± 0.002 , 0.188 ± 0.008 to 0.200 ± 0.009 , 0.312 ± 0.014 to 0.370 ± 0.017 , 0.516 ± 0.256 to 0.584 ± 0.010 , respectively. There was no significant difference ($P > 0.05$) in body weight of progenies of the four genetic group at week 1, they were statistically similar. However, at 6 weeks of age, there was significant differences ($P < 0.05$) in the body weight of the progenies of the genetic group. The body weight of the white, barred and black strain was statistically similar and significantly higher ($P < 0.05$) than bodyweight of progenies of white strain. At 8 weeks, there was significant difference ($P < 0.05$), body weight values for progenies of golden, black and white strain was statistically similar and significantly higher than bodyweight value of progenies of barred strain.

The mean body length of the progenies of 4 genetic groups of the heavy ecotype local chicken for 4 weeks ranges from 6.410 ± 0.094 to 6.450 ± 0.116 , 10.910 ± 0.167 to 11.160 ± 0.482 , 14.200 ± 0.251 to 14.830 ± 0.265 , 15.600 ± 0.385 to 17.020 ± 1.255 respectively. There was no significant differences ($P > 0.05$) in body length of progenies of the four genetic group at weeks 1, 4 and 6, they were statistically similar. But at 8 weeks of age, there were significant differences ($P < 0.05$) in the body length as body length values of progenies of barred, black and golden strain are statistically similar and significantly higher ($P < 0.05$) than body length values of progenies of white strain. The mean and standard error for the chest girth of 4 genetic groups of heavy ecotype local chicken varies from 5.910 ± 0.200 to 6.390 ± 0.198 , 10.150 ± 0.05 to 10.290 ± 0.065 , 16.010 ± 0.545 to 16.280 ± 0.396 respectively. There was no significant difference ($P > 0.05$) for the chest girth values observed at week 1 and 4, and week 8 of age. The mean outcome of the research on shank length for white strain, black strain, barred strain, golden strain were 2.230 ± 0.085 to 2.310 ± 0.200 for week one, 4.980 ± 0.129 to 5.000 ± 0.210 , for week 4, 6.500 ± 0.290 to 7.360 ± 0.160 for week 8 respectively. No significant differences ($P > 0.05$) were observed at week 1 and week 4. However, at week 8, the mean value of the shank length of progenies of golden strain, white strain were statistically similar and significantly higher ($P < 0.05$) than the shank length values of the progenies of black and barred strain.

Heritability estimates for biometric traits in 4 strains of the Nigerian Heavy Local Chicken Ecotype (NHLCE).

Body weight:

Table 2 below shows the heritability estimates for biometric traits in 4 strains of the Nigerian Heavy Local Chicken Ecotype (NHLCE).

Table 2. Heritability estimates for biometric traits of 4 strains of the Nigerian heavy local chicken ecotype

TRAITS	Vs	Vt	Heritability
Bw(kg)			
Week 1	0.000	0.000	0.000
Week 4	0.001	0.02	0.2
Week 6	0.005	0.055	0.36
Week 8	0.009	0.09	0.4

The heritability of body weight ranged from low to high. Low (0.00) at week 1 and high (0.4) at 8 week. The result revealed that heritability estimates for body weight at week 1,4,6 and 8 were 0.00,0.2,0.36 and 0.4 respectively. It shows that body weight of heavy ecotypes were low at week 1, moderate at week 4 and 6, and high at 8 weeks, indicating that selection of body weight of heavy ecotypes should be done at 8 weeks of age, and it reveals that mass selection at 8 weeks could lead improve their genetic performances.

Heritability estimates for Allometric traits in 4 strains

The heritability estimates for allometric traits in 4 strains of the Nigerian Heavy Local Chicken Ecotype (NHLCE) are presented in the tables 3a, 3b and 3c below.

Table 3: The heritability estimates for body length in 4 strains of the Nigerian heavy local chicken ecotype (NHLCE)

TRAITS	Vs	Vt	Heritability
BL(CM)			
Week 1	0.027	0.381	0.29
Week 4	0.061	0.8571	0.28
Week 6	0.323	0.607	0.15
Week 8	0.1835	1.7905	0.4

The result of the study revealed that the heritability for body length at week 1,4,6,8 of ages were 0.29 (moderate), 0.28 (moderate), 0.15 (Low) and 0.40 (high) respectively.

The heritability estimates of body length observed in this study ranged from 0.15 (low) at week 6 and 0.40 (high) at 8 weeks of age and moderate (0.29,0.28) at week 1 and 4. The result shows that selection for body weight at 8 weeks could improve genetic performance of the heavy ecotype local chicken.

Chest girth:

Table 4: The heritability estimates for chest girth in 4 strains of the Nigerian heavy local chicken ecotype (NHLCE)

TRAITS	Vs	Vt	Heritability
CG(CM)			
Week 1	0.0046	0.4306	0.04
Week 4	-0.007	1.09	-0.25
Week 8	-0.323	3.042	-0.42

The result in table 4 revealed that the heritability estimates for chest girth of heavy ecotype local chicken at 1, 4, 8 weeks were 0.04, -0.25, -0.42. The heritability estimates of chest girth ranged low (0.04), indicating that selection family/pedigree selection could be done and negative at week 4 and 8 which could be due to small sampling size.

Shank length:

Table 5: The heritability estimates for shank length in 4 strains of the Nigerian heavy local chicken ecotype (NHLCE)

TRAITS	Vs	Vt	Heritability
SL(CM)			
Week 1	-0.034	0.572	-0.23
Week 4	-0.092	0.8378	-0.44
Week 8	0.083	0.690	0.62

The heritability estimates of the study shows that the shank length of heavy ecotype local chicken at week 1, 4, 8 were -0.23, -0.44 and 0.62 respectively indicating that heritability estimates for shank length was negative due to small sampling size and high at the week 8 inferring that mass selection can be used to improve shank length of heavy ecotypes at the age of 8 weeks. The result obtained from 0 to 8 weeks in this study for biometric traits: body weight, body length, chest girth and shank length amongst four strains or genetic groups of the heavy ecotype local chicken namely white strain, black strain, barred strain and

golden strain shows an increasing trend supporting that age is a major determinant of growth and physiological development. This report formally agrees with Momoh (2007) that the body weight of the heavy ecotype local chicken ranges from 0.9kg to 2.5 kg.

Discussion:

The heritability estimates of body weight at week 1 is in agreement with the report of Aslam *et al.* (2011) who reported 0.00 at day old to 6 days. It is lower than the value given by Frank (2018) who reported 0.02 and Ohagenyi *et al.* (2012) who reported 0.2 in NHLCE. The heritability estimates at week four reported in this study was within the same range (0.21 and 0.25) reported by Ajayi (2010), Ohagenyi *et al.* (2012) and Ogbu (2012) in NHLCE. It corresponds to Ndofor-Foleng *et al.* (2006) who reported heritability estimates of body weight as 0.40 for heavy ecotypes of Nigerian local chicken at week 8. However, it was higher than the values (0.26 and 0.35) given by Ogbu (2012) and Ohagenyi *et al.* (2012) among the NHLCE and lower than the values (0.60) and (0.54) reported by Agu (2012) for heavy ecotypes and Osei Amposah *et al.* (2013) for the local chicken of Ghana at the age of 8 weeks.

These differences in heritability estimates for body weight could be due to differences in strains of chicken used and environment, this agrees with the fact that heritability is affected by population.

The heritability estimates for body length of NHLCE at weeks 4 and 8 were higher than the values (-0.06 and 0.27) reported by Ohagenyi *et al.* (2012).

The estimated value for shank length reported in this study at weeks 4 and 8 were lower than the values (0.23 and 0.18) reported by Ohagenyi *et al.* (2012) for NHLCE. The shank length and chest girth reported in this study at week 4 and week 6 are in contrast with the values reported by Ige (2013) and Agbo (2012), who reported a heritability estimate of 0.73 for shank length and 0.89 for chest girth in Fulani ecotypes. It is higher than the value (0.42) given by Osei- Amposah *et al.* (2013) for shank length at week 8 for local chicken of Ghana. The existing Variances found in the estimated values could be due to environment (differences in techniques used, differences in strains of chicken used and management).

Generally, the heritability estimates for body weight fall within range of previously reported heritability estimates. This study also reveals that fitness traits such as BWT and SHL tend to have higher heritability estimates compared to traits that are not connected to reproductive efficiency of animals. The low heritability at younger ages indicates that body weight and other traits at these ages are to a very large extent a function of environmental factors, the implication of this is that the selection of traits of interest may not result in any appreciable improvement, this explains why day old body weights was not heritable. The high heritability estimates for body weight as well as moderate to high heritability estimates for body length implies that additive genetic variance made a greater contribution to the total phenotypic variance compared to environmental and gene combination variance. It's an indication that variability due to additive gene action is probably higher than non additive component, implying that a larger proportion of superiority of parents will be retained in the offspring and genetic progress can be made through mass selection. Selection should be carried out at ages where heritability estimates were high so as to ensure genetic improvement.

Conclusion

This study shows that the heavy ecotypes vary significantly in their biometric and allometric traits. The heritability estimates reported for traits studied (body weight, body length, chest girth and shank length) were low at 1 week old; low and moderate at 4 and 6 weeks of age, but high at the age of 8 weeks. High heritability estimate at week 8 is an indication that variability due to additive gene action is probably higher than non additive component and implies that a larger proportion of superiority of parents will be retained in the offspring and genetic progress can be made through mass selection at week 8.

References

- Agbo, M.C. (2018). Genetic response of productive traits of the Nigerian heavy local chicken ecotype using selection index from fourth to sixth generation
- Agu, C., Ndofor-Foleng, H.M. and Nwosu, C.C. (2012). Evaluation of economic traits in progenies of Nigerian heavy ecotype chicken as genetic material for development of rural poultry production. *African journal of biotechnology*.
- Ajayi FO 2010 Nigerian indigenous chicken: A valuable genetic resource for meat and egg production. *Asian Journal of Poultry Science* v.4, p.164-172.
- Ajayi FO and BO Agaviezor 2009 Phenotypic characteristics of indigenous chicken in selected local government areas in Bayelsa state, Nigeria. 3rd Nigeria International Poultry Summit, Feb. 22-26, 2009 Abeokuta, Proceedings p.75-78.
- Asadu C LA 2002 Fluctuations in the characteristics of an important short tropical season, august break in eastern Nigeria, discovery and innovation .IITA England, v.14n.1.
- Aslam Mlbastiaansen, JWM, Croojimans RPMA, Ducro B, JVerrijken A and Groenend AM 2011 Genetic variances, heritabilities and maternal effects on body weight, breast meat yield, meat quality traits and the shape of the growth curve in turkey birds. *BMC Genetics*, v.12, p.14-23.
- Dana N and Vanderwaaij E H 2011 Genetic and phenotypic parameter estimates for bodyweight and egg production in Horro chicken of Ethiopia. *Tropical Animal Health and Production* 43v.1p. 21-28
- Ebangi A L and Ibe SN 1994 Heritabilities of and genetic correlations between some growth traits in Nigerian local chickens. *Nigerian Journal of Animal Production* v.21p.19-24.
- FAOSTAT 2007 Food and Agricultural Organization statistical databases.
- Ohagenyi IJ, Iregbu FC and Udeh VC 2021 Estimation of genetic parameters for body weight and some colour traits in index selected Nigeria heavy local chicken ecotype. *Agricultural science Digest*, 10.
- Hardiman J 2007 "How 90 years of poultry breeding has shaped today's industry" (PDF). *Poultry International*. Archived from the original (PDF) on May 25, 2012. Retrieved July 1, 2012.
- Ige AO 2013 Estimation of genetic parameters in Yoruba and Fulani ecotypes indigenous chickens of Nigeria, *Transitional journal of source of science and technology*.
- Kinney TB 1969 A summary of reported estimates of heritabilities and of genetic and phenotypic correlations for traits of chickens.
- Momoh OM and CC Nwosu 2008 Genetic evaluation of growth traits in crosses between two ecotypes of Nigeria Local Chicken Livestock Resources. *Rural Dev.* v.20.
- Momoh OM, Ehiobu NG and Nwosu CC 2007 Egg production of two Nigerian local chicken ecotypes under improved management. 32nd Annual conference of Nigerian Society for Animal Production, March 18-22, University of Calabar, Nigeria, Proceedings p.278-281.
- Ogbu CC, Udeh F and Nwakpu PC 2012 Comparative performance of two commercial egg strains, the indigenous chickens and their random bred progenies. *Livestock research for rural development*.
- Ohagenyi IJ 2009 Some biometric and allometric growth traits of purebred heavy ecotype of the Nigerian local chicken. Msc. thesis submitted to University of Nigeria, Nsukka.
- Ohagenyi IJ, Emennaa CI, Adeloowo OV, Brenshak SB, Udokainyang AD, Ogbu KI and Amah NE 2012 Heritability Estimates of growth traits of the Nigerian Heavy Ecotype Chicken. The 17th animal science association of Nigeria, 2012. Proceedings
- Osei-Amponsah R, Kayang BB, Naazie A, Osie YD, Youssao AK, Yapi-Gnaore VC, Tixier-Boichard M and Rognom X 2010 Genetic diversity of forest and savannah chicken populations of Ghana as estimated by microsatellite markers. *Animal Sci G* 1994 *Animal breeding. Tropical Agriculturist*, Macmillan Education Limited London.