Economic evaluation, carcass characteristics and organ weights of broiler chickens fed graded levels of raw tropical sickle pod (*Senna obtusifolia*) seed meal based-diets

Augustine, C.,^{*1} Kwari, I. D.², Igwebuikwe, J. U.², Adamu, S. B.², Medugu, C. I.² Obidah, L. U.¹ Khobe, D.¹ and Babale, D. A. M.¹

¹Department of Animal Production, Adamawa State University, Mubi,

²Department of Animal Science University of Maiduguri.

Abstract Corresponding author audaggai@gmail.com, GSM 08132946167

A study was conducted to investigate the effect of raw *Senna obtusifolia* seed meal (RSOSM) on carcass characteristics, organ weights and economic performance of broiler chickens in a 9 weeks trial. One hundred and eighty (180) broiler chicks were managed on deep litter pens. Six (6) experimental diets were formulated to contain RSOSM at concentrations of 0, 5, 10, 15, 20 and 25%, and designated as T1, T2, T3, T4, T5 and T6, respectively. The broiler chicks were randomly allotted to the 6 dietary treatments in a randomized complete block design with pen location serving as the blocking factor. Data were collected on final live weight, plucked and dressed weight, cut-up parts and economic performance of broiler chickens fed RSOSM based-diets. The data collected were analysed using a computer statistical package (Statistix, 9.0). Carcass parameters such as dressed weight and dressing percentage were observed to be depressed as the level of RSOSM increases in the diets. Dressing percentage decreased from 78.69 to 52.38% in T1 to T6, respectively. The weight of cut-up parts followed similar trend as above. Thigh and drum stick significantly ($P < 0.05$) decreased from 10.16 to 7.88 g and 9.01 to 6.11g in T1 to T6, respectively. The result of organ weights indicated significant ($P < 0.05$) reduction in weights as the level of RSOSM increases in the experimental diets except for the liver which indicated significant ($P < 0.05$) increase in weight as the levels of RSOSM increases in the diets. On economic ground, inclusion of RSOSM beyond 5% in the diets of broiler chickens is not cost effective. This is because feed cost per kilogram weight gain was observed to increase as the level of RSOSM increases in the diets. Feed cost per kilogram body weight gain was observed to increase from N223.84 to N290.18 in T1 to T6, respectively. In conclusion, incorporation of RSOSM beyond 5% in the diets of broiler chickens showed depressive effects on carcass yield and organs weight. Furthermore, inclusion of RSOSM beyond 5% in the diets of broiler chickens is not cost effective. It is recommended that the seeds of RSOSM be thoroughly processed before incorporation in the diets of broiler chickens. There is also urgent need to conduct more studies on the best processing method(s) that will effectively detoxify RSOSM and improve its utilization.

Keywords: Broiler chickens, raw *Senna obtusifolia*, toxicity depressive effect

Introduction

The rapid growth and expansion of the Nigerian poultry industry is limited by the scarcity and high cost of conventional feed resource. In this regard, there is urgent need to look inward and exploit alternative feed resources that are available, cheap and can serve as good replacers for the costly feed resources. Adegbenro *et al.* (2011) further buttressed on the utilization of under-

exploited. *Senna obtusifolia* is one of such ideal lesser-known legumes that have long been under-utilized as feed ingredient for poultry in Nigeria *Senna obtusifolia* is a short-lived (annual or biennial) shrub growing up to 2.5 m tall, but usually less than 2 m in height. The leaves are pinnate and alternately arranged along the stem, born on petioles 15-20 mm long. The flowers are yellow (10-15 mm across) while the fruit is

slender, strongly curved downward (sickle shape), pod (6-18 cm long and 2-6 mm wide). The seeds (3-6 mm long) are dark brown in colour, shiny in appearance with rhomboid or irregular shape (Agnes, 2011). The chemical composition of the seeds as reported by Ingweye *et al.* (2010) and Augustine *et al.* (2017) revealed that they have good protein content (29.54 and 23.40%) and fair distribution of amino acids. However, the authors further revealed that the seeds contain anti-nutritional factors which might limit nutrient utilization and adversely affect overall animal performance and carcass yield. At the moment, base-line information on the adverse effects of raw *Senna obtusifolia* seed meal based-diets on carcass characteristics, organ weights and economic performance of broiler chickens seems to be inadequate. Therefore, more studies are required to bridge this information gap. It is in view of the above, that this study was conducted to evaluate the effects of feeding raw *Senna obtusifolia* seed meal on carcass yield organ weight and economic performance of broiler chickens.

Materials and methods

Location of the study area

The research was conducted at the Poultry Unit of the Department of Animal Production Livestock Teaching and Research Farm, Adamawa State University, Mubi. Mubi is located between latitudes 9°30' and 11° North of the equator and longitudes 13° and 13° 45' East of the Greenwich meridian. The temperature regime in Mubi region is warm to hot throughout the year however, there is usually a slight cold period between November and February. There is a gradual increase in temperature from January to April. The minimum and maximum

temperatures of the area are 18.1°C and 32.8 °C. The mean annual rainfall ranges from 900-1050 mm (Adebayo, 2004).

Experimental chickens and their management

One hundred and eighty broiler (180) chicks were used for the experiment which lasted sixty three (63) days. The broiler chickens were managed on deep litter pens and vaccinated against Gumboro (2 and 4 weeks of ages) and Newcastle disease (3 and 5 weeks of age). The experimental broiler starter diet was fed *ad-libitum* to the chicks for forty two (42) days while the broiler finisher diet was fed for the remaining twenty one (21) days, respectively. Clean drinking water was similarly supplied *ad-libitum* throughout the period of the experiment.

Experimental diets

Six experimental diets for both the starter (23% CP) and finisher (20% CP) phases, respectively were formulated to contain 0, 5, 10, 15, 20 and 25% of raw *Senna obtusifolia* seed meal replacing soybean meal in T₁, T₂, T₃, T₄, T₅ and T₆, respectively. The 0% replacement level (T₁) served as the control diet. The ingredient and calculated chemical composition of the experimental diets are presented in Tables 1 and 2.

Experimental design

The 180 chicks were individually weighed and randomly allotted to the six (6) dietary treatments in a randomized complete block design (RCBD) with pen location serving as the blocking factor. Each treatment group consisted of thirty (30) birds replicated three times with ten (10) birds per replicate.

Parameters measured

Carcass and organ evaluation

At the end of the experiment, three (3) broiler chickens were randomly selected from each replicate in each treatment. The broiler chickens were deprived of feed overnight but allowed access to water and

the fasted live weights of individual chickens were recorded. The birds were slaughtered and bled by severing the jugular vein. The chickens were dipped in warm water and plucked. The plucked carcass weights were recorded and thereafter eviscerated through a slit made between the end of the keel bone and the rectum. Dressed weight and cut-up parts (head, neck, wings, breast, shanks, drumsticks, thighs and back) and the organs

(heart, lungs, liver, gizzard, and kidney), abdominal fats and intestine were weighed and recorded.

Dressed weight was determined by weighing the dressed chickens and dressing percentage (DP) was determined by dividing the dressed weight by the live weight expressed as percentage. It can be mathematically expressed as follows:

$$DP = \frac{\text{Carcass weight}}{\text{Live weight}} \times 100$$

Table 1: Ingredient Composition and Calculated Analysis of the Experimental Broiler Starter Diets

Ingredient	Level of inclusion of raw SOSM					
	T1(0%)	T2(5%)	T3(10%)	T4(15%)	T5(20%)	T6(25%)
Maize	50.00	50.00	50.00	50.00	50.00	50.00
Roasted soya bean	27.00	22.00	17.00	12.00	7.00	2.00
Groundnut cake	10.00	10.00	10.00	10.00	10.00	10.00
RSOSM	0.00	5.00	10.00	15.00	20.00	25.00
Fish meal	5.00	5.00	5.00	5.00	5.00	5.00
Maize offal	4.00	4.00	4.00	4.00	4.00	4.00
Salt (NaCl)	0.30	0.30	0.30	0.30	0.30	0.30
Bone meal	3.00	3.00	3.00	3.00	3.00	3.00
Methionine	0.30	0.30	0.30	0.30	0.30	0.30
Lysine	0.15	0.15	0.15	0.15	0.15	0.15
Premix*	0.25	0.25	0.25	0.25	0.25	0.25
Total	100.00	100.00	100.00	100.00	100.00	100.00
Calculated analysis						
ME energy (kcal/kg)	2892.14	2857.50	2822.14	2788.21	2753.57	2718.93
Crude protein (%)	23.47	23.07	22.67	22.55	21.20	20.45
Crude fibre (%)	3.09	3.87	3.92	3.99	4.11	4.33
Methionine (%)	0.72	0.72	0.77	0.79	0.82	0.87
Lysine (%)	1.24	1.25	1.26	1.27	1.28	1.29
Calcium (%)	1.69	1.70	1.72	1.73	1.74	1.76
Phosphorus (%)	0.75	0.72	0.70	0.68	0.65	0.63

*2.5kg Vitamin–Mineral Premix (Animal Care) supplies the following: Vitamin 12,000,000 vit. D₃ 3000,000, vit. E 30,000 mg, vit. K₁ 2,500 mg, folic acid 1,000 mg, niacin 40,000, folic acid, 1,000 mg, calpan, 10,000 mg vit. B₂ 5000 mg, vit. B₁₂ 20 mg, vit. B₁ 2,000 mg, vit. B₆ 3,500 mg, biotin 80 mg and antioxidant 125,000 mg, cobalt 250 mg, selenium 250 mg, iodine 1,200 mg, iron 40,000 mg, manganese 70,000mg, copper 8,000 mg, zinc 60,000 mg, chloride 200,000 mg Metabolizable energy (ME) calculated according to the formula of Ponzenga, (1985) ME= 37 × %CP + 81 × % EE + 35.5 × % NFE , RSOSM = Raw *Senna obtusifolia* seed meal

Economic evaluation, carcass characteristics and organ weights of broiler chickens

Table 2: Ingredient Composition and Calculated Analysis of the Experimental Broiler Finisher Diets

Ingredient	Level of inclusion of raw SOSM					
	T1(0%)	T2(5%)	T3(10%)	T4(15%)	T5(20%)	T6 (25%)
Maize	54.00	54.00	54.00	54.00	54.00	54.00
Roasted soya bean	27.00	22.00	17.00	12.00	7.00	2.00
Groundnut cake	7.00	7.00	7.00	7.00	7.00	7.00
Fish meal	3.00	3.00	3.00	3.00	3.00	3.00
Maize offal	5.00	5.00	5.00	5.00	5.00	5.00
RSOSM	0.00	5.00	10.00	15.00	20.00	25.00
Salt (NaCl)	0.30	0.30	0.30	0.30	0.30	0.30
Bone meal	3.00	3.00	3.00	3.00	3.00	3.00
Methionine	0.30	0.30	0.30	0.30	0.30	0.30
Lysine	0.15	0.15	0.15	0.15	0.15	0.15
Premix*	0.25	0.25	0.25	0.25	0.25	0.25
Total	100.00	100.00	100.00	100.00	100.00	100.00
Calculated analysis						
ME energy (kcal/kg)	2948.38	2900.03	2879.09	2856.46	2821.01	2778.37
Crude protein (%)	21.01	20.61	20.21	19.85	19.76	19.45
Crude fibre (%)	3.69	4.11	4.23	4.38	4.49	4.51
Methionine (%)	0.70	0.72	0.72	0.73	0.75	0.77
Lysine (%)	1.04	1.13	1.38	1.41	1.44	1.42
Calcium (%)	1.14	1.14	1.16	1.18	1.21	1.23
Phosphorus (%)	0.68	0.61	0.59	0.51	0.49	0.45

*2.5 kg Vitamin–Mineral Premix (Animal Care) supply the following: Vitamin 12,000,000 vit. D₃ 3000,000, vit. E 30,000 mg vit. K₃ 2,500 mg, folic acid 1,000 mg, niacin 40,000, folic acid, 1,000 mg, calpan, 10,000 mg vit. B₂ 5000 mg vit. B₁₂ 20 mg, vit. B₁ 2,000 mg, vit. B₆ 3,500 mg, biotin 80 mg and antioxidant 125,000 mg, cobalt 250 mg, selenium 250 mg, iodine 1,200 mg, iron 40,000 mg, manganese 70,000mg, copper 8,000 mg zinc 60,000 mg chloride 200,000 mg Metabolizable energy (ME) calculated according to the formula of Ponzenga, (1985) ME= 37 × %CP + 81× % EE + 35.5 × % NFE, RSOSM = Raw *Senna obtusifolia* seed meal

Cost-benefits of using *Senna obtusifolia* as feed ingredient for broiler chickens

The cost-benefits of using *Senna obtusifolia* as feed ingredient for broiler chickens was assessed by computing feed cost per kilogram, cost of total feed intake and feed cost per kilogram body weight gain using prevailing current prices as at 2015:

Feed cost per kilogram was calculated using the prevailing market prices of each feed ingredient at the time of study.

Cost of total feed intake (N/kg) = Total feed intake x feed cost per kilogram.

Feed cost/ body weight gain(N /kg gain) = $\frac{\text{Cost of total feed consumed N}}{\text{Total body weight gain (kg)}}$

Total body weight gain (kg)

Statistical analysis

Data obtained were subjected to analysis of

variance (ANOVA) of the randomized complete block design (RCBD) using Statistix 9.0 (Statistix, 2003). Least Significant Difference (LSD) was used to separate the treatment means where significant differences occur. Significant difference was considered at 0.05 level of significance.

Results and discussion

Proximate composition of experimental diets

The proximate compositions of the experimental diets are presented in Tables 3 and 4. The composition of the starter and finisher diets indicated that the nutrients profile of the diets are adequate for optimal performance of broiler chickens at all phases of growth. The tannin content of the experimental diets revealed that the levels of

tannins in the diets increased as the levels of RSOSM increases from 0 to 25%. Similar observation was made by Kaya *et al.* (2011). The authors reported that diets containing raw Vetch seed (*Vicia sativa*) at

higher level (25%) showed higher tannin content. The levels of tannins in these diets are enough to cause depressive effects on the performance of broiler chickens as reported by Hagerman and Klucher (1986)

Table 3: Proximate composition and levels of anti-nutritional factors of experimental broiler starter diets

Parameters	Level of inclusion of raw RSOSM					
	T1(0%)	T2(5%)	T3(10%)	T4(15%)	T5(20%)	T6(25%)
Dry matter (%)	93.18	92.29	93.15	93.02	93.49	93.57
Crude protein (%)	22.15	20.08	21.25	22.92	21.00	20.89
Crude fibre (%)	4.51	6.16	6.23	6.28	6.51	6.58
Ether extract (%)	6.12	5.98	6.08	7.02	7.09	7.21
NFE (%)	50.75	50.51	50.46	49.66	49.58	48.87
Tannin (%)	0.00	0.16	0.38	0.47	0.58	0.61
Total phenols (%)	0.00	0.46	0.57	0.68	0.86	0.97
ME energy (kcal/k)	3116.89	3026.71	3070.06	3179.59	3182.42	3091.83

Metabolizable energy (ME) calculated according to the formula of Ponzenga, (1985) $ME = 37 \times \% CP + 81 \times \% EE + 35.5 \times \% NFE$; NFE = Nitrogen -free extract; RSOSM = Raw *Senna obtusifolia* seed meal

Table 4: Proximate composition and levels of anti-nutritional factors of experimental broiler finisher diets

Parameters (%)	Level of inclusion of raw RSOSM					
	T1(0%)	T2(5%)	T3(10%)	T4(15%)	T5(20%)	T6(25%)
Dry matter	93.05	93.40	92.77	93.40	93.26	93.54
Crude protein	20.05	19.06	19.64	19.31	19.26	20.00
Crude fibre	8.78	8.42	8.63	8.77	8.81	8.95
Ether extract	7.03	7.45	7.52	7.63	8.43	8.93
NFE	54.70	53.47	50.23	49.66	49.19	46.13
Tannins	0.00	0.24	0.40	0.47	0.51	0.68
Total phenols	0.00	0.43	0.61	0.73	0.79	0.86
ME energy (kcal/kg)	3253.28	3206.86	3118.97	3095.43	3141.70	3100.95

Metabolizable energy (ME) calculated according to the formula of Ponzenga, (1985) $ME = 37 \times \% CP + 81 \times \% EE + 35.5 \times \% NFE$; NFE = Nitrogen-free extract; RSOSM = Raw *Senna obtusifolia* seed meal

Carcass characteristics of broiler chickens fed Raw Senna obtusifolia seed meal

The impact of the treatment diets on carcass yield and visceral organs of broiler chickens fed raw *Senna obtusifolia* seed meal (RSOSM) are presented in Table 5. Carcass yield is an indication of the quality and utilization of the ration (Bamgbose and Niba, 1998). The values of fasted live weight, plucked weight, dressed weight

and dressing percentage revealed significant ($P < 0.05$) variations among the treatments. Broiler chickens fed 0 and 5% RSOSM recorded superior ($P < 0.05$) dressed weight and dressing percentage compared to the groups fed 10, 15, 20 and 25% RSOSM. This is a clear indication that, the chickens fed high concentrations of the seed meal poorly utilized their diets which consequently affected their dressed weight and dressing percentage. This suggests that inclusion of

Economic evaluation, carcass characteristics and organ weights of broiler chickens

RSOSM at levels higher than 5% resulted to high density of anti-nutritional factors such as tannins in the diets. The implication is poor nutrient utilization which affected weight gain and carcass yield. This finding agrees with the report of Uchegbu *et al.* (2004) who observed low dressed weight and dressing percentage of broiler chickens fed raw *Napoleona imperialis* seed meal. Similar effects were observed in the cut-up parts. Higher body weight was said to attract higher cut-up parts (Ojewole *et al.*, 2000).

The dressing percentages for the broiler

chickens fed 0 to 20% RSOSM falls within the range of 60.96 to 66.09% reported by Odetola *et al.* (2011). This range was earlier reported by Bamgbose and Niba (1998) as the carcass yield for eight weeks old broiler chickens. Chickens fed up to 10% RSOSM had the highest breast, thorax, thighs and drum stick while broiler chickens fed 20 and 25% RSOSM recorded the lowest values of cut-off parts. The values for wings, thighs and drumstick falls within the range of 7.88 to 8.79 g, 10.35 to 11.06 g and 9.89 to 12.22 g, respectively reported by Adetola *et al.* (2012) for broiler chickens fed *Sesamum indicum* based diets.

Table 5: Carcass Characteristics of Broiler Chickens Fed Raw *Senna obtusifolia* Seed Meal (RSOSM)
Level of inclusion of raw SOSM

Parameters	T1(0%)	T2 (5%)	T3 (10%)	T4(15%)	T5(20%)	T6(25%)	SEM
Live Wt. (g)	2033.30 ^a	1883.30 ^{ab}	1600.00 ^c	1300.0 ^d	1100.30 ^e	1050.65 ^e	6.65*
Plucked wt (g).	1883.30 ^a	1800.00 ^a	1616.70 ^b	1216.70 ^c	1200.00 ^c	1033.30 ^{cd}	155.25*
Dressed wt. (g)	1600.00 ^a	1350.00 ^b	1100.00 ^c	850.00 ^d	700.00 ^d	550.00 ^e	44.36*
Dressing %	78.69 ^a	71.68 ^b	68.75 ^c	65.38 ^{cd}	63.64 ^d	52.38 ^e	4.12*
Cut-up parts as percentage of slaughter weight							
Head	3.24 ^a	2.49 ^b	2.51 ^b	2.42 ^c	2.40 ^c	3.26 ^a	1.21*
Neck	4.67 ^a	4.42 ^a	4.76 ^a	4.69 ^a	3.78 ^b	3.88 ^b	1.19*
Breast	14.75 ^a	14.29 ^a	14.75 ^a	11.00 ^c	13.91 ^b	13.03 ^b	4.69*
Back	8.07 ^b	7.46 ^c	7.57 ^c	6.37 ^d	7.39 ^c	9.68 ^a	3.43*
Thorax	5.59 ^a	5.52 ^a	5.85 ^a	4.96 ^b	4.43 ^b	5.30 ^a	1.42*
Wings	8.40 ^a	8.33 ^a	8.32 ^a	8.14 ^a	7.66 ^b	5.13 ^c	3.47*
Shanks	4.92 ^b	4.20 ^b	4.69 ^b	4.88 ^b	3.42 ^c	3.24 ^c	0.30*
Thighs	10.16 ^a	10.64 ^a	10.05 ^a	8.89 ^b	7.13 ^c	9.88 ^b	4.7*
Drumstick	9.01 ^a	9.69 ^a	10.21 ^a	8.69 ^b	8.88 ^b	8.11 ^b	1.14*
Internal organs and other visceral components as percentage of slaughter weight							
Proventriculus	0.57 ^a	0.54 ^a	0.55 ^a	0.49 ^b	0.49 ^b	0.33 ^c	0.03*
Full gizzard	3.75	3.69	4.27	4.10	4.38	4.40	1.80 ^{NS}
Empty gizzard	2.13	2.23	2.01	2.52	2.03	2.07	0.43 ^{NS}
Heart	0.44	0.47	0.42	0.49	0.49	0.46	0.03 ^{NS}
Liver	1.85 ^b	1.66 ^b	1.72 ^b	1.88 ^b	1.90 ^b	2.02 ^a	0.09*
Kidney	0.48 ^b	0.59 ^a	0.46 ^b	0.55 ^a	0.68 ^a	0.63 ^a	0.09*
Lungs	0.64 ^a	0.52 ^b	0.62 ^a	0.52 ^b	0.57 ^b	0.59 ^b	0.06*
Pancreas	0.22	0.29	0.29	0.24	0.24	0.29	0.01 ^{NS}
Spleen	0.13	0.10	0.21	0.10	0.12	0.11	0.04 ^{NS}
Abdominal fat	2.97 ^a	2.94 ^a	1.41 ^b	1.03 ^b	1.24 ^b	1.09 ^b	0.12*
I.L(cm)	189.00 ^b	209.00 ^a	218.33 ^a	191.00 ^b	201.00 ^a	197.06 ^b	3.09*
Intestinal wt.	4.68 ^a	1.19 ^b	2.02 ^b	1.13 ^c	1.33 ^c	1.98 ^b	0.39*

a,b,c,d = means in the same row with different superscripts are significantly different (P<0.05), NS = Not significant (P>0.05) * = Significant at 0.05% level of confidence; SEM = Standard error of means and I.L = Intestinal Length

The result of organ weights showed significant ($P<0.05$) differences among the treatments except for the gizzard. The general trend revealed that organ weights decreased as the level of RSOSM increased in the diets indicating positive correlation between organ weights and live body weight. This was supported by Young *et al.* (2001) who earlier explained that live weight is among the factors that affects yield of parts and therefore large body size may attract larger organs. However, the relative weight of liver was higher at 20 and 25% level of inclusion. This could be due to high concentration of anti-nutritional factors. This agreed with report of Uchegbu *et al.* (2004) who reported a linear increase in the weight of organs (liver, heart and gizzard) as the level of raw *Napoleona imperialis* seed meal increases in the diets of broiler chickens. They attributed this effect to higher physiological activities of these organs triggered by the presence of anti-nutritional factors and their concomitant effects.

The result of intestinal weight, length and caecal length were significantly ($P<0.05$) different among the treatment groups. They exhibited a decreasing pattern as the concentration of RSOSM increased in the diets. This could be due to the body sizes relative to their gut capacity. Bigger chickens are expected to possess larger digestive tract to accommodate more feed materials. The impact of the dietary treatments on abdominal fat content of broiler chickens fed dietary treatments indicated that the chickens fed 0 and 5% RSOSM significantly ($P<0.05$) recorded high abdominal fat compared to the other treatment groups. The general trend revealed decline in abdominal fat content as the level of RSOSM increased in the diets. This showed that anti-nutritional factors such as saponins and tannins can negatively

affect fat deposition in broiler chickens. This was buttressed by Ueda (2001) and Dong *et al.* (2007) who explained that saponin plays a role in lowering cholesterol and fat due to its ability to form insoluble complexes bond with fat and cholesterol from food in the gut, so that they cannot be reabsorbed by the intestine. Wardah *et al.* (2012) further pointed out that anti-nutritional factors have the ability to reduce intracellular lipid accumulation in the tissues of animals. It is therefore evident that raw *Senna obtusifolia* seed meal may be used as a natural feed supplement to reduce fat content and cholesterol levels in broiler chicken meat. This may be of health benefit to consumers that are conscious of cholesterol in their diets.

Cost-benefits of feeding broiler chickens with raw *Senna obtusifolia* seed meal

The cost-benefits of feeding broiler chickens with graded levels of RSOSM is presented in Table 6. The results revealed that cost of feed per kilogram, which is one of the major economic indices, significantly ($P<0.05$) reduced as the level of RSOSM increased in the diets. However, feed cost per kilogram body weight gain steadily increased as the level of RSOSM increased in the experimental diets. This indicated poor nutrient utilization as the concentration of RSOSM increased from 0 to 25% hence the ability of the chickens to convert nutrients to flesh declined. This effect may be attributed to the presence of anti-nutritional factors in the RSOSM. Similar observation was made by Yisa *et al.* (2012) who reported poor feed conversion ratio and increasing feed cost per kilogram body weight gain of broiler chickens fed raw and boiled pigeon pea seed meal. On economic grounds, feeding RSOSM at 15, 20 and 25% is not cost effective.

Economic evaluation, carcass characteristics and organ weights of broiler chickens

Table 6: Cost-Benefit of Feeding Broiler Chickens with Graded Levels of Raw *Senna obtusifolia* Seed Meal (RSOSM)

Parameters	Level of inclusion of raw SOSM						SEM
	T1(0%)	T2(5%)	T3(10%)	T4(15%)	T5(20%)	T6(25%)	
Total FI (g)	5.62 ^a	5.44 ^a	5.36 ^a	4.79 ^b	4.49 ^b	4.53 ^b	0.03*
Feed cost (₦/kg)	77.22	73.97	70.72	64.47	63.22	61.97	-
Total feed cost (₦)	433.40	402.39	379.05	308.81	283.85	280.72	-
Total weight gain (kg)	1.94 ^a	1.67 ^b	1.40 ^c	1.10 ^d	1.06 ^d	1.05 ^d	3.21*
Feed cost/kg gain (₦)	223.94	241.14	270.86	281.08	290.18	286.92	-

a,b,c,d,e = means in the same row with different superscripts are significantly different (P<0.05), * = significant at 0.05, SEM = standard error of means, & FI = Feed intake

References

- Adebayo, A. A. 2004.** *Mubi region a geographical synthesis* paraclete publishers Yola, Nigeria 133 Pp.
- Adegbenro, M., Onibi, G. E., Agbede, J. O., Aletor, V. A., Aro, S. O. and Adeyeye, S. A. 2011.** Evaluation of the nutrients and anti-nutritional factors in some under-utilized seeds. *Proceedings of the 16th annual Conference of Animal Science Association of Nigeria (ASAN)*, September 12th – 15th 2011, Kogi State University, Anyigba, Nigeria, Pp96-98.
- Adetola, O. O., Sajo, A. P., Ogunwole, O. A. and Omojola, A. B. 2012.** Growth carcass characteristics and organ weights of broiler chickens fed toasted sesame seed (*Sesamum indicum*, linn) base diets. *Proceedings of the Annual Conference of Animal Science Association of Nigeria* 13th – 19th September, Abuja, Nigeria. Pp 500 – 502.
- Agnes, L. 2011.** Invasive species compendium online data sheet. Sickie pod(*Senna obtusifolia*). CABI publishing, 2011. www.cabi.org/isc. Accessed 21st August, 2017.
- Augustine, C., Kwari, I. D., Igwebuike, J. U. and Adamu, S. B. 2017b.** Evaluation of chemical composition of raw and processed tropical sickle pod (*Senna obtusifolia*) seed meal. *Agricultural Science and Technology*, 9(2):110-113.
- Bamgbose, A. M. and Niba, A. T. 1998.** Performance of broiler chickens fed cotton seed cake in starter and finisher rations. *Proceedings of the the 3rd Annual Conference of Animal Science Association of Nigeria*, September, 22nd – 24th Lagos Pp 84 - 37.
- Dong, X. F., Gao, W. W., Tong, J. M., Jia, H. Q., Sa, R. N. and Zhang, Q. 2007.** Effect of polysavone (Alfalfa extract) on abdominal fat deposition and immunity in broiler chickens. *Poultry Science*, 86:1955-1959.
- Hagerman, A. E. and Klucher, K. M. 1986.** Tannin protein interaction. In: *Plant Flavanoids in Biology and Medicine: Biochemical Pharmacological and Structure-activity Relationships*. Editors: Cody, V., Middleton E.Jr., Harbone, J. and Alanr-Liss. New York, Pp. 67-76.
- Kaya, H., Celebi, S., Macit, M. and Geyikolgu, F. 2011.** The effects of raw and physically processed common Vetch seed (*Vicia sativa*) on laying performance, egg quality, metabolic parameters and liver histopathology of laying hens. *Asian-Australasian Journal of*

- Animal Science*, 24(10): 1425-1434.
- Odetola, O. M., Eruvbetine, D., Oduguwa, O. O., Onagbesan, O. M. and Adeolu, M. 2011.** Performance and carcass characteristics of broilers fed varying levels of raw kenaf meal as replacement for full fat soya meal. *Proceeding of the 16th Annual Conference of the Animal Science Association of Nigeria (ASAN)* September, 12th – 15th, Kogi State University, Anyigba, Nigeria. Pp391-392.
- Ojewole, G. S., Uka, S. N. and Onyenuchey, F. 2000.** Comparative carcass characteristics of indigenous poultry fed different agro-industrial by-products. *Tropical Journal of Animal Science*, 3(2):159 – 161.
- Pauzenga, U. 1985.** Feeding parent stock. *Zootecnia International* Pp22-25.
- Statistix 2003.** Statistix for windows manual. Analytical Software. Version 8.0.
- Uchegbu, M. C., Okoli, I. C., Anyanwu, C. E., Etuk, E. B., Esonu, B. O. and Udedibie, A. B. I. 2004.** Performance, carcass and organ characteristics of finisher broilers fed graded levels of raw *Napoleona imperialis* seed meal. *Livestock Research for Rural Development*, volume 16 www.org/Irrd/6/16/Uche/6041.htm retrieved 2nd August, 2013.
- Ueda, H. 2001.** Short-term feeding response in chicks to tea saponin. *Journal of Poultry Science*, 86:1955-1959.
- Wardah, T. S., Bimo A. E. and Kusurningrum, H. 2012.** Reduction of intracellular lipids accumulation serum leptin and cholesterol levels in broilers fed diet supplemented with powder leaves of *Phyllanthus buxifolius*. *Asian Journal of Agricultural Science*, 6:106 – 117.
- Yisa, A. G., Yisa, M. K., Edache, J. A. and Ogedegbe, S. A. 2012.** Effect of graded levels of boiled and dried pigeon pea (*Cajanus cajan* (L) millsp) on growth performance of broiler starters. *Proceedings of 14th Annual Conference of Animal Science Association of Nigerian (ASAN)* 9th- 13th Abuja, Nigeria. Pp387 – 390.
- Young, L. L., Northcutt, J. K., Buhr, R. J., Lyon, C. E. and Ware, G. O. 2001.** Effect of post mortem ageing on percentage yield of parts from broiler chicken carcasses. *Poultry Science*, 80:376-379.

Received: 20th July, 2017

Accepted: 13th December, 2017