

CARCASS YIELD AND ECONOMICS OF PRODUCTION OF GROWING RABBITS (*Oryctolagus cuniculus*) FED DIFFERENT TYPES AND LEVELS OF RUMEN CONTENT

^{1*}ABDULLAHI, A. M., ¹MAIDALA, A., ²DOMA, U.D., ¹LAWAN, A., ¹ABDULLAHI, U.,
¹CHANA, I. M., ¹MAKINDE, O., ³OYAWOYE, E., & ⁴TAMBURAWA, M. S.

¹Department of Animal science, Faculty of Agriculture, Federal University Gashua, Yobe State

²Department of Animal Production, Abubakar Tafawa Balewa University, P.M.B. 248. Bauchi State

³Department of Animal Production, Landmark University, Omu Aran. Kwara state

⁴Department of Animal Production, Kano State University, Wudil. Kano State

Corresponding author: alhajimusaabdullahigadaka@mail.com+2348024991014

*Presenting author

Abstract

An experiment was conducted to investigate the carcass yield and the economics of rabbit production. Rabbit were fed different types and levels of rumen content. Forty-eight (48) mongrel rabbits were randomly allotted to six diets containing rumen content of *caprine*, *ovine* and *bovine* each had 30 and 40% levels of inclusion in 3x2 factorial arrangements completely randomized design (CRD). Results showed that the interaction effect did not affect live weight (1075.00-1487.50g), carcass weight (456.25-775.00g) and dressed weight (38.36-52.86%). There were significant different ($P>0.05$) in the grow performance of rabbit fed different levels of rumen content. Most of the organs were not affected by the interaction effect ($P<0.05$) except the lungs (0.42-0.63%), head (7.72-9.25%) and small intestine (3.50-4.38%) which were affected by the interaction effects ($P<0.05$). Most of the parameters assessed were not affected by the main effect ($P>0.05$) of types of rumen contents except the kidney (0.15-0.64%), stomach (1.2-1.58%) and small intestine (0.93-1.19%) which were affected ($P<0.05$). The economics of production showed that feed cost per kg decreased slightly at higher levels of inclusion for the three types of rumen content. The feed cost per kg gain also decreased as the level of rumen content increase. Therefore rumen content should be included at higher levels in the diet of rabbits for enhanced carcass yield and reduce cost of production

Keywords: Rumen content, Carcass yield, Caprine, Ovine, Bovine

Introduction

The dietary protein intake in Nigeria is 4.5g grossly short of the recommended 35g of animal protein intake per person per day (FAO, 2001). This observed low animal protein intake may be attributed to the declining animal production occasioned by the high cost of feeds which usually accounts for 70-85% of total cost of production (Sanni and Ogundipe, 2005). Possible and most appropriate measure for shortage of animal protein for human consumption lies in the production of fast growing animals like rabbits. This is because livestock like cattle, camels, sheep, goats and pigs take longer time to mature. Rabbits as potential animal with short generation interval, high prolificacy, easy management strategy, ability to utilize waste and other unconventional feed sources for maximum weight gain (efficient feed conversion) to good high quality and quantity animal protein and the ability to thrive on forage with little concentrate will reduced the shortage of protein from animal sources. Rabbit meat is also known to be low in sodium, cholesterol and higher proportion of polyunsaturated linoleic and linolenic fatty acids. All these attributes in addition to less feed cost of production, make rabbit desirable livestock that can help increase and improve protein intake of an average income earning in Nigeria (Mohammed, *et al.*, 2011)

There is an urgent need for low cost feed ingredient especially locally available and cheap agro-industrial products particularly those that do not attract competition in consumption between humans and livestock or have no direct relevance in human food channel. Rumen digesta is one of these wastes that pose a problem to the environment. It is an abattoir by-product obtained from cattle, sheep and goat as well as camel (Maigandi *et al.*, 2004). Magaindi and Tukur (2002) reported the proximate analysis of rumen content for camel, cattle sheep and goat slaughtered at the Sokoto abattoir. In their studies, camel rumen content was reported to have 8.39% crude protein (CP), 11.98% ether extract (EE), and 36.74% (CF) while 8.18% CP, 2.68 EE and 33.31% CF were documented for cattle. Sheep rumen digesta had 9.18% CP, 5.57% EE and 33.31% CF while 8.03 % CP, 5.57% EE and 30.83% CF were values for goat. Sun-dried rumen content of various ruminants animal by-product which is found mostly in abattoirs when the rumen of the animals is cut open after slaughter. The rumen is a unique organ. Its content (rumen content or digesta) is very heterogeneous. It is made up of digested feed at different stages of degradation, saliva (making up the rumen liquor) micro-organisms and the products of their metabolic activities such as proteins, peptides, amino acids, lipids, vitamins and Volatile Fatty Acids (VFA). The aim of this research work is to ascertain higher inclusion levels of rumen content of cattle, sheep and goats on carcass characteristics and economics of production of rabbits.

Materials and Methods

Experimental Site

The experiment was conducted in Katagum Local Government Area of Bauchi State, Nigeria. It is located between latitudes 11° 42' and 11° 40' and longitude 10° 31' and 10° 11' east (Anon, 2009). It shares common boundary with Itas/Gadua local government in north-west, Jama'are to the west, Dambam to the east, Misau to the south-west, Giade to the south and Shira to the southwest (Azare, 2013). It has a landmass of 1,120 square kilometres (NPC, 2009). The climate of the study area is controlled by the intertropical convergent zone (ITCZ) which is marked by the rainy and dry season. The major climate elements that influence the weather of the study area and affecting the farming system were temperature and precipitation (rainfall), the annual temperature ranged between 22-33°C from April to May (Bashir *et al.*, 2001). The mean annual rainfall ranged between 615.6-985mm with peak between July-Augusts.

Sources and feed Processed

Different rumen contents inclusion from caprin, ovine and bovine were obtained from Azare Abattoir and suns dried by constant raking at an intervals of 2hours until dry materials were obtained. Other feed ingredients were procured at Azare main market. The rabbits used in this study were mongrels obtained from the rabbit farmers in Azare. The three types of rumen content were used for formulation of six diets each at 30 and 40% levels. The percentage composition of the experimental diet was shown in table 2. Forty-eight mongrel rabbits were randomly allotted to the six experimental diets. There were eight rabbits per treatment and replicated 4 times (two rabbits per replicate) in a completely randomized design (CRD) using 3x2 factorial arrangements.

Proximate composition of different types of rumen content

The samples of different types of rumen content were taken to biochemistry laboratory, School of Agriculture and Agricultural Technology of Abubakar Tafawa Balewa University Bauchi for proximate analysis. Samples were analyzed for crude protein (CP), crude fibre (CF), ash, crude fat (ether extract) and nitrogen free extract (NFE) accurate to the procedures of AOAC (1990).

Carcass evaluation

At the end of the feeding trial, 4 rabbits were randomly sampled from each treatment, weighed, starved overnight and slaughtered by severing the jugular vein. During evisceration, the internal organs and other gut contents were removed and weight, the dressed carcass and internal organs were weight and expressed as percentage of the live weight.

$$\text{Dressing \%} = \frac{\text{Dressed carcass weight}}{\text{Live weight}} \times 100$$

The economics of production was based on the prevailing market price at Azare as of the time of the experiment. The experiment lasted for six weeks and both the interaction effects as well as main effects were tested. The data set generated was analyzed according to Steel and Torrie 1980 using the Minitab Statistical software. LSD was used to separate the means.

Results and Discussion

Table 1 showed the percentage composition of the different types of rumen content fed to rabbits. The crude protein, crude fibre and ash content ranged of 16.00%-16.06%, 8.34%-17.83% and 3.56%-7.93% were adequate for growing rabbits in the tropics (Aduku, 2004). The proximate composition of different types of rumen content shown in table 2, results showed variation in different types of rumen content for caprine, ovine and bovine CP 15.93%, 12.29% and 11.23%, CF 32.58%, 33.56% and 34.61%, EE 0.86%, 1.15% and 1.23%, Ash 11.21%, 13.42% and 12.54% and NFE 40.85%, 34.42% and 35.59% respectively. The results were similar to those reported by Abubakar (1998), and Dairo (2005). The high CP in caprine rumen content can be attributed to grazing habits of goat which prepare to browse of young shoot, twigs and flowers which are more nutritious than the mature plant. Similarly Mohammed *et al.* (2008) showed a wide variation between rumen content of cattle, camel, sheep and Goat. The effect of interaction of carcass of rabbits fed rumen content is shown in Table 3, results showed that final live weight, carcass weight and dressed weight were not affected by the types and levels of rumen content ($P>0.05$), and this findings partly concord with the findings of Ani *et al.* (2015) who reported non-significant difference in rabbits fed dehydrated rumen content. Most of the gut content of different rumen content fed to rabbits were not affected by the types and levels of rumen content except the head, and small intestine which were affected by interaction effect ($P<0.05$). Olabanji *et al.*, (2007) reported a significant difference in small intestine of rabbits fed rumen content. Similarly Esonu *et al.*, (2006) reported that there were no significant differences in organs of rabbits fed rumen content. The main effects of different types of rumen content are shown in Table 4. Results showed that most of the carcass parameters were not affected by the types of rumen content ($P>0.05$), this results partly agrees with the findings of Okpanchi *et al.*, (2010) who reported that there was no significant difference between carcass yield and organs of rabbits fed sun dried rumen content. The economics of production of rabbits fed different rumen content is shown in table 5. Results showed that the feed cost per kg steadily declines from 30-40% level. Similarly the feed cost per kg gain decreased from 30-40% for the three types of rumen content and this is in line with the earlier reports of Mohammed *et al.*, (2011) who reported decline in cost per kg gain of rabbits fed bovine rumen content. The highest feed cost per kg gain was observed in caprine rumen content (₦181.60) while the lowest feed cost was observed in 40% rumen content of bovine (₦137.75)

Conclusion

Rumen content of cattle sheep and goat can be included up to 40% level in the diet of growing rabbits without adverse effect on carcass characteristics with reduction in cost of production.

References

- Abubakar, M.M.(1998). Utilization of unconventional feedstuff for sustainable livestock production. Abubakar Tafawa Balewa University, Bauchi, Inaugural lectures series NO.9, 44pp
- Aduku, A. O. (2004). **Animal Nutrition in the Tropics**. Davcon and Business bureau, Samaru, Zaria, Nigeria
- Ani, A. O., Udeh, F. U., Uzochukwu, I.E. and Onuora, S.I. (2015). Haematological indices, carcass and organ characteristics of crossbred weaner rabbits fed dehydrated bovine rumen digesta. Proc. 40th Ann. Conf. Nigerian Society for Animal Production, 15-19th march, 2015, NAPRI/ABU, Zaria
- Association of Official Analytical Chemists (AOAC), (1990). Official Methods of Analysis, Washington D.C. USA.
- Anonymous (2009). Mapsofworld. Com. Available at <http://www.mapsofworld.com/Nigeria/cities/azare/html> mapXL inc. 10s third street Suite 310 San Jose.
- Azare, I.M. (2013).Evaluation of farmers response strategies to climate change in Katagum local government area of Bauchi state. *Journal of Environment Technology and Sustainable Agriculture* 2 (10):26-33
- Bashir, M.M., Bala, A., Mohammed, I.T., Isa, H.J., Adamu, M.B., Hamisu, M.S., and Abdullahi, A.(2001).Request for the creation of Katagum state out of the present Bauchi state of Nigeria. A memorandum submitted to the speakers, house of representatives, national assembly, Abuja, Nigeria pp 1-28.
- Cheeke, P.R. (1987) Rabbits feeding and nutrition. Academy Press Inc. London pp 41-51.
- Esonu, B.O., Ogbonna, U.D., Anyanuwn, G.A., Emenalom, O.O., Uchegbu, M.C., Etux, E.B. and Udedibie, A.B.I. (2006) Evaluation of performance, Organs characteristics and Economic Analysis of Broiler finisher fed Dried Rumen Digesta. *Inter. J. Poult. Sci.* 5(12): 1116 – 1118.
- Dairo, F.A.S. and Aina, O.O, (2000) Ruminant content as a protein source in diet of growing rabbits. Paper presented at the 25th Annual Conference of the Nigerian Society for Animal Production, Michael Okpara University of Agriculture, March 2000
- Maigandi, S.A. Adebisi, B.Z. and Manga, S.B. (2004).Chemical composition of Ensiled Forestomach digesta (FSD) from Animals slaughter at Sokoto abattoir *Proceedings of the 29th Annual conference of Nigeria Society of Animal Production*, 29:149-151.
- Maigandi, S. A. and Tukur, H.M. (2002). Potentials of fore-stomach Digesta as an unconventional feed ingredients.*Journal Agriculture and Enviroment*. 3(1): 55-64.

- Mohammed, G., Igwebuike J.U. Ubosi, C.O. and Alade, N.K. (2008). Comparative study of the nutrient composition, Amino acid profile and microbial Assay of fresh and dry cattle, camel, sheep, and goat, rumen contents. Proc. of the 13 Ann. Conf. of .Anim.sci.Assoc. of Nig. (ASAN) Sep 15-19, 2008. p-518-520
- Mohammed, G., Igwebuike, J.U and Alade, N.K. (2011). Performance of growing rabbits fed graded levels of bovine blood-rumen content mixture. Agriculture and Biology Journal of North America 2(4):720.723
- Okpanachi, U., Aribido, S.O. and Daikwo, S.I. (2010). Growth and heamatological response of growing rabbits to diets containing graded levels of Sundried Bovine Rumen content. *Africa Journal of Food and Agriculture Nutrition and Development*. 10(11): 4444 – 4457
- National Population Commission (NPC, 2006).National Population Commission Bulletin. Printed and published by federal government printers, Abuja. FGP
- Olabanji, R. O., Farinu, G. O., Akinlade, J. A. and Ojebiyi, O. O. (2007). Growth Performance, Organ Characteristics and Carcass Quality of Weaner Rabbits Fed Different Levels of Wild Sunflower (*Tithonia diversifolia* Hemsl A. Gray) Leaf-Blood Meal Mixture. *International Journal of Agricultural Research*, 2: 1014-1021.
- Sanni, S.A. and Ogundipe, S.O. (2005). Economics of some modules of poultry production inKaduna state. *Nigerian Journal of Animal Production*, 32(1):102-107.
- Steel, R. G. D. and Torrie, J. H. (1980). Principles and procedures of Statistics: A biometric Approach New York: McGraw Hill

Table 1: Percentage composition of rumen content fed rabbits

Parameters	Goat rumen content		Sheep rumen content		Cattle rumen content	
	30%	40%	30%	40%	30%	40%
Maize	42.33	34.33	38.33	29.00	35.33	25.00
Rumen content	30.00	40.00	30.00	40.00	30.00	40.00
Soybean	14.67	12.66	18.67	18.00	21.67	22.00
Maize offal	10.00	10.00	10.00	10.00	10.00	10.00
Bone meal	2.00	2.00	2.00	2.00	2.00	2.00
Salt	0.50	0.50	0.50	0.50	0.50	0.50
Premix	0.50	0.50	0.50	0.50	0.50	0.50
Total	100.00	100.00	100.00	100.00	100.00	100.00

Calculated analysis (%)

Crude protein	16.01	16.07	16.01	16.00	16.01	16.00
Crude fibre	8.34	13.63	13.63	16.97	14.30	17.83
Ether extract	6.05	7.93	5.78	7.40	3.56	4.48
Ash	6.14	7.93	5.78	7.40	3.56	4.48

Premix supply the following per 2.5 contain vitamin of 15,000,000 I.U., Vitamin D₃ 3,000,000 I.U, Vitamin E 30,000,000, Vitamin K₂ 500mg, Thiamine B₁ 2000mg, Riboflavin B₂ 6000mg, Pyridoxine B₆ 400mg, Niacin 40,000mg, Pantothenic acid 10,000mg, Folic acid 1000mg, Biotin 80mg, Choline chloride 500mg, Biotin 80mg, antioxidant 125g, Manganese, 96g, Zinc 60g, Iodine 1.4g, Selenium 240g, Cobalt 24mg.

Table 2: Proximate composition of the different types of rumen content

Types of rumen content	Parameters					
	DM	CP	CF	EE	Ash	NFE
<i>Caprine</i>	95.43	15.93	32.58	0.86	11.21	40.85
<i>Ovine</i>	95.23	12.29	33.56	1.15	13.42	34.42
<i>Bovine</i>	96.51	11.23	34.61	1.23	12.54	35.59

Table 3: Types and level interaction effect on carcass, organs and gut characteristics (% body weight) of rabbits

Diets									
Parameters		<i>Caprine</i>		<i>Ovine</i>		<i>Bovine</i>		SEM	LSD
		30%	40%	30%	40%	30%	40%		
Final live weight (g)		1293.30	1200.00	1075.00	1130.00	1368.80	1487.50	126.74	NS
Carcass weight (g)		593.75	456.25	531.25	668.75	718.75	775.00	64.13	NS
Dressed weight %		44.97	38.36	50.08	51.75	52.25	52.86	3.17	NS
Liver %		2.52	2.50	3.25	2.35	2.29	2.77	0.31	NS
Lungs %		0.63 ^b	0.54 ^b	1.34 ^a	0.51 ^b	0.42 ^b	0.60	0.19	0.46
Heart %		0.21	0.13	0.19	0.17	0.15	0.13	0.03	NS
Kidney %		0.53	0.56	0.59	0.50	0.16	0.67	0.25	NS
Abdominal fat %		0.53	0.56	0.59	0.50	0.16	0.67	0.25	NS
Head %		8.23 ^b	8.48 ^b	9.25 ^a	7.73 ^b	7.72 ^b	7.95 ^b	0.38	0.90
Feet %		2.19	2.57	2.66	2.12	2.06	1.93	0.19	NS
Pelt %		7.29	6.15	6.68	6.87	5.92	5.80	0.48	NS
Stomach %		1.39	1.60	1.87	1.62	1.48	1.36	0.12	NS
Small intestine %		3.88	3.50	5.42	3.50	4.35	4.38	0.36	NS
Large intestine %		0.96	0.89	1.13	0.77	1.05	1.32	0.23	NS
Caecum %		2.75	2.32	3.04	2.75	2.50	2.26	0.24	NS

Note: Means bearing different superscripts are statistically different ($P < 0.05$), NS= Not significant

Table 4: Main effects of rumen content type and levels on carcass, organs and gut characteristics (% body weight) of rabbits

Parameter	Diets			LSD	30%	40%	LSD
	Caprine	Ovine	Bovine				
Final live weight (g)	1246.90	1187.50	1428.10	NS	1245.80	1329.20	NS
Carcass weight (g)	525.00	600.00	746.87	157.60**	614.58	633.33	NS
Dressed weight %	41.67 ^a	50.92 ^b	52.55 ^c	7.76**	49.10	47.66	NS
Liver %	2.51	2.80	2.54	NS	2.69	2.54	NS
Lungs %	0.59	0.93	0.53	NS	0.79	0.56	NS
Heart %	0.17	0.18	0.14	NS	0.18	0.14	NS
Kidney %	0.44 ^a	0.64 ^c	0.48 ^b	0.15	0.57	0.49	*
Abdominal fat %	0.45	0.55	0.41	NS	0.43	0.51	NS
Head %	8.35	8.49	7.83	NS	8.39	8.05	NS
Feet %	2.88	2.39	1.99	NS	2.31	2.21	NS
Pelt %	6.72	6.78	5.86	NS	6.63	6.27	NS
Stomach %	1.49	1.75	1.42	NS	1.58	1.53	*
Small intestine %	3.69	4.46	4.36	NS	4.57	3.79	*
Large intestine %	0.93	0.95	1.19	NS	1.05	0.99	NS
Caecum %	2.54	2.89	2.38	NS	2.76	2.44	NS

Note: Means bearing different superscripts are statistically different ($P < 0.05$), NS= Not significant

Table 5: Economics of production of rabbits fed rumen content based diet

Parameters	Diets					
	Caprine 30%	40%	Ovine 30%	40%	Bovine 30%	40%
Total feed intake (kg)	2.59	3.29	2.52	2.75	2.64	2.57
Feed cost (₦/kg)	24.54	21.23	25.77	22.87	26.61	24.12
Total feed cost (₦)	63.68	69.85	64.94	62.89	70.25	61.99
Total weight gain (kg)	0.35	0.45	0.37	0.39	0.42	0.45
Feed cost (₦/kg) gain	181.60	155.22	175.51	161.26	167.26	137.75