



NSAP

**47th Annual
Conference
(JOS 2022)**

**CONFERENCE
PROCEEDINGS**

THEME
**SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES**

**GROWTH PERFORMANCE OF TURKEY (*Meleagris gallopavo*) POULTS FED BUTYRIC ACID
SUPPLEMENTED DIETS AS AN ALTERNATIVE TO ANTIBIOTICS GROWTH
PROMOTERS**

*¹Mbakwe, C. C., ²Agboola, A. F. and ²Omidwura, B. R. O

*¹Department of Animal Science, Faculty of Agriculture,
Federal University Dutsin-ma, Kastina State.

²Department of Animal Science, Faculty of Agriculture, University of Ibadan, Oyo State

*Corresponding author: frimbia75@gmail.com. +2349035827553

Abstract

The effects of butyric acid supplemented diets on growth performance as an alternative to antibiotic growth promoters was examined in a 28-day experiment using one hundred and sixty (160) (one-day old)



Nicholas turkey poult. The experimental design was a Completely Randomised Block Design. Poults were brooded for 7 days, after which they were allotted to 4 dietary treatments with 5 replicates of 8 birds per replicate. The experimental feeds consist of four diets; treatment 1 was the Corn-SBM diet (basal diet/negative control); treatment 2 was the positive control, consisting of the basal diet and antibiotics (Oxytetracycline hydrochloride at 0.02g/1000g feed), treatment 3 had the basal diet and 0.2% butyric acid, while treatment 4 was the basal diet and 0.4% butyric acid. The inclusion of butyric acid to the corn-soya based diets did not positively affect the final weight and weight gain of birds at the end of the first 28 days of the feeding trial.

Keywords: Butyric acid, Growth performance, Turkey poult

Introduction

The continued indiscriminate use of antibiotics in both man and birds poses a challenge of antibiotic resistance caused by antibiotic residues in meat and milk. This led to a ban on antibiotics use in feed formulation in Europe (Maron *et al.*, 2013). This ban has brought about the need for researchers to find alternatives to the use of antibiotics in poultry production. Short-chain fatty acids (acetic, butyric and propionic acids) have been reported (Agboola *et al.*, 2015; Agboola *et al.*, 2018) to have a positive impact on broiler growth (Hassan *et al.*, 2010). It has been documented that these acids exhibit antimicrobial properties when dissociated by reducing intestinal pH (Van Immerseel *et al.*, 2006). Notwithstanding the progress made, there has been diverse recommended levels of organic acids that would successfully replace antibiotics. Therefore, the purpose of this study is to determine the potency of butyric acid and to establish a specific level of inclusion that will provide optimum performance and improve the integrity of the gut in turkey poult.

Materials and methods

Experimental site

This study was carried out in the Poultry Unit of the Teaching and Research Farm, University of Ibadan, Oyo State in the South West geopolitical zone of Nigeria, within the tropical rain forest region.

Experimental birds, design and management

One hundred and sixty (160) one-day old unsexed Nicholas turkey poult were obtained from a reputable commercial hatchery and kept in a deep litter system. The poult were mass brooded for seven (7) days during which they were fed the negative control (basal diet). After which they were weighed, tagged and randomly allotted to four dietary treatments. Each dietary treatment had five (5) replicates of eight (8) poult per replicate in a Completely Randomised Design (CRD). The birds were reared in a well ventilated and illuminated standard poultry house.

Experimental diets and feeding

The experimental feeds consist of four diets; treatment 1 was the Corn-SBM diet (basal diet/negative control); treatment 2 was the positive control, consisting of the basal diet and antibiotics (Oxytetracycline hydrochloride at 0.02g/1000g feed); treatment 3 had the basal diet and 0.2% butyric acid, while treatment 4 was the basal diet and 0.4% butyric acid. The basal diet was a corn-soyabean diet formulated to meet the nutrient requirements (NRC, 1994) for starter phase 1 (0 to 4 weeks) phases. Experimental diets and water were provided *ad libitum* in a feeding trial that lasted for 4 weeks.

Table 1: Gross Composition (g/1000gDM) of experimental diets

INGREDIENTS g/kg	Starter 1 (0-4) weeks
------------------	-----------------------



NSAP

47th Annual Conference
(JOS 2022)

CONFERENCE PROCEEDINGS

THEME
SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES

Corn	410.0
Soyabean meal	480.0
Fish meal	40.0
Wheat Offal	6.0
Dicalcium phosphate	20.0
Premixes	2.5
Limestone	10.0
DL-methionine	2.5
L-lysine	3.5
Salt	2.5
Butyric acid	0.0
Antibiotics	0.0
Soya oil	23.0
TOTAL	1000.0

Calculated Nutrients

Crude protein g/kg	260.12
ME Kcal/kg	2811.23
Fat g/kg	35.264
Crude fibre g/kg	43.41
Calcium g/kg	11.6283
Total phosphorus g/kg	8.9417
Non-phytate P, g/kg	5.0627
Ca:NPP	2.298083

Composition of premix per kg of diet: Vitamin A, 12500 I.U; vitamin D3, 255000 I.U; vitamin K3, 2mg; vitamin B1, 3mg; vitamin B2, 5.5mg; calcium pantothenate, 11.5mg; vitamin B12, 0.025mg; choline, chloride, 500mg; folic acid, 1mg; biotin, 0.08mg; manganese, 120mg; iron, 100mg; zinc, 80mg; copper, 8.5mg; iodine, 1.15mg; cobalt, 0.3mg; selenium, 0.12mg; anti-oxidant, 120mg. Over the top inclusion of 0.2, 2 and 4 were added for antibiotics, 0.2% butyric acid and 0.4% butyric acid respectively.

Parameters measured and calculated

The growth performance parameters calculated include feed intake and body weight gain. Feed conversion efficiency was calculated using the weight gain and feed intake values.

Feed intake

The weekly feed intake was determined by giving a known quantity of feed to the birds and collecting the leftovers. Feed remnants were weighed and subtracted from the initial weight of feed offered to determine the feed intake per replicate per week.

Body weights

The chicks were weighed at the beginning of the trial and subsequently weighed on weekly basis throughout duration of the experiment. The weekly weight gain was obtained by subtracting the weight of the preceding week from that of the present.

Feed Conversion Ratio

**NSAP****47th Annual
Conference
(JOS 2022)****CONFERENCE
PROCEEDINGS****THEME
SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES**

This was calculated per pen as the ratio of weight gained to the feed consumed.

$$\text{Feed Conversion Ratio (FCR)} = \frac{\text{Average feed intake(g)}}{\text{Average weight gain(g)}}$$

Data analysis

Samples collected were subjected to Analysis of Variance using (SAS,2008). Mean differences was separated using Duncan's Multiple Range Test

Results and Discussion

The result on the influence of butyric acid supplementation on the performance of turkey poult as shown in Table 1, initial weight of birds on antibiotic supplemented diet was ($p < 0.05$) significantly higher than the control, 0.2 and 0.4% butyric acid supplemented diets. Final weight and weight gain of birds on the control and antibiotic supplemented diet were similar ($p < 0.05$) and significantly higher than those on 0.2 and 0.4% butyric acid supplemented diets. Birds on the control, antibiotic supplemented and 0.2% butyric acid supplemented diet consumed more feed than those on 0.4% butyric acid supplemented diet

Table 1: Effect of butyric acid supplementation on performance and cost analysis of turkey poult (1 – 28 days)

Parameters	T1	T2	T3	T4	SEM
Initial weight (g/bird)	43.48 ^b	44.97 ^a	43.89 ^b	43.06 ^b	0.44
Final weight (g/bird)	350.27 ^a	360.07 ^a	315.41 ^b	309.71 ^b	11.19
Weight gain (g/bird)	306.79 ^a	315.09 ^a	271.52 ^b	266.65 ^b	11.12
Feed intake (g/bird)	734.58 ^a	746.07 ^a	694.76 ^a	629.64 ^b	31.25
FCR	2.40	2.37	2.56	2.36	0.15

^{a, b} Means within the same row with the same superscripts are not significantly different ($p > 0.05$). T1 = Control (basal diet); T2 = basal diet + antibiotics fed for 21 days; T3 = basal diet + 0.2% butyric acid fed for 21 days; T4=basal diet+ 0.4% FCR=Feed Conversion ratio. SEM =Standard Error of Mean

The results of this study showed that the inclusion of butyric acid to the corn-soya based diets did not positively affect the final weight and weight gain of birds at the end of the 28 days of the feeding trial. This agrees with the work of Adil *et al* (2011) who reported no significant body weight gain difference between the control diet and butyric acid (2 and 3 %) supplemented diets in broiler chickens. The result of this experiment is however, contrary to the findings of Panda *et al* (2009) and Taherpour *et al* (2009). Panda *et al* (2009) reported that butyric acid inclusion at 0.4 and 0.6% produced significant increase in body weight gain when compared to the control and 0.2 butyric acid inclusion level while Taherpour *et al* (2009) reported that a supplementation with butyrate acid glycerides showed an increase in final body weight (42 days) of broiler chickens when compared to those on the basal diet. However, the authors also reported a lower feed intake and feed conversion ratio for birds on the butyric acid supplemented diet when compared to those on supplement-free diet. Agboola *et al* (2015) reported an improved weight gain at 21 days when 0.4 % organic acid (formic and propionic acids) were added to the feed of broiler chicken.

Conclusion and recommendation

The result obtained from this study showed that butyric acid at the inclusion level up to 0.4 % had no detrimental effects on growth performance of turkey poult (0-4 weeks) however, further research is needed to better ascertain the best inclusion level in turkey nutrition for optimal productivity. Also, varying levels of butyric acid may be explored and further work can be carried out to finisher phase.

**NSAP****47th Annual
Conference
(JOS 2022)****CONFERENCE
PROCEEDINGS**THEME
**SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES****References**

- Adil, S., Banday, M. T., Bhat, G. A., Qureshi, S. D. and Wani, S. A. (2011). Effect of supplemental s acids on growth performance and gut microbial population of broiler chicken. *Livestock Research for Rural Development* 23 (1): 1-8.
- Agboola, A. F., Omidwura, B. R. O., Odu, O., Popoola, I.O., and Iyayi, E. A. (2015). Effects of organic acid and probiotic on performance and gut morphology in broiler chickens. *South African Journal of Animal Science* 45 (No. 5):494-501
- Agboola, A. F., Omidwura, B. R. O., Osakue, P. O., and Ogunbiyi, K. O., (2018). Effects of Sodium Acetate and Sodium Propionate Supplemented Diets on Growth Performance and Gut Histomorphology of Broiler Starters. *Nigerian Journal of Animal Science* 20.2:183–196.
- Hassan, H. M. A., Mohamed, M. A., Youssef, A. W. and Hassan, E. R. (2010). Effect of using organic acids to substitute antibiotic growth promoters on performance and intestinal microflora of broilers. *Asian Aus Journal of Animal Science*. 23 (10):1348-1353
- Maron, D.F., Smith, T.J. and Nachman, K.E. (2013). Restrictions on antimicrobial use in food animal production: an international regulatory and economic survey. *Global Health* 9,48.
- National Research Council (1994). Nutrient Requirement of Poultry: Ninth Revised Edition. Washington, DC: The National Academies
- Panda, A. K., Raju, M. V., Rama, L. N., Rao. S.V., Shyam S. G., Reddy, M.R. (2009). Effect of graded levels of formic acid on gut microflora count, serum biochemical parameters, performance, and carcass yield of broiler chickens. *Indian Journal of Animal Science*; 79:1165-1168.
- SAS (2008). Statistical Analysis System, SAS users guide: statistics. SAS institute Inc. Cary, N.C. USA
- Taherpour, K., Moravej, H., Shivazad, M., Adibmoradi, M., and Yakhchali, B. (2009). Effects of dietary probiotic, prebiotic and butyric acid glycerides on performance and serum composition in broiler chickens. *African Journal of Biotechnology*. 8 (10); 2329-2334.
- Van Immerseel F., Russell J. B., Flythe M. D., Gantois I., Timbermont L., Pasmans F., Haesebrouck F. and Ducatelle, R. (2006). The use of organic acids to combat salmonella in poultry: a mechanistic explanation of the efficacy. *Avian Pathol*. 35:182-188.

**EFFECT OF COOKING DURATION ON PROXIMATE COMPOSITION AND PHYTOCHEMICAL CONTENTS OF
KIDNEY BEAN SEED (*PHASEOLUS VULGARIS*)**

R.J.Thani, S.E.Alu, I. K. Owuna and A.Yakubu

Nasarawa state, Department of Animal Science, Faculty of Agriculture, Nasarawa State University Keffi,
Shabu- Lafia Campus, Nasarawa State, Nigeria.

E- mail: rahabthani3@gmail.com

Abstract

This study aimed at evaluating the effect of cooking duration on proximate composition and phytochemical contents of kidney bean seed (Phaseolus vulgaris). The seeds were cleaned and extraneous materials like dry leaves, stones, and dirt were removed and divided into five portions. They