

**NSAP****47<sup>th</sup> Annual Conference**  
(JOS 2022)**CONFERENCE PROCEEDINGS**THEME  
**SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES****EFFECTS OF PLANTING METHODS AND FERTILIZER APPLICATION RATES ON PROXIMATE COMPOSITION OF *CENTROSEMA PASCUORUM* IN DUTSE LOCAL GOVERNMENT AREA OF JIGAWA STATE****\*Gumel, I. A<sup>1</sup>., Mohammed, J<sup>2</sup>.,Abdurrahman, S.L<sup>1</sup>., Babandi, B<sup>3</sup>., Kiri I. Z<sup>4</sup>., Bukar, A.I<sup>1</sup>., Umar, A.M<sup>1</sup>., Umar, A.M<sup>3</sup>., Umar, Z<sup>5</sup>., Danzabuwa, A.I<sup>6</sup>.,Tajuddeen, I<sup>3</sup>., Babangida, L<sup>1</sup>.,Naziru, B<sup>1</sup>., Murtala, I<sup>7</sup>. and Umar M.I<sup>8</sup>**<sup>1</sup>Department of Animal Science Federal University, Dutse Jigawa State P.M.B 7156<sup>2</sup>Government Science and Technical College Nguru, Yobe State P.M.B 150<sup>3</sup>Department of Animal Health and Production Binyaminu Usman polytechnic, Hadejia P.M.B 103<sup>4</sup>Department of Biology,Sule Lamido University,Kafin Hausa PMB 048 Jigawa State, Nigeria<sup>5</sup>Department of Biology Jigawa State College of Education Gumel P.M.B 1002<sup>6</sup>Department of Animal Science Federal University Dutsin-ma P.M.B 5001<sup>7</sup>Jigawa State Ministry of education Science and Technology, Dutse.<sup>8</sup>Jigawa State Agricultural and Rural development Authority,Dutse**\*Corresponding Author**[abdullahiibrahim887@gmail.com](mailto:abdullahiibrahim887@gmail.com) +2348034695413**Abstract**

A study was conducted to evaluate effects of planting methods and fertilizer application rates on proximate composition of *Centrosema pascuorum*, at experimental field of crop science Department, Faculty of Agriculture Federal University Dutse, Jigawa State. The research was laid out in 2x5 factorial arrangement in a randomized completely block design (RCBD) consisting of two (2) methods of planting (Drilling and Broadcasting) and five (5) fertilizer rate (0kg/ha<sup>-1</sup>, 150kg/ha<sup>-1</sup>, 200kg/ha<sup>-1</sup>, 250kg/ha<sup>-1</sup> and 300kg/ha<sup>-1</sup>). Fertilizer used was single super phosphate (SSP). Proximate components were all determined. Result revealed that Proximate composition had no effect on planting method, ( $P>0.05$ ) numerically Moisture content and Ash (17.34) (15.15) was higher in drilling method compared to broadcasting. However, broadcasting method numerically NFE was higher (29.89). Effect of fertilizer on proximate composition Moisture content, Dry matter, Crude fibre, and Ash was significant ( $P<0.05$ ).at 150 and 300kg/ha fertilizer. The effect of interaction between planting method and fertilizer application rates Crude fibre was highly significant ( $P<0.05$ ) at 150 kg/ha fertilizer rates drilling (23.96), broadcasting (24.72) 150kg/ha,) and (25.23)300kg/ha respectively. The crude protein was numerically higher in drilling method (17.01) at 250kg/ha fertilizer rates.Based on the results obtained on the proximate composition of *C. pascuorum* in this study, it can be recommended that any farmer intending to go into pasture production particularly production of *C. pascuorum* can adopt any of the two levels of fertilizer (150 and 300kg/ha fertilizer) for higher dry matter, crude fibre and ash. 250kg/ha fertilizer for drilling can be recommended for crude protein from this study.

**Keywords:** Broadcasting method, *Centrosema pascuorum*, Drilling method, Fertilizer application and proximate composition

**Introduction**

Nutrition is one of the important management practices in ruminant production and it is the bed rock of performance in animal, but if there are limited forages; and reduction in nutrient composition of the available forage grasses and legumes during dry season, this poses a problem (Amole *et al.*, 2011). the major challenge to livestock production in Nigeria is ensuring adequate feed supply throughout the year in terms of quality and quantity in the traditional setting, where animals are predominantly raised on natural grasses (Kallah *et al.*1997). There are many pasture crops used in feeding animals made up of grasses and legumes. Among the forage is *Centrosema pascuorum* (centurion). *C. pascuorum* is an annual twining leguminous herb that can root from the node of trailing herb under moist condition. *C. pascuorum* is suitable for mixture with grasses or pure stands for lay-farming and adapted to wide range of soil ranging from sand to heavy clay and from slightly acidic to alkaline

**NSAP****47<sup>th</sup> Annual Conference**  
(JOS 2022)**CONFERENCE PROCEEDINGS**THEME  
**SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES**

soil pH (5 – 8.5). It is adapted to tropical areas with pronounced dry season up to 8 months (Feedipedia, 2019). In grassland farming, the *C. pascuorum* phase is able to provide 80-100 kg/ha N to the system, (Thiagalingam *et al.*, 1997). However, the study aims to determine the effect of planting methods and fertilizer rate on proximate composition of *C. pascuorum*..

## **Materials and methods**

### **Experimental sites**

The experiment was conducted at the experimental field of Crop Science Department, Faculty of Agriculture, Federal University Dutse, Jigawa state of Nigeria. Located between latitude 11.00°N to 13.00°N and longitudes 8.00°E to 10.15°E High temperature are normally recorded between the month of April and September with average temperature range of 15°C and 35°C. Raining season last from May to September with average rainfall of between 600mm to 1000mm. The ecological zone is sudan savannah vegetation which is made up vast grazing land suitable for livestock production and free from tsetse fly infestation. The environment is conducive to different species of livestock production, such as cattle, sheep, goats, Rabbits donkeys, horses and poultry (JARDA, 2012).

### **Materials use in the research**

The seeds of *C. pascuorum* were purchased at National Animal Production Research Institute (NAPRI) Zaria. The single super phosphate fertilizer was used. Weighing scale and tape were used for the weighing and measuring purposes respectively.

### **Experimental design**

The research was laid out in 2x5 factorial arrangement in randomize completely block design (RCBD) consisting of two (2) methods of planting (Drilling and Broadcasting) and five (5) fertilizer rate (0kg/ha<sup>-1</sup>, 150kg/ha<sup>-1</sup>, 200kg/ha<sup>-1</sup>, 250kg/ha<sup>-1</sup> and 300kg/ha<sup>-1</sup>). A total land area of (120m<sup>2</sup>) was used, the field was prepared by ploughing and harrowing to soften and enhance early plant germination. The experimental field was 2mx2m each up to 30 plots, inter-row and intra-row of 0.5m, 1m was used respectively.

### **Chemical Analyses**

Harvesting was carried out at 8 weeks post sowing, materials were then oven dried at 65°C for 72 h, dried samples was hammer (comprising leaves and stem) to pass through 2 mm sieve, the ground sample was used for proximate composition (DM, ash, CP, CF, EE and NFE) according to the procedures described by AOAC (2005).

### **Data analysis**

Data collected were subjected to analysis of variance (ANOVA) using System Analytical Statistics package (SAS) package and differences between mean was separated using Duncan Multiple Range at probability level of 0.05 (SAS,2009)

### **Results and discussion**

Table1. Presents proximate composition of *C. pascuorum*, had no effect on planting method, ( $P>0.05$ ) numerically Moisture content and Ash (17.34) (15.15) was higher in drilling method compared to broadcasting. However, broadcasting method in numerically NFE was higher (29.89)

The effect fertilizer on proximate composition of *C. pascuorum* is presented Table 2. Show that Moisture content, Dry matter, Crude fibre, and Ash was significant ( $P<0.05$ ).at 150 and 300kg/ha fertilizer. The significant difference observed in proximate composition of *C pascuorum* as may be influence by varying levels of fertilizer applied.

Table 3.shows the effect of interaction between planting method and fertilizer application rates Crude fibre was highly significant ( $P<0.05$ ) at 150 kg/ha fertilizer rates drilling method (23.96), broadcasting method (24.72) 150kg/ha,) and (25.23)300kg/ha respectively. The crude protein was numerically higher in drilling method (17.01) at 250kg/ha fertilizer rates.

This finding is contrary with the work of (Baba, M., *et al.*, (2018) on “Dry matter yield and proximate composition of grass legume mixtures as influenced by sowing date” who reported that the CP, NFE, and CF were highly significant at ( $P<0.05$ ) level of significance. *C. pascuorum* on broadcasting method, CF was highly



significant at ( $P < 0.05$ ) among the crude fibre with 25.23 and 24.27% at 300 1nd 150 kg/ha of fertilizer level. However, The CP content recorded for this study was greater than  $< 7$  to 8 % CP suggested as threshold for sufficient utilization of feed by (Lamidi and; MacDonald 1995; Osarobundo, 2020). Therefore, *C. pascuorum* would provide the adequate nitrogen requirement for the rumen microorganisms to maximally digest the main components of dietary fibre leading to the production of volatile fatty acids (Trevaskis et al 2001; Lamidi and Ogunkunle 2016) which in turn assist microbial protein synthesis is Lamidi and Aina (2013). The dry matter in Table 3. ranged between 79.84 and 84.24 g/100 g in *C. pascuorum* this finding is agreed with finding of (Lamidi and; and Osarobundo, 2020). Who reported dry matter value of 87.66 and 90.79% *C. pascuorum* at four different cutting regimes in humid respectively.

**Table 1: The effect planting methods on proximate composition of *Centrosema pascuorum***

| Planting methods    | MC%                | DM%   | CP%   | CF%   | OIL% | ASH%               | NFE%               |
|---------------------|--------------------|-------|-------|-------|------|--------------------|--------------------|
| Drilling method     | 17.34 <sup>a</sup> | 81.71 | 16.75 | 23.07 | 1.54 | 15.15 <sup>a</sup> | 27.46 <sup>b</sup> |
| Broadcasting method | 15.55 <sup>b</sup> | 82.67 | 16.54 | 22.40 | 1.71 | 13.76 <sup>b</sup> | 29.89 <sup>a</sup> |
| SEM                 | 0.481              | 0.80  | 0.18  | 0.43  | 0.12 | 0.481              | 1.01               |
| LOS                 | NS                 | NS    | NS    | NS    | NS   | NS                 | NS                 |

<sup>ab</sup>= mean values with different superscripts along the same column are significant ( $P < 0.05$ ) MC=Moisture content, DM=Dry matter, CP= Crude protein, CF= Crude fibre, NFE= Nitrogen free extracts, SEM=Standard error mean, LOS= Level of significance. NS= Not significant

**Table 2: The effect of fertilizer on proximate composition of *Centrosema pascuorum***

| Fertilizer rate (kg/ha) | MC%                 | DM%                 | CP%   | CF%                 | OIL% | ASH%                | NFE%               |
|-------------------------|---------------------|---------------------|-------|---------------------|------|---------------------|--------------------|
| 0                       | 17.16 <sup>a</sup>  | 82.90 <sup>a</sup>  | 16.37 | 21.49 <sup>c</sup>  | 1.43 | 13.91 <sup>c</sup>  | 30.14 <sup>a</sup> |
| 150                     | 16.61 <sup>ab</sup> | 82.04 <sup>ab</sup> | 16.51 | 24.11 <sup>ab</sup> | 1.64 | 14.15 <sup>bc</sup> | 26.56 <sup>b</sup> |
| 200                     | 15.68 <sup>c</sup>  | 81.08 <sup>bc</sup> | 16.60 | 21.96 <sup>bc</sup> | 1.72 | 14.21 <sup>b</sup>  | 27.68 <sup>a</sup> |
| 250                     | 15.81 <sup>bc</sup> | 81.93 <sup>b</sup>  | 16.90 | 21.98 <sup>b</sup>  | 1.70 | 14.88 <sup>b</sup>  | 29.04 <sup>a</sup> |
| 300                     | 16.98 <sup>ab</sup> | 82.99 <sup>a</sup>  | 16.83 | 24.12 <sup>a</sup>  | 1.64 | 15.13 <sup>a</sup>  | 29.96 <sup>a</sup> |
| SEM                     | 1.09                | 1.26                | 0.28  | 0.68                | 0.18 | 0.76                | 1.60               |
| LOS                     | *                   | *                   | NS    | *                   | NS   | *                   | NS                 |

<sup>ab</sup>= mean values with different superscripts along the same column are significant ( $P < 0.05$ ) MC=Moisture content, DM=Dry matter, CP= Crude protein, CF= Crude fibre, NFE= Nitrogen free extracts, SEM=Standard error mean, LOS= Level of significance. NS=Not significant \* significant

**Table 3: The effect of interaction of planting methods and fertilizer rate on proximate composition (g/100 g) of *Centrosema pascuorum***

| Planting methods    | Fertilizer rate (kg/ha) | MC%   | DM%                | CP%   | CF%                 | OIL% | ASH%  | NFE%  |
|---------------------|-------------------------|-------|--------------------|-------|---------------------|------|-------|-------|
| Drilling method     | 0                       | 16.12 | 83.86 <sup>a</sup> | 16.30 | 22.76 <sup>bc</sup> | 1.26 | 14.19 | 30.29 |
|                     | 150                     | 18.42 | 79.84 <sup>b</sup> | 16.69 | 23.96 <sup>ab</sup> | 1.62 | 15.45 | 23.97 |
|                     | 200                     | 16.22 | 81.03 <sup>a</sup> | 16.80 | 22.70 <sup>bc</sup> | 1.82 | 15.32 | 26.62 |
|                     | 250                     | 16.84 | 81.90 <sup>a</sup> | 17.01 | 22.92 <sup>bc</sup> | 1.51 | 15.57 | 27.48 |
|                     | 300                     | 19.04 | 81.93 <sup>a</sup> | 16.93 | 23.01 <sup>bc</sup> | 1.51 | 15.21 | 28.96 |
| Broadcasting method | 0                       | 18.12 | 81.95 <sup>a</sup> | 16.44 | 20.23 <sup>d</sup>  | 1.60 | 13.63 | 29.98 |
|                     | 150                     | 14.80 | 84.24 <sup>b</sup> | 16.33 | 24.27 <sup>ab</sup> | 1.66 | 12.85 | 29.14 |
|                     | 200                     | 15.13 | 81.14 <sup>a</sup> | 16.39 | 21.22 <sup>cd</sup> | 1.62 | 13.10 | 28.74 |

**NSAP****47<sup>th</sup> Annual  
Conference  
(JOS 2022)****CONFERENCE  
PROCEEDINGS**THEME  
**SECURING ANIMAL  
AGRICULTURE AMIDST  
GLOBAL CHALLENGES**

|     |     |       |                    |       |                     |       |       |       |
|-----|-----|-------|--------------------|-------|---------------------|-------|-------|-------|
|     | 250 | 14.79 | 81.97 <sup>b</sup> | 16.80 | 21.04 <sup>cd</sup> | 1.90  | 14.18 | 30.61 |
|     | 300 | 14.93 | 84.05 <sup>b</sup> | 16.72 | 25.23 <sup>a</sup>  | 1.763 | 15.04 | 30.96 |
| SEM |     | 1.546 | 1.785              | 0.40  | 0.960               | 0.26  | 1.075 | 2.263 |
| LOS |     | NS    | NS                 | NS    | **                  | NS    | NS    | NS    |

abcd= mean values with different superscripts along the same column are significant (P<0.05) MC=Moisture content, DM=Dry matter, CP= Crude protein, CF= Crude fibre, NFE= Nitrogen free extracts, SEM=Standard error mean, LOS= Level of significance.

## 11.5 Conclusion

The study concluded that planting methods has no effect on proximate composition of *C. pascuorum* However; fertilizer and interaction of planting method and fertilizer affect proximate composition of *C. pascuorum*.

## 11.6 Recommendations

.Based on the results obtained on the proximate composition of *C. pascuorum* in this study, it can be recommended that any farmer intending to go into pasture production particularly production of *C. pascuorum* can adopt any of the two levels of fertilizer (150 and 300kg/ha fertilizer application) for higher dry matter, crude fibre and ash. 250kg/ha fertilizer for drilling can be recommended for optimum crude protein from this study.

## References

- AOAC. (2005). Association of Official Analytical Chemists. Official Methods of Analysis 18th Edition AOAC Inc, Arlington, Virginia USA. 1094pp.
- Amole, T.A., Ojo, V.O.A., Dele, P.A., Adeoye, S.A., Onifade O.S., Jolasho, A.O., Olanite, J.A and Oyewole, S. T. (2011). Nutrient Digestibility of Calves Fed Maize - Lablab Silage Supplement 2011, Proceeding 36th Conference, Nigeria Society for Animal Production. 13-16 March. Pp. 485-487.
- Baba, M., Sani, A.A, Ibrahim, Muhammad, I.R., (2018), "Dry matter yield and chemical composition of grass legume mixtures as influenced by sowing date" .pp: 136
- Bamikole, M. A. (1998). Performance of goats fed *Panicum maximum* and *Stylosanthes hamata* mixture and *Panicum sward* with or without N-fertilizer. Ph.D Thesis, University of Ibadan, Nigeria.
- Ettu, R. O., and Mako, A. A. (2020). Evaluation of nutrient content and in Vitro gas fermentation of some crop residues as feed supplements for ruminants. *ADAN Journal of Agriculture*, 1 (1): 123-130
- Feedipedia (2019) Online encyclopedia of animal feeds Retrieved from <http://www.feedipedia.org/node/16072> on 16/03/2019
- JARDA. (2012). Jigawa State Agricultural and Rural development Authority metrological Station Report: Temperature and Rainfall Record Book and management. Pp. 1-2.
- Lamidi, A. A. and Osarobundo, E. N. (2020). Growth indices, dry matter yield and nutritive value of *Centrosema pascuorum* at four different cutting regimes in humid tropics. *Nigerian Journal of Animal Science* 2020 Vol 22 (3): 128-138 (ISSN:1119-4308)
- Lamidi, A.A., and Aina, A. B. J. (2013). Presowing treatment, agronomic parameters and nutritive potential of *Tephrosia bracteolata* (Guill. Et Perr.) at four different stages of growth. *African Journal of Livestock Extension*. 11:31-36.
- Lamidi, A.A., and T. Ogunkunle. (2016). Nutritional potential of poultry dropping meal as feed resources for ruminant production in Niger Delta. Nigeria. *Global Agricultural Science & Technology*. 1:1-11.
- MacDonald, P., Edwards, R.A. Greenhalgh, J. F. D. and Mergan, C.A. (1995). Animal Nutrition. EIBS Longman, London and New York. Pp. 155-159, 524
- SAS. (2009) Statistical Analysis System Institute, Inc.Cary, North Carolina,USA
- Thiagalingam, K., Zuill, D. and Price, T. (1997) A review of *Centrosema pascuorum* (centurion) cvv. Cavalcade and Bunday as a pasture legume in the ley farming system studies in North West Australia.



# NSAP

**47<sup>th</sup>** Annual  
Conference  
(JOS 2022)

## CONFERENCE PROCEEDINGS

THEME  
SECURING ANIMAL  
AGRICULTURE AMIDST  
GLOBAL CHALLENGES

Proceeding, XVIII International Grassland Congress, Winnipeg-Saskatoon, Canada. Vol. 1:10, 43-44.

Trevaskis, L.M., Fulkerson, W.J. and Gooden, J.M. (2001). Provision of Certain Carbohydrate based Supplement to Pasture fed Sheep, As Well as Time of Harvesting of the Pasture, Influences pH, Ammonia Concentration and Microbial Protein Synthesis in The Rumen. Australian Journal Experimental Agriculture. 41:21- 27