

**NSAP****47th Annual
Conference
(JOS 2022)****CONFERENCE
PROCEEDINGS**THEME
**SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES****RELATIONSHIP BETWEEN BODY WEIGHT AND LINEAR BODY
MEASUREMENTS IN MATURED MUSCOVY AND KHAKI CAMBELL DUCKS****Abdu, I¹., Salisu. A, D¹., Yahaya B²., Jafar³, A. M. and Shuaibu, A⁴.**¹Department of Animal Production, Abubakar Tafawa Balewa University, Bauchi²Department Animal production Technology, Bauchi State Collage of Agriculture, Bauchi³Department of Animal Science, Bauchi State University, Gadau⁴Department of Animal Science, Federal University Kashere**Corresponding author:** iabdu@atbu.edu.ng**ABSTRACT**

The study was conducted at Bauchi local government area of Bauchi state, Nigeria to determine the relationship between body weight and body linear measurements in two breeds of ducks. Data on 150 ducks comprising 78 (fourty male and thirty-eight female) Muscovy and 72 (thirty-two male and fourty female) Khaki Campbell were obtained and subjected general linear model, Pearson correlation and regression analysis. Significant ($P < 0.001$) effects of breed and sex on body weight and body linear measurements were detected. Muscovy duck had higher values in all the parameters compared to khaki Campbell. Males exhibited higher values among the parameters measured than their female counterpart. Most of the correlation coefficients among body weight and linear body measurements were strong positive and significant. The most reliable prediction from regression analysis was between body weight and body length + chest width + chest circumference ($R^2 = 0.919$). The research has shown that body linear measurements could be used to predict body weight in matured ducks. Under field conditions, body length, chest width and chest circumference can be combined in a multiple regression equation to predict the body weight of mature ducks with some level of accuracy.

Keywords: Breed, Ducks, Muscovy, Khaki Cambell Body weight, Body Linear Measurement**INTRODUCTION**

There is continues increased demand for animal protein to meet the nutritional requirements of growing world population of humans. Increased duck rearing would be a great supplement to total poultry production. The relationship existing among body characteristics provide useful information on performance and that these quantitative measures of size and shape are necessary for estimating genetic parameters in animal (Chineke *et al.*, 2002). correlation between weight and body measurement can be used for assessment and selection of body weight (Raji *et al.*, 2009). Therefore, this study is aimed to determine the relationship between body weight and linear body measurements among two breeds of ducks reared under the Nigeria environmental conditions.

Materials and Method**Description of Study Location**

This study was carried out in Bauchi local government area of Bauchi State, Nigeria. Bauchi is located between latitude 10.18° and 57.00° N of the Equator. Longitudinally, longitude 9.50° and 39.00° E of the Greenwich. the climatic condition of the area is characterized by two distinct seasons, dry and wet. It receives rains late, usually around May or June and the average rainfall is about 1091.4mm per annum (BASADP, 2007)

Management of Ducks

The ducks were managed extensively. They are allowed roam freely during the day and scavenge for food and sometime their meals are supplemented with kitchen wastes.



Data Collection

Measurements on body weights (BW) and body linear measurements (BLM) of 150 ducks comprising 78 Muscovy and 72 khaki Campbell were collected within Bauchi local government area. Using a weighing scale, the BW of matured ducks were individually taken in kilograms (kg). The measurements of various body components of matured ducks were taken using a measuring tape calibrated in centimeters (cm). The parameters taken were body length (BL), chest girth (CG), chest width (CW), femur length (FL), femur circumference (FC), metatarsus length (ML), metatarsus circumference (MC), beak length (BKL) and wing length (WL).

Statistical Analysis

Data collected were analyzed using general linear model of SPSS version 25 (2017). Effect of breed and sex on BW and body BLM were determined. Significantly different means were compared using the Duncan multiple range test (DMRT). Pearson correlation analysis was used to determine the relationships among BW and BLM. Separate models (Linear and Multiple) for body measurements singly and combined were enumerated. The regression model adopted was as follows:

$Y = a + b_1 X_1 + b_2 X_2 + b_3 X_3$, Where Y = body weight (kg), X_1 to X_3 = body measurements, a = Intercept $b(1-3)$ = regression coefficients of Y on X ($i = 1, 2, 3$).

Results and Discussion

Effect of Breed on Body Weight and Body Linear Measurement

Effect of breed on BW and BLM are presented in Table 1. Significant ($P < 0.001$) effect of breed on BL, BW, CW, CG, FL, FC, ML, MC, BKL and WL was noticed. Muscovy duck recorded the higher values in all the parameters recorded except BKL in which the higher value was observed among Khaki Campbell. The observed influence of breed in current study is in line with the findings of Purwantini *et al.* (2013) who reported the effect of breed on BW and BLM among ducks. This may be attributed to the genetic differences among the birds. However, the result contradicts the findings of Padhi (2010) who noticed non-significant effect of genotype between pure and crossbreeds.

Effect of sex on body weight and body linear measurement of duck

Means and standard error of the effect of sex on BW and BLM of ducks are presented in Table 2. The duck exhibited a high degree of sexual dimorphism. The drakes were superior ($P < 0.001$) to the ducks for BW and all the body characteristics recorded. The average BW for males and females were 2.60kg and 2.02kg respectively. Higher BW and BLM in drakes than ducks in the present study agreed with the finding of Raji *et al.*, (2009) and Taguia *et al.*, (2007) who reported sexual dimorphism in favour of drakes. This may be attributed to the differential hormonal action between the sexes which invariably leads to differential growth rates (Baeza *et al.*, 2001).

Table 1: Effect of breed on body weight and linear measurement of duck

	BREEDS		LOS
	Muscovy	Khaki Campbell	
BW	2.58±0.07 ^a	1.85±0.30 ^b	***
BL	45.26±0.87 ^a	40.31±0.26 ^b	***
CW	25.35±0.20 ^a	18.52±0.12 ^b	***
CG	39.06±0.54 ^a	28.49±0.19 ^b	***
FL	12.31±0.13 ^a	10.39±0.09 ^b	***



NSAP

47th Annual Conference (JOS 2022)

CONFERENCE PROCEEDINGS

**THEME
SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES**

FC	11.16±0.16 ^a	9.17±0.12 ^b	***
ML	5.50±0.11 ^a	4.54±0.04 ^b	***
MC	4.79±0.07 ^a	3.86±0.02 ^b	***
BKL	5.54±0.07 ^b	6.29±0.06 ^a	***
WL	30.88±0.26 ^a	25.41±0.16 ^b	***

*** = Significant (P < 0.001) CG = Chest Circumference CW = Chest width ML = Metatarsus length BL = Body length MC = Metatarsus circumference BKL = Beak length FL = Femur length BW = Body weight WL = Wing length FC = Femur circumference

Table 2: Effect of sex on body weight and body linear measurement of duck.

	SEX		
	Male	Female	LOS
BW	2.60±0.99 ^a	2.02±0.23 ^b	***
BL	46.95±0.97 ^a	39.97±0.26 ^b	***
CW	23.68±0.49 ^a	21.50±0.38 ^b	***
CG	38.03±0.94 ^a	31.76±0.42 ^b	***
FL	12.42±0.18 ^a	10.72±0.09 ^b	***
FC	11.38±0.20 ^a	9.42±0.10 ^b	***
ML	5.73±0.12 ^a	4.55±0.03 ^b	***
MC	4.80±0.09 ^a	4.10±0.04 ^b	***
BKL	6.19±0.06 ^a	5.60±0.08 ^b	***
WL	30.30±0.42 ^a	27.14±0.30 ^b	***

*** = Significant (P < 0.001) CG = Chest Circumference CW = Chest width ML = Metatarsus length BL = Body length MC = Metatarsus circumference BKL = Beak length FL = Femur length BW = Body weight WL = Wing length FC = Femur circumference

Correlation between Body Weight and Body Measurements

Correlation coefficients of body weight and linear measurements of local ducks are presented in Table 3. Positive and significant (P < 0.01) correlations were observed. The association among the parameters ranged from 0.57 (between CW and WL) to 0.95 (between ML, BW, CC, FC and BKL). The highest correlation value (0.95) was between BL and CG and ML while the least (0.57) was between WL and CW. The result is in agreement with findings of Raji *et al*, (2009) who reported high and positive correlation between linear measurements and body weights among Muscovy ducks and Ogah (2011) who observed strong and positive correlation between body weight and body linear measurement in turkey.

Regression Analysis

Table 4 presented the regression equations and coefficient of determination (R²) of linear measurements of local ducks. The R² values observed in this study ranged from 0.485 to 0.919. The highest value (0.919) was obtained when BL and CW and CG were combined in a multiple regression equation. The least R² (0.485) value was determined between BW and BL. When used singly, CG had the highest R² of



0.899 followed by ML and FL ($R^2 = 0.897$ and 0.886). Raji *et al.* (2009) and Tulki (2017) reported similar

Variable	Regression equation	R^2	LOS	SE of estimate
----------	---------------------	-------	-----	----------------

findings among Muscovy ducks and Turkish geese. The higher association of body weight with chest girth was possibly due to relatively large contribution to body weight by chest girth consisting of bones, muscles and viscera (Raji *et al.*, 2009).

Table-3. Correlation between Body Weight and Body Linear Measurements among Two Breeds of Ducks.

Parameter	BW	BL	CW	CC	FL	FC	ML	MC	BKL	WL
BW										
BL	0.70**									
CW	0.74**	0.58**								
CC	0.95**	0.66**	0.70**							
FL	0.94**	0.62**	0.72**	0.94**						
FC	0.93**	0.63**	0.65**	0.95**	0.92**					
ML	0.95**	0.66**	0.65**	0.95**	0.91**	0.95**				
MC	0.91**	0.64**	0.68**	0.91**	0.86**	0.90**	0.90**			
BKL	0.93**	0.62**	0.67**	0.95**	0.91**	0.94**	0.92**	0.88**		
WL	0.73**	0.60**	0.57**	0.79**	0.69**	0.74**	0.81**	0.68**	0.78**	

** = Significant ($P < 0.01$) CC = Chest Circumference CW = Chest width ML = Metatarsus length BL = Body length MC = Metatarsus circumference BKL = Beak length FL = Femur length BW = Body weight WL = Wing length FC = Femur circumference

Table 4: Regression equations and coefficient of determination (R^2) of body weight and body linear measurements of ducks.



NSAP

47th Annual Conference (JOS 2022)

CONFERENCE PROCEEDINGS

THEME
SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES

BL	$Y = 0.37 + 0.056 \text{ BL}$	0.485	**	0.444
CW	$Y = -4.131 + 0.265 \text{ CW}$	0.550	**	0.415
CC	$Y = -2.210 + 0.123 \text{ CC}$	0.899	**	0.197
FL	$Y = -3.482 + 0.492 \text{ FL}$	0.886	**	0.209
FC	$Y = -1.942 + 0.405 \text{ FC}$	0.859	**	0.233
ML	$Y = -0.858 + 0.624 \text{ ML}$	0.897	**	0.199
MC	$Y = -2.113 + 0.978 \text{ MC}$	0.832	**	0.254
BKL	$Y = -2.752 + 0.961 \text{ BKL}$	0.855	**	0.234
WL	$Y = -3.584 + 0.199 \text{ CW}$	0.538	**	0.420
CC, FL	$Y = -2.925 + 0.071 \text{ CC} + 0.223 \text{ FL}$	0.917	**	0.181
FC, ML	$Y = -1.253 + 0.121 \text{ FC} + 0.451 \text{ ML}$	0.904	**	0.193
MC, BKL	$Y = -2.779 + 0.563 \text{ MC} + 0.566 \text{ BKL}$	0.898	**	0.199
BL, CW	$Y = -3.492 + 0.032 \text{ BL} + 0.182 \text{ CW}$	0.657	**	0.365
BL, CC	$Y = -2.254 + 0.011 \text{ BL} + 0.111 \text{ CC}$	0.909	**	0.188
CC, CW	$Y = -3.065 + 0.108 \text{ CC} + 0.055 \text{ CW}$	0.911	**	0.186
BL, CW, CC	$Y = -2.965 + 0.008 \text{ BL} + 0.047 \text{ CW} + 0.102 \text{ CC}$	0.919	**	0.177

** = Significant ($P < 0.01$) CC = Chest Circumference CW = Chest width ML = Metatarsus length BL = Body length MC = Metatarsus circumference BKL = Beak length FL = Femur length BW = Body weight WL = Wing length FC = Femur circumference.

Conclusion and Recommendation

Muscovy drakes were superior to the ducks for body weight and all the body characteristics studied. Therefore, Prediction of body weight by use of body linear measurements should be by sex within breed rather than by breed only.

REFERENCES

- Baeza, E., Dessay, C., Wacrenier, N., Marche, G., Listrat, A. (2001). Effect of Selection for Improved Body Weight and Composition on Muscle and Meat Characteristics in Muscovy duck. *British Poultry Science*, 43: 560-56
- Chineke, C. A., Agaviezor, B., Ikeobi, C. O. N and Ologun, A. G. (2002). Some Factors Affecting Body Weight and Measurements of Rabbit at Pre and Post Weaning Ages. *Proceedings in the 27th Annual Conference of Nigeria Society for Animal Production*, FUT Akure, Nigeria. 17th-27th march. Pp. 1-3
- National Research Council (NRC), (1994). Nutrient Requirements of Poultry. National Academic Press. Washington, DC.
- Ogah, D. M. (2011). Assessing Size and Conformation of the Body of Nigerian Indigenous Turkey. *Slovak Journal of Animal Science*, 44(1): 21-27.



NSAP

**47th Annual
Conference
(JOS 2022)**

**CONFERENCE
PROCEEDINGS**

THEME
SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES

- Padhi, M. K. (2010). Production Benefits of the Crossbreeding of Indigenous and Non- Indigenous Ducks, Growing and Laying Period Body Weight and Production Performance. *Tropical Animal Health and Production* **42**: 1395-1403.
- Pingel, H. (2009). Waterfowl Production for Food Security. Proceedings of the 4th World Waterfowl Conference 11 – 13 November, 2009, Thrissur, India. Pp. 5-15.
- Purwantini, D., Yuwanta, T., Hartatik, T., Ismoyowati, (2013). Morphology and Genetic Diversity of Mitochondrial DNA D-LOOP Region Using PCR-RFLP Analysis in Magelang Duck and Other Native Duck. *Journal of Indonesian Tropical Animal Agriculture*, **38**: 1-9.
- Raji. A. O., Igwebuike, J. U. And Usman, M. T. (2009). Zoometrical Body Measurements and Their Relation with Live Weight in Matured Local Muscovy Ducks in Borno State, Nigeria. *Agricultural and Biological Science*, **4**(3): 58-62.