
IMPACT OF COMBINATION OF PHYTOGENIC PLANT MATERIALS ON THE GROWTH PERFORMANCE OF STARTER BROILER BIRDS

Uzoma, C., Olabode, A.D., Azodo, L.N. and Adetutu, O.I.

Department of Animal Production Technology,

Federal college of Agriculture, Ishiagu, Ebonyi state.

Corresponding author: uzomachinenye015@gmail.com/Adeoladavid2005@yahoo.com:

ABSTRACT

The research was carried out to determine the impact of combination of different levels of moringa leaf and black plum leaf meal on the growth performance of starter broiler birds. One hundred and twenty (120) day old "Ross 308 strains" broiler chicks were randomly allocated to five dietary treatments, each replicated three times with eight birds per replicate in a completely randomized design (CRD). Five diets were formulated such that diets 2, 3, 4 and 5 contained moringa leaf meal and black plum leaf meal at 5% inclusion level in the ratio 1:4, 2:3, 3:2 and 4:1. Feed and water were given ad-libitum throughout the research period of twenty-one days. Values obtained for final body weight was superior ($P < 0.05$) in treatment 1 with 1321.90g, which did not differ ($P > 0.05$) from 1283.09g in treatment 2, but was different from the values of 1202.86g and 1204.29g obtained in treatment 3 and 4. While, the least value of 1164.10g was observed in treatment 5. Body weight gain was superior ($P < 0.05$) for birds in treatment 1 (1162.73g), while the least value of 1003.68g was obtained in treatment 5. Feed conversion ratio was superior in treatment 1 (1.75) in terms of performance. From the results obtained, it can be concluded that the inclusion of moringa leaf meal and black plum leaf meal at 5% inclusion level and ratios 1:4, 2:3, 3:2 and 4:1 is viable at starter phase without any declining effect on the growth performance of the birds, with the best performance obtained in treatment 1 (control), respectively.

Keywords: Growth performance, impact, starter broiler birds, moringa leaf meal and black plum leaf meal

INTRODUCTION

Natural antioxidants such as vitamin C, tocopherols, flavonoids and other phenolic compounds are known to be present in certain plants. Moringa (*Moringa oleifera*) is one of such plant that has been identified to contain natural antioxidants (Siddhuraju and Becker, 2003). *Moringa oleifera* is rich in bioactive compounds and may be a potential feed source as a phytogenic feed additive. Moringa leaves contain vitamins, flavonoids and carotenoids, which not only serve as essential nutrients, but also enrich poultry meat and eggs, and intensify the pigmentation of the shanks and egg yolk. It is also a good source of dietary fiber, potassium and manganese (Adenike, 2014). The leaves are good sources of digestible protein, β -carotene, vitamin C, calcium, iron and potassium. Moreover, the antioxidant effect of *Moringa oleifera* leaf is due to the presence of polyphenols, tannins, anthocyanin, glycosides and thiocarbamates, which remove free radicals, activate antioxidant enzymes and inhibit oxidases (Luqmans *et al.*, 2012).

Black plum (*Vitex doniana*) is among plant leaves with potential for improving performance in livestock and poultry nutrition. It is an indigenous tropical plant distributed across tropical sub-saharan, Africa's coastal savannas and savanna woodland. The tree is none domesticated, but it is often found at the centre of West African villages. *Vitex doniana* is commonly known as Black plum (English), 'Dinya' (Hausa), 'Oriri' (Yoruba) and 'Uchakoro' (Igbo) where the bark, leaves and roots of the plant are used in ethno-medicine for the management and treatment of numerous disorders such as microbial infection, cancer, rheumatism, hypertension and inflammatory diseases (Atawode, 2005). *Vitex doniana*, as an indigenous leafy vegetable is inexpensive but is a high-quality nutritional source, especially when used in the diets of broiler birds They are found to contain much more potassium and phosphate than calcium, magnesium, zinc and iron (Nnamani, *et al.*, 2007). In view of this, the present study is conducted to determine the growth performance of starter broilers feed replacement level of Black plum and Moringa leaf meal.

MATERIALS AND METHODS

The experiment was carried out at the poultry unit of Animal Production Technology Department, Federal College of Agriculture, Ishiagu, Ebonyi State. The moringa and black plum leaf that was used for the experiment was sourced from Ishiagu town and environment all within Ebonyi state.

Table 1: Experimental diet for starter broilers fed supplemental levels of moringa and black plum leaf meal

Ingredients	Treatments				
	T1	T2	T3	T4	T5
Maize	52.00	50.00	50.00	50.00	50.00
Wheat offal	7.75	4.75	4.75	4.75	4.75
Soyabean meal	8.15	8.15	8.15	8.15	8.15
Groundnut cake	20.00	20.00	20.00	20.00	20.00
Fish meal (72%)	3.50	3.50	3.50	3.50	3.50
Blood meal	3.50	3.50	3.50	3.50	3.50
Black plum leaf meal	0.00	1.00	2.00	3.00	4.00
Moringa leaf meal	0.00	4.00	3.00	2.00	1.00
Limestone	1.50	1.50	1.50	1.50	1.50
Bonemeal	2.50	2.50	2.50	2.50	2.50
Methionine	0.35	0.35	0.35	0.35	0.35
Lysine	0.15	0.15	0.15	0.15	0.15
Starter premix	0.35	0.35	0.35	0.35	0.35
Common salt	0.25	0.25	0.25	0.25	0.25
Total	100	100	100	100	100
Calculated value					
Crude protein (CP)	23.85	23.92	23.90	23.88	23.86
MEnergy (Kcal/kg)	2830.40	2829.70	2829.35	2828.92	2829.70

The leaves were obtained fresh, washed, air-dried and then sun-dried for an hour to get a crisp leafy material. The crispy leaves were then turn to powder by grinding and then incorporated in the diets of the birds.

Table 2: Proximate composition of moringa and black plum leaf meal

Component (%)	Moringa leaf meal	Black plum leaf meal
Dry matter	91.86	89.06
Moisture	8.14	10.94
Crude protein	26.49	11.27
Crude fiber	7.92	7.37
Ether extract	4.31	2.60
Ash	9.62	9.38
Nitrogen free extract	43.52	58.44
Metabolizable energy (Kcal/kg)	2,816.70	2,660.85

One hundred and twenty (120), 1- day old “Ross 308 strain” broiler birds were used for the research work. Each treatment had twenty-four birds with three replicates consisting of eight birds each distributed in a completely randomized design (CRD). The birds were brooded for one week. Feed and water were given *ad-libitum* and vaccinations were given as at when due according to standard practices. The initial weight of the birds was taken at the beginning of the study and then subsequently on a weekly basis. Feed intake was also recorded as the difference between the quantity of feed given the previous day and the quantity that was left the next day. Feed conversion ratio was obtained as the ratio of feed intake divided by the body weight gain. Data collected were subjected to analysis of variance (ANOVA). Significantly different means were separated according to the method of Duncan multiple range test. Proximate analysis of moringa and black plum leaf meal were carried out using the standard procedure of AOAC (2005).

RESULTS AND DISCUSSION

Results of the performance characteristics of broiler birds fed supplemental levels of moringa and black plum leaf meal at starter phase is shown in Table 2. Birds in treatment 1 had a superior ($P<0.05$) final body weight of 1321.90g which was similar ($P>0.05$) to those of 1283.09g observed for birds in treatment 2. While, the least value of 1164.10g was observed in treatment 5. Birds in treatments 3 and 4 had similar ($P>0.05$) values of 1202.86g and 1204.29g. The progressive decrease of final body weight across the group could be due to the inability of the birds to extract necessary nutrients like vitamin and minerals from the test ingredients for their growth. Other contributing factors include the high fibre content of the test ingredients and imbalance of nutrient content and digestibility of the diet. Also, it connotes that the bioactive substance in moringa and black plum could not support increase in body weight to a greater extent when compared to the control. The present study contradicts the findings of Nnamani *et al.* (2007) who observed

Table 3: Growth performance of starter broiler birds fed supplemental levels of Moringa and Black plum leaf meal

Parameters	Treatments					SEM
	T1	T2	T3	T4	T5	
Initial body weight (g)	159.17	160.84	161.25	159.59	160.42	-
Final body weight (g)	1321.90 ^a	1283.09 ^a	1202.86 ^b	1204.29 ^b	1164.10 ^c	18.12
Body weight gain (g)	1162.73 ^a	1122.25 ^b	1041.61 ^c	1044.70 ^c	1003.68 ^d	15.71
Feed intake (g)	2029.38 ^a	1983.76 ^b	1964.69 ^b	1932.51 ^c	1940.94 ^c	9.88
Daily body weight gain (g)	55.37 ^a	53.44 ^a	49.60 ^b	49.75 ^b	47.79 ^c	0.96
Daily feed intake (g)	96.64 ^a	94.47 ^a	93.56 ^a	92.02 ^b	92.43 ^b	0.60
Feed conversion ratio	1.75 ^b	1.77 ^b	1.89 ^a	1.85 ^a	1.93 ^a	0.02

^{abcd} Means on the same row with different superscripts are significantly ($p<0.05$) different.

higher weight in birds fed *Vitex doniana* leaf meal-based diets. Also, Khan *et al.* (2017) and Abdulsalam *et al.* (2015) observed higher body weight when dietary moringa leaf meal was used in broiler birds. But, was similar to the findings of Olugbemi *et al.* (2010), where they observed lower body weight in diets fortified with moringa leaf meal in broiler birds. Treatment effects showed that daily feed intake had superior ($P<0.05$) value of 96.64g in treatment 1, which did not differ ($P>0.05$) from the values of 94.47g and 93.56g obtained for daily feed intake in treatments 2 and 3. The least value of 92.02g was observed in treatment 4, which was similar ($P>0.05$) to that of 92.43g obtained in treatment 5. There was a progressive decrease in the values obtained for daily feed intake across the treatment group. These could be due to the presence of anti-nutritional factors in the moringa and black plum leaves used during the research work. Similar observation was reported by Gakuya *et al.* (2014) who working with *Moringa oleifera* leaves in broiler birds noted a decreased feed consumption. Also, Olabode *et al.* (2022) observed decline in feed consumption across the treatment group when black plum leaf meal was used to fortify diet offered to broiler birds at starter phase. Better turn over of feed to meat was observed in treatment 1 with a value of 1.75 which was followed by birds in treatments 2 with 1.77, while those in treatments 5, 4 and 3 had values which were similar ($P>0.05$) to each other with values of 1.93, 1.85 and 1.89. These results disagree with the report of Adeyina *et al.* (2017) who observed better performance in birds fed *Vitex doniana* based diet. Also, Atuahene *et al.* (2008) reported higher feed conversion efficiency where moringa leaf meal was used to fortify diets given to broiler birds.

CONCLUSION

It can be drawn from the research work that the inclusion of moringa and black plum leaf meal is viable in broiler birds at starter phase at the ratio of 1:4, 2:3, 3:2 and 4:1 respectively as seen from the results obtained in the experiment with the best performance in treatment 1.

REFERENCES

- Abdulsalam, S., Yahaya, M. and Yakasai, M. (2015). Performance of broiler chickens fed on *Moringa oleifera* leaf meal supplemented poultry feed. *Nigeria Agricultural Journal*. 46:139-146.

- Adenike, A.B (2014).** The effects of drying on some nutrients of *Moringa oleifera* leaves. *Food Science and Technology* 15(2): 246-250.
- Adeyina, A.O., Akanbi, A.S., Sanusi, S.B., Olaniyi, B.T., Adegoke, A.G., Hassan, K.T., Olaoye, T.S., Salako, A.O. and Adeyina, O.A. (2014).** Reproductive response to inclusion of graded levels of *Ipomea purpurea* leaf meal (morning glory) in diets of laying chickens. *Journal of Agricultural Science and Environment*. 14, 36-41.
- Agu, C.I., Uzoma, C., Okelola, O.E., Olabode, A.D. and Ebiaku, V. (2021).** Influence of supplemental levels of turmeric meal (*Curcuma longa*) on the growth performance and serum biochemistry indices of finisher broiler birds (A case study in Ishiagu, Ivo, LGA of Ebonyi state, Nigeria). *International Journal of Agriculture and Bioscience*. 10(4): 229-232.
- AOAC (2005).** Official Methods of analysis. 18th edition. Published by the Association of Official Analytical Chemists; Arlington, Virginia.
- Atawodi, S. E (2005).** Comparative *in-vitro* trypanocidal activities of petroleum Ether, chloroform, methanol and aqueous extracts of some Nigerian savannah plants. *African journal of Biotechnol.* 4 177-182.
- Atuahene, C. C., Attoh-Kotoku, V., Fosu, K. D., Amissah, S. E., Sarfo, F. K. and Mensah, J. J. (2008).** Preliminary study of the effect of feeding *Moringa Oleifera* leaf meal as a feed ingredient on the growth performance of broiler chickens. Proceedings of the Ghana Animal Science Association held at the 9 University of Education, Winneba, Mmpong Campus, 28th and 29th August, 2008. pp.72-75.
- Fasuyi, A.O., Fajemibhin, K.S.O. and Omojola, A.B. (2005).** The egg quality characteristics of layers fed varying dietary inclusions of Siam weed (*Chromolaena odorata*) leaf meal. *International Journal of Poultry Science*. 4 (10): 752-757.
- Gakuya, D.W., Mbugua, P.N., Kavoi, B. and Kiama, S.G. (2014).** Effect of supplementation of *Moringa oleifera* leaf meal in broiler chicken feed. *International Journal of Poultry Science*. 13:208-213.
- Khan, I., Zaneb, H., Masood, S., Yousaf, M.S., Rehman, H.F. and Rehman, H. (2017).** Effect of *Moringa oleifera* leaf powder supplementation on growth performance and intestinal morphology in broiler chickens. *Animal physiology and Animal nutrition*. 101:114-121.
- Luqmans, S., S. Srivastava, R. Kumar, A.K. Maurya and D. Chanda, (2012).** Experimental assessment of *Moringa oleifera* leaf and fruit for its antistress, antioxidant, and scavenging potential using *in vitro* and *in vivo* assays. Evidence-Based Complement. *Alternative Medicine*
- Nnamani, C. V., Oscelebe, H. O. and Agbatutu, A. (2007).** Assessment of Nutritonal value of three underutilized Indigenous Leafy Vegetable of Ebonyi State, Nigeria. *African Journal of Biotechnology*, 8 (9): 2321-2324.
- Olabode, A.D., Agu, C.I., Adetutu, O.I., Onyishi, P.I., Olorunfunmilola, N.A and Uzoma, C. (2022).** Growth performance and cost benefit analysis of broiler starter chickens fed diets containing graded levels of Aloe vera (*Aloe barbadensis*) leaf meal. Proceedings of Nigerian Society for Animal Production. 47 (1).
- Olugbemi, T.S., Mutayoba, S.K. and Lekule, F.P. (2010).** Evaluation of *Moringa oleifera* leaf meal inclusion in cassava chip-based diets fed to laying birds. *Livestock Research and Rural Development*. 22: 118-124.
- Siddhuraju, P. and K. Becker, (2003).** Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *Journal of Agriculture and Food Chemistry*, 51: 2144-2155.