

In Vitro gas production assesment of *Panicum maximum* incubated with *Leucaena leucocephala* at varying proportions to predict the nutritional value for ruminants

Falola, O. O. and Olufayo, O. O.



¹Federal College of Animal Health and Production Technology, Ibadan

²Federal Polytechnic Ilaro, Ogun state

Corresponding author: oyenikelubunmi@yahoo.com

Abstract

The inclusion of multipurpose trees with grass such as *Panicum maximum* in the diet of ruminants may enhance productivity. Thus, the study was carried out to determine the proximate composition, in vitro gas production characteristics and parameters of *Leucaena leucocephala* and *Panicum maximum* at varying proportions. Five diets were formulated such that *Panicum maximum* was substituted with *Leucaena leucocephala* at different ratio: T1 (100% *Panicum maximum*), T2 (100% *Leucaena leucocephala*), T3 (50% *P. maximum* + 50% *L. leucocephala*), T4 (75% *P. maximum* + 25% *L. leucocephala*) and T5 (25% *P. maximum* + 75% *L. leucocephala*). Data were subjected to analysis of variance. Values obtained for dry matter (34.43 – 35.95g/100g) decreased with the increased inclusion of *Leucaena leucocephala* in the diets. The crude fiber values (14.33 – 30.75g/100g) also followed the same trend while crude protein (CP) content (10.70 – 26.78g/100g) increased the inclusion of *Leucaena leucocephala* in the diets. There were significant ($P < 0.05$) differences in the treatment means of organic matter digestibility (OMD 31.93 – 37.07%), Metabolisable energy (ME 3.62 – 4.33 MJ/kgDM), short chain fatty acids (SCFA 0.04 – 0.16mL) and methane (ME 1.00 – 2.50 mL). The values (2.00 – 4.67 ml/200mgDM) obtained for immediate soluble 'a' was significantly ($P < 0.05$) different among the treatments. The extent of gas production (a+b) ranged from 2.00 – 4.67 mL/200mgDM, T1 (100% *Panicum maximum*) recorded the lowest while highest was observed in T2 (100% *Leucaena leucocephala*). The insoluble but degradable fraction 'b' ranged from 2.67 – 5.67ml/200mgDM, while the rate of gas production 'c' ranged from 0.04 – 0.14ml/hr. In conclusion, the enhanced values of crude protein, OMD, SCFA, and ME in the *Panicum maximum* and *Leucaena leucocephala* mixture indicate that the diets is able to meet the nutrients requirements of small ruminants in the tropics especially during the dry season.

Keywords: *In vitro*, gas production, nutrients, *Leucaena leucocephala*, *Panicum maximum*

Introduction

Browse species play a major role as feed for ruminants in arid and semi-arid regions, particularly during the dry season when poor quality forage and crop residues are common (Kibon and Orskov, 1993). The inclusion of multipurpose trees like *Leucaena leucocephala*, *Gliricidia sepium* among others is one of the ways of improving the utilization of low quality forage. *Panicum maximum* is one of the common grasses in the derived savannah region of Nigeria. Under good conditions,

its nutritional value is high having 9 – 12.5% crude protein, total digestible nutrients (TDN) of 10.2%, calcium of 0.05% and magnesium of 0.06% (McDonald *et al.*, 1995). However these values can be reduced during the dry season because of poor soil fertility and low rainfall which reduces the growth rate of the grass resulting into high cell wall contents and thereby lowering the quality of the forage. According to Andera and Pablo (1999) a sustainable way of improving feeding value of poor quality pastures is through supplementation with

In Vitro* gas production assesment of *Panicum maximum* incubated with *Leucaena leucocephala

browse plants. Browse plants can be fed to ruminants at varying levels with *Panicum maximum* especially during the dry season as source of supplementary protein. This is because they survive long dry season by the possession of deep rooting system, low transpiration xerophytic foliage and often have good water harvesting architecture. *Leucaena leucocephala* is a promising feedstuff for goats when compared with other legumes such as alfalfa, *Lablab purpureus* and *Gliricidia sepium*. It is rich in nutrients, resulting in better DM intake, weight gain and reproductive performance (Kanani *et al.*, 2006). The in-vitro gas production technique is a laboratory estimation of rate and extent of nutrient disappearance in livestock feed sample for the purpose of assessing the potential nutritive value of the feed. It is also a method that is reproducible and parameters obtained correlate well with in vivo trials (Fajemisin, 2002). In vitro gas production technique has the advantage of not only being less expensive and less time consuming but it allows for more precision in experimental conditions than the in vivo method (Makkar, 2002). It is convenient, fast and allows a large number of samples to be handled at a time. It is based on the quantification of substrate degraded and of gas or short chain fatty acid produced in rumen fermentation system (McDonald, *et al.*, 1995). In vitro gas production is a technique that involves volumetric measurement of gas production in phosphate and bicarbonate buffered in vitro incubation in syringes. Although gases produced during rumen fermentation are waste products and of no nutritive value to the ruminants, gas production tests are used routinely in feed research as gas volumes are related to both the extent and rate of substrate degradation (Blummel *et al.*, 1997a). Thus, the study was designed to

evaluate the nutritional composition of *Panicum maximum* incubated with *Leucaena leucocephala* at different ratio by in vitro gas production technique.

Material and methods

Site of the experiment

The experiment was carried out at Ruminant Laboratory of the Department of Animal Science, University of Ibadan, Nigeria.

Collection of forage samples

The forage samples were harvested from the Federal College of Animal Health and Production Technology, Moor Plantation, Ibadan. The plant samples were air-dried at room temperature for seven days and milled to pass through 1mm mesh and stored in different air-tight labeled container.

Experimental diets preparation

Five experimental diets were formulated such that *Panicum maximum* was substituted with *Leucaena leucocephala* at ratio T1 = 100% *Panicum maximum*, T2 = 100% *Leucaena leucocephala*, T3 = 75% *P. maximum* + 25% *L. leucocephala*, T4 = 50% *P. maximum* + 50% *L. leucocephala*, T5 = 25% *P. maximum* + 75% *L. leucocephala*.

Chemical analysis

Known weight of each air-dried sample was oven dried at 65°C to a constant weight for dry matter determination (DM). Proximate analysis; crude protein, crude fibre, ether extract and ash were analysed according to the standard methods of (AOAC, 1995).

***In vitro* gas production and statistical analysis**

Rumen fluid was obtained from three West African dwarf goats using stomach tube before the morning feed. The animals were fed with concentrate diets (40% corn, 10% wheat offal, 10% palm kernel cake, 20% groundnut cake, 5% soya bean meal, 10% dried brewers grain, 1% common salt, 3.75% oyster shell and 0.25% fish meal). The use of rumen liquor and buffer (9.8g

Falola and Olufayo

NaHCO₃+2.77g Na₂HPO₄+0.57g KCl+0.47g NaCl+0.12g, MgSO₄.7H₂O+0.16g/litre CaCl₂.H₂O) (1:4, v/v) under continuous flushing with CO₂ for incubation was as reported by Babayemi, 2004. The gas production was measured at 3, 6, 9, 12, 15, 18, 21 and 24 hour. After 24 hour post incubation 4ml of 10M NaOH solution was introduced as described (Fievez *et al.*, 2005) to estimate methane. Metabolizable energy (ME, MJ/KgDM) and organic matter digestibility (OMD %) were estimated as established by (Menke and Steingass, 1988) and short chain fatty acids (SCFA) was calculated as reported by Getachew *et al.*, (1998) using 24 h post incubation. ME = 2.20 +0.136?Gv+ 0.057?CP +0.0029?CF, OMD = 14.88+ 0.889Gv +0.45CP + 0.651XA; SCFA = 0.0239Gv – 0.0601, where Gv, CP, CF, and XA are net gas production (ml/200mgDM), crude protein, crude fibre and ash of the incubated samples respectively. Data obtained were subjected to analysis of variance and means were compared where significant using Duncan multiple range F- test (SAS, 2002).

Results and discussion

The result of proximate composition is presented in Table 1. It was observed that

there were significant variations in the values of determined nutrients with the exception of dry matter (DM). The DM of the experimental diets ranged from 34.43 in T2 to 35.95g/100g in T1 while CP of the diets ranged from 10.70 to 26.78g/100g. Result showed that CP increased as the level of *L. leucocephala* increased, implying the diets could be good sources of protein that may meet the requirement of small ruminants for growth and production. The improved CP values observed agreed with the findings of Ezenwa and Aken'ova (1988) that the effect of legume components in a grass-legume mixture would increase the protein quality of the mixture. The crude fibre values observed in this study ranged from 14.33 (T2) – 30.75g/100g (T1) and these values were lower than the values (47.80 – 51.20%) reported for other forage leguminous plants (Ifut, 1987). T1 had the highest ash value (11.05g/100g) which was lower than the value reported by Fajemisin *et al.* (2015). The lowest NFE value (29.69g/100g) was observed in T2 (100% *L. leucocephala*) which was significantly (P < 0.05) different from the values recorded for other *P. maximum L. leucocephala* diets, this implies that the soluble carbohydrates could support the production of volatile fatty acids in the rumen during fermentation (Blummel *et al.*, 1997b).

Table 1: Proximate composition of mixture of *Panicum maximum* and *Leucaena leucocephala*

Treatments	Dry matter	Crude protein	Crude fibre	Ash	Ether extract	NFE
T1	35.95	10.70 ^d	30.75 ^a	11.05 ^a	10.44 ^c	37.06 ^a
T2	34.43	26.78 ^a	14.33 ^d	9.85 ^b	15.35 ^a	29.69 ^c
T3	34.78	15.74 ^c	26.17 ^b	10.45 ^a	15.90 ^a	31.74 ^b
T4	34.95	13.23 ^{cd}	30.08 ^{ab}	10.75 ^a	11.65 ^c	34.29 ^b
T5	34.60	18.26 ^b	20.23 ^c	10.15 ^a	14.12 ^b	37.24 ^a
SEM	0.52	0.08	0.42	0.09	0.46	0.95

T1 = 100% *Panicum maximum*, T2= 100%*L.leucocephala*, T3 = 75% *P.maximum* + 25% *L. leucocephala*,

T4 = 50% *P.maximum* + 50% *L. leucocephala*, T5 = 25% *P.maximum* + 75% *L. leucocephala*.

SEM means standard of means, ^{a, b, c} means with different superscripts on the same row are significantly different p > 0.05

In Vitro gas production assesement of *Panicum maximum* incubated with *Leucaena leucocephala*

The results of the OMD, ME, SCFA and methane are presented in Table 2. Organic matter digestibility (OMD) could be said to be a measure of degradability potentials of the microbes on the substrates especially in the presence of sufficient ammonia nitrogen (NH₃-N) which has influence on bacteria fermentation. The OMD was highest (37.07%) at T1 and lowest (31.93%) at T2. The SCFA or volatile fatty acid (VFA) is a reflection of energy availability in a feed stuff, its level indicates the energy available to the animal; it contributes up to 80% of animal daily energy requirement (Fellner, 2004). This is one of the end products of rumen fermentation. The SCFA is very important for relating feed composition to production parameters and to net energy values of the forages, therefore production of SCFA from in vitro gas measurement will be increasingly important in a developing country. The metabolizable energy (ME) obtained from this study were significant ($P < 0.05$) the values ranged from 3.62-4.33MJkgDM. The in vitro gas production method has been widely used to evaluate the energy value of several classes of feed (Getachew *et al.*, 1998). The result obtained in this study were within the range (2.99 – 4.75) reported by Babayemi and Bamikole, (2006). Methane production observed in this study ranged from 1.50 – 2.00ml 200mg/DM T1 and T4 recorded the highest while the lowest was observed in

T5. Methane production is said to be energy loss to ruminants and also contribute to global warming (Babayemi and Bamikole, 2006). In most cases, feedstuffs that show high capacity for gas production are also observed to be synonymous for high methane production; also most tropical feedstuffs have been implicated to increase methanogenesis (Babayemi *et al.*, 2004; Babayemi and Bamikole, 2006) as an integrated part of carbohydrate metabolism. Although gas production is a nutritionally wasteful product (Mauricio *et al.*, 1999) but provides a useful basis from which ME, SCFA and OMD may be predicted. The SCFA or VFA such as butyrate and acetate suggest a potential to make energy available to ruminants. Table 3 shows the in vitro gas production characteristics of *Panicum maximum* incubated with *L. leucocephala* at different proportions.

The intercept value 'a' at 24 hours ranged from 2.00 – 4.67ml200mg/DM. The value 'a' used ideally reflect the fermentation of soluble fraction. The values observed for 'a' were significant ($P < 0.05$) among the treatments. The soluble fraction makes the attachment by rumen micro-organism to be done easily and lead to much gas production. There were significant differences in the extent of gas production 'a+b' which ranged from 4.67 to 9.34ml200mg/DM. The potential degradability a+b of a diet depicts the level at which the diets could be degraded if it were in actual rumen of the

Table2: in vitro gas production characteristics (200mg/DM) of the forage mixtures

Treatments	OMD%	SCFA(μmol)	ME(Kg/DM)	Methane(ml)
Treatment1	31.93	0.07	3.62	2.00
Treatment 2	37.07	0.16	4.33	1.50
Treatment 3	32.57	0.07	3.68	1.50
Treatment 4	35.93	0.08	3.74	1.50
Treatment5	33.56	0.08	4.01	1.50
SEM	1.28	0.03	0.20	1.00

T1 = 100% *Panicum maximum*, T2= 100% *Leucaena leucocephala*, T3 = 75% *P.maximum* + 25% *L. leucocephala*

T4 = 50% *P.maximum* + 50% *L. leucocephala*, T5 = 25% *P.maximum* + 75% *L. leucocephala*

SEM means standard of means, ^{a, bc} means with different superscripts on the same row are significantly different $p > 0.05$

Falola and Olufayo

Table3: in vitro parameters (200mg/DM) of *P.maximum* and *L. leucocephala* at different ratio

Treatments	A	A+b	b	C
T1	2.00 ^c	4.67 ^c	2.67 ^c	0.04 ^c
T2	4.67 ^a	9.34 ^a	5.67 ^a	0.14 ^a
T3	4.33 ^b	7.66 ^b	3.33 ^b	0.07 ^b
T4	2.67 ^c	5.19 ^c	3.52 ^a	0.08 ^b
T5	4.66 ^a	9.18 ^{ab}	4.52 ^b	0.12 ^a
SEM	0.87	0.03	0.79	1.44

T1 = 100% *Panicum maximum*, T2= 100% *Leucaena leucocephala*, T3 = 75% *P.maximum* + 25% *L. leucocephala*

T4 = 50% *P.maximum* + 50% *L. leucocephala*, T5 = 25% *P.maximum* + 75% *L. leucocephala*

SEM means standard of means,^{a, b, c} means with different superscripts on the same row are significantly different $p > 0.05$

a = gas produced from immediately soluble fraction, b= extent of gas production from insoluble but degradable fraction

a + b = potential extent of gas production, c = rate of gas production at time (t).

animal (*in vivo*). This largely depends on how much of the fiber fraction (NDF and ADF) have been broken down for easy access of the microbes to the nutrients available in the diets. The extent of gas production 'b' described the fermentation of insoluble but degradable fraction of the samples, result obtained in this study ranged from 2.67 – 5.67 ml/200mg/DM. Blummel and Orskov (1993) found that 'b' value could account for 88% for voluntary feed intake. The rate of gas production 'c' ranged from 0.04 – 0.14ml/hr the values were significantly ($P < 0.05$) different among the treatments, T2 recorded the highest while T1 had the lowest. Babayemi *et al.*, (2004) reported that factors that determine the rate of gas production during fermentation depend on the nature and level of the fibre, the presence of secondary metabolite and the potency of rumen liquor for incubation. Orskov and Ryle, (1990) also reported that the rate 'c' determines digestion time and consequently how long a potentially digestible material would occupy space in the rumen.

Conclusion

The enhanced values obtained in terms of in vitro gas production parameters and gas characteristic indicated that *Panicum maximum*- *Leucaena leucocephala* diets can meet the nutrients requirement of small ruminants particularly during the dry

season.

References

- Andera, M. M. and Pablo, E. C. 1999. A tropical forage solution to poor quality ruminant diet. A review of *Lablab purpureus*, *Livestock Research for Rural Development*, 11(2): 34–56.
- AOAC, 1995. Official Methods of Analysis of the Association of Official Analytical Chemistry. 16th Edition., AOAC International, Washington DC, USA., Page:1141.
- Babayemi, O. J. and Demeyer, D. and Fievez, V. 2004. In vitro fermentation of tropical browse seeds in relation to their content of secondary metabolites. *J. Anim. Feedsci* 13(1): 13-34.
- Babayemi, O. J. and Bamikole, M. A. 2006. Effects of *Tephrosia candida* DC leaf and its mixture with Guinea grass on in vitro fermentation changes as feed for ruminants in Nigeria. *Pak. J. Nutr.* 5(1): 14–18.
- Blummel, M., Makkar, H. P. S., Changa, G., Mtimuni, J. and Becker, K. 1997b. The prediction of dry matter intake of temperate and tropical forages from in vitro digestibility/gas production data and the sry matter intake and in vitro digestibility of African roughages in

In Vitro gas production assesment of *Panicum maximum* incubated with *Leucaena leucocephala*

- relation to ruminants live weight gain. *Animal feed science and technology* 69;131–141
- Blummel, M., Makkar, H. P. S. and Becker, K. 1997a.** In vitro gas production: a technique revisited. *Journal of animal physiology and animal nutrition*, 77: 24-34.
- Blummel, M. and Orskov, E. R. 1993.** Comparison of gas production and Nylon bag degradability of roughages in predicting feed intake in cattle. *Animal Feed Science and Technology*, 40:109–119.
- Ezenwa, L. and Aken'ova, M. E. 1988.** Performance of mixtures of selected grasses and herbaceous legumes in southwest Nigeria. *Tropical grasslands*, 32; 131–138.
- Fajemisin, A. N. 2015.** Evaluation of nutritional composition of *Panicum maximum* *Moringa oleifera* diets by in vitro gas production techniques. *Nigeria Journal of Animal Science* 17(2):214–224.
- Fajemisin, A. N. 2002.** Mineral Utilization by West African Dwarf Goats Fed Rumen Epithelium Based Diets. Ph.D. thesis submitted at Department of Animal Science, University of Ibadan, Ibadan, Nigeria.
- Fellner, V. 2004.** In vitro versus in vivo estimates of ruminal fermentation. Department of animal science, North Carolina state university.
- Fievez, V., Babayemi, O. J. and Demeyer, D. 2005.** Estimation of direct and indirect gas production in syringes: A tool to estimate short chain fatty acid production requiring minimal laboratory facilities, *Animal Feed Science Technology* (123-124): 197-210.
- Getachew, G., Makkar, H. P. S. and Becker, K. 1998.** Stoichiometric relationship between short chain fatty acid and in vitro production in presence and absence of polyethylene glycol for tannin containing browses, EAAP Satellite Symposium, Gas production; fermentation kinetics for feed evaluation and to assess microbial activity 18-19 August, Wageningen, The Netherlands.
- Ifut, O. J. 1987.** The nutritional value of *Gliricidia sepium*, *Panicum maximum* and peels of *Manihot* spp. Fed west African dwarf goats. Ph.D thesis, University of Ibadan, Ibadan, 181Pp.
- Kanani, K., Lukefahr, S. D. and Stanko, R. L. 2006.** Evaluation of tropical forage legumes (*Medicago sativa*, *Dolichos lablab*, *Leucaena leucocephala* and *Desmanthus biconotus*) for growing goats. *Small ruminant research*. 65(12): 1–7.
- Kibon, A. and Orskov, E. R. 1993.** The use of degradation characteristics of browse plants to predict intake and digestibility by goats. *Anim.prod.* 57: 247–251.
- Makkar, H. P. S. 2002.** Application of the in vitro gas method in the evaluation of feed resources and enhancement of nutritional value of tannin-rich tree/browse leaves and agro-industrial by-products. Proceedings of the final review meeting of an IAEA Technical.
- Mauricio, R. M., Mould, F. L., Abdalla, A. I. and Owen, E. 1999.** The potential nutritive value for ruminants of some tropical feed stuffs as indicated by in vitro gas production and chemical analysis. *Animal Feed*

Falola and Olufayo

- Science and Technology* 79: 321 – 330.
- McDonald, P., Edward, R. A., Grenhalgh, J. F. D. and Morgan, C. A. 1995.** Animal Nutrition. 5th edition. Pearson Education Limited, Edinburg gate, Harlow Essex CM202JE, United Kingdom.
- Menke, K. H. and Steingass, H. 1988.** Estimation of the energetic feed value from chemical and in vitro gas production using rumen fluid. *Animal Research Development*, 28:7-55.
- Orskov, E. R. and Ryle, M. 1990.** Energy nutrition in ruminants. Elsevier applied science. London.
- SAS, 2002.** Statistical analytical system institute, SAS/STAT User's guide statistical, SAS Institute, Cary. North Carolina, USA.

Received: 11th February, 2017

Accepted: 27th July, 2017