



## **Age Based Classification of Morphometric Traits of Different Nigeria Goat Breeds**

**Shoyombo<sup>1\*</sup>, A.J., Alabi<sup>1</sup>, O.O., Falana<sup>1</sup>, B.M., Ajala<sup>1b</sup>, A.O., Musa<sup>2</sup>, A.A., Popoola<sup>3</sup>,  
M.A., Ogunniyi<sup>4</sup>, E., and Jubril, A.E**

**<sup>1</sup>Department of Agriculture, Animal Science Programme, College of Agricultural Sciences, Landmark University Omu-Aran, Kwara State, Nigeria**

**<sup>1b</sup>Department of Agriculture, Agricultural Economics and Extension Programme, College of Agricultural Sciences, Landmark University Omu-Aran, Kwara State, Nigeria**

**<sup>2</sup>Department of Animal Science, Faculty of Agricultural Sciences, Kogi State University Anyigba, Kogi State, Nigeria.**

**<sup>3</sup>National Biotechnology Development Agency, Abuja**

**<sup>4</sup>Department of Animal Production, school of Agriculture and Agricultural Technology, Minna, Niger State, Nigeria**

**<sup>5</sup>Department of Animal Science, Faculty of Agriculture, University of Agriculture, Abuja, Nigeria**

**Corresponding Author: [shoyombo.ayoola@lmu.edu.ng](mailto:shoyombo.ayoola@lmu.edu.ng) +23408036926455**

### **Abstract**

The objective of this study was to determine the effect of age on the biometric traits of different Nigeria goat breeds. The effects of age, breed, and sex on linear measurement were estimated using the GLM procedure of the statistics analysis software package. Means were separated within age groups using Tukey test. Sahelian breed was superior to other breeds across all age groups in body weight, while the red Sokoto was intermediate followed by the West African Dwarf. However, variations in morphometric measurements did not follow this trend and indicates that body parts are influenced differently by age, breed and sex. The variability of body weight suggests possibility of improving this trait through careful selection. Present results showed that age has strong influence with a linear response as the animals aged. It is concluded, that age had significant influence on the improvement of size and shape of goat breeds in Nigeria.

**Key words:** Age, Morphometric, Goats, Breed

### **Introduction**

Goat population in Nigeria are estimated as the largest among the ruminant animals totaling about 58.3 million (FAOSTAT, 2008). Goats, predominantly are kept by the rural household and ownership which spread across human diversity (Akpa, 1999). The importance of goat stem from its ability to convert forages and household by-product into high quality human food and their role as locally available source of food, fibre, power and other products to suit local community needs. Several scholars (Adu and Ngere, 1979; Dettmers, 1983; Bunge *et al.*, 1990 and Hall, 1991) has done a lot of work on the genetic improvement of goats. But majority of these were carried out on herds from research institute and universities while there exist little information on morphological and morphometric characteristics and their differentiation in the Nigerian breeds of goat based on on-farm (small holder herds) since these farm herds make up the bulk of the goat population of Nigeria, they serve as a true reflection of the genetic potential of this specie, therefore this study is designed to supplement existing literatures on Nigerian



goats using on-farm herds from smallholder stocks. The influence of age on production characteristics cannot be overemphasized, reports on the impact of age on puberty, kidding, growth and growth rate, birth weight etc. has been documented (Banerjee *et al.*, 2000; Mamabolo and Webb, 2005; Akpa *et al.*, 2013 ). Ojedapo *et al.* (2007) reported that age had strong, positive and significant correlations with morphometric traits in goats. It is on this basis that the study was designed to characterize age based variations of morphological traits within the different breeds of goats.

### **Materials and Methods**

The animals used for this study were sampled in the abattoir, of Borno, Sokoto and Ogun states. The sampling was purposive and consisted of nine hundred (900) goats of different breeds which consisted of fifty males and fifty females distributed in the following age groups <1, 1-2 and 2-3 years. Age was classified based on dentition methods. The following metric characters used in this study were described by Salako and Ngere (2002).

### **Data analysis**

Fixed factor effect of age, breed and sex were estimated using the GLM procedure of the statistics analysis software (SAS, 1990) package. These were computed on the basis of interaction with age groups. Means were separated within age groups using Duncan multiple range test.

### **Results and Discussions**

Mean body weight of goat breeds less than one year are presented in Table 1. The Sahel breed is superior to the other breeds with a body weight of  $13.56 \pm 0.56\text{Kg}$  and  $14.94 \pm 0.52\text{Kg}$  in does and bucks respectively. Within the Red Sokoto population, body weight in doe and buck was  $13.73 \pm 0.51\text{Kg}$  and  $12.86 \pm 0.42\text{Kg}$  respectively while in West African Dwarf,  $10.44\text{Kg} \pm 0.22$  and  $10.62 \pm 0.21\text{Kg}$  for does and bucks. Coefficient of variation ranges between 13.98% and 29.40% suggesting the influence of major genes on this trait. The standard deviation estimates associated with the mean liveweight were generally small, (1.49-3.90). In Horn length, the Sahel had the highest ( $7.21 \pm 0.33\text{cm}$  and  $7.42 \pm 0.31\text{cm}$ ), while the least values ( $5.24 \pm 0.30\text{cm}$  and  $4.99 \pm 0.28\text{cm}$ ) for does and bucks respectively. Averagely, Red Sokoto showed an intermediate distribution ( $6.20 \pm 0.32\text{cm}$  and  $5.22 \pm 0.23\text{cm}$ ) respectively. Associated coefficient of variation and standard deviation ranges between 29.10% – 40.93% and 1.60-2.27 respectively, suggesting that the measurement is almost independent of the environment. Ear length in the Sahel had the highest values ( $11.52 \pm 0.25\text{cm}$  and  $11.06 \pm 0.27\text{cm}$ ), while the least values ( $9.56 \pm 0.15\text{cm}$  and  $9.43 \pm 0.17\text{cm}$ ) for does and bucks respectively. Averagely, Red Sokoto showed an intermediate distribution ( $10.22 \pm 0.17\text{cm}$  and  $9.86 \pm 0.15\text{cm}$ ) respectively. Associated coefficient of variation and standard deviation ranged between 10.81% – 16.96% and 1.07-1.88 respectively. Red Sokoto goats had the highest values for height at withers ( $52.06 \pm 0.87\text{cm}$  and  $51.63 \pm 0.81\text{cm}$ ), while the least values ( $42.47 \pm 1.42\text{cm}$  and  $45.80 \pm 1.34\text{cm}$ ) for does and bucks respectively was observed in Sahel. West African Dwarf buck had withers height of  $47.93 \pm 1.13\text{cm}$  and does have  $48.07 \pm 1.07\text{cm}$ . Associated coefficient of variation and standard deviation ranges between 11.03% – 23.64% and 5.69-10.04 respectively. For mean pouch girths, Red Sokoto had the highest values ( $55.37 \pm 1.32\text{cm}$  and  $59.61 \pm 1.11\text{cm}$ ), while the lowest values of  $51.42 \pm 0.77\text{cm}$  and  $50.99 \pm 0.71\text{cm}$  were observed for does and bucks respectively in WAD. In Sahel goats, pouch girth measured  $52.20 \pm 0.49\text{cm}$  in bucks and  $50.63 \pm 0.66\text{cm}$  in does. The coefficient of variation for this trait ranged from 6.70% to 16.83% while standard deviation was from 3.50 to 9.32. Red Sokoto goats had the highest HG values



(53.33±1.31cm and 57.53±1.17cm), while Sahel had 49.90±0.49cm and 48.77±0.66cm for does and bucks respectively. WAD bucks had a mean heart girth of 50.15±0.78cm and 49.71±0.73cm for does. Coefficient of variation and standard deviation for heart girth were between 6.74%–17.33% and 3.46 – 9.24 respectively.

WAD had the highest BL values (42.04±0.83cm and 41.88±0.73cm), while the least values (35.36±1.01cm and 39.56±1.31cm) were for Sahel does and bucks respectively. Red Sokoto bucks and does had body lengths of 41.92±0.67cm and 41.97±0.56cm respectively. Coefficient of variation between 9.50%–23.39% and standard deviation from 3.99 to 9.25 were obtained for this trait. WAD goats had the highest NC values (23.47±0.59cm and 24.39±0.59cm), while the least values (21.02±0.52cm and 23.31±0.64cm) for does and bucks respectively were observed in Sahel goats. Neck circumference was 22.69±0.72cm in does and 23.43±0.61cm in bucks of Red Sokoto. Associated coefficient of variation and standard deviation ranges were between 17.04%–22.48% and 3.67-5.10 respectively. Red Sokoto goats had the highest tail length 11.71±0.30cm (bucks) and 12.08±0.26cm (does). The lowest length of 10.29±0.19cm and 10.27±0.21cm for doe and buck respectively was observed in WAD goat, while Sahel does and bucks had 11.62±0.32cm and 11.97±0.32cm respectively. Associated coefficient of variation and standard deviation were between 12.90%–19.52% and 1.33 – 2.28 respectively. Shoulder width in Sahel does and bucks were 18.18±0.43cm and 18.59±0.36cm, 13.34±0.56cm and 13.83±0.48cm in WAD and 12.75±0.43 and 12.99±0.48 in Red Sokoto does and bucks respectively. Coefficient of variation for this trait was between 13.80% – 29.57% and standard deviation from 2.57-3.94.

Summarized in Table 2 are Sahel goats had a mean weight of 17.13±0.43Kg and 16.86±0.49Kg followed by West African Dwarf goats with 15.45Kg±0.54 and 17.65±0.44Kg for does and bucks respectively. Red Sokoto goats weighed 13.60±0.46Kg (does) and 13.51±0.43Kg (bucks). Coefficient of variation from 17.56% – 23.85% suggest the influence of major genes on body weight. The standard deviation estimates associated with the mean liveweight were generally small, (3.01-3.79). Results for horn length in goats showed that Sahel had 9.45±0.31cm and 8.64±0.40cm, Red Sokoto (6.91±0.31cm and 6.70±0.24cm) and WAD (6.23±0.33cm and 5.64±0.23cm) for does and bucks respectively. Coefficient of variation (23.27% – 37.42%) was moderate and standard deviation ranged from 1.63 to 2.82. Mean ear lengths indicated the Sahel goats had ear lengths of 13.82±0.40cm and 12.81±0.38cm, 12.24±0.30cm and 11.88±0.32cm for Red Sokoto goats and 11.25±0.22cm and 10.52±0.23cm for WAD does and bucks respectively. Associated coefficient of variation for ear lengths was 13.70% – 20.80%. Sahel recorded highest values (60.58±0.87cm and 60.12±0.88cm), while the least values (52.20±1.04cm and 49.37±1.20cm) for doe and buck respectively was observed in WAD. Red Sokoto showed an intermediate distribution of 55.47±0.77cm for does and 53.60±0.91cm in bucks. Associated coefficient of variation and standard deviation ranges between 9.78–17.24 and 5.42%-8.51% respectively. The mean pouch girth in Red Sokoto had the highest values (55.37±1.32cm and 59.61±1.11cm), followed by Sahel with 52.20±0.49cm and 50.63±0.66cm while the least values of 51.42±0.77cm and 50.99±0.71cm for doe and buck respectively were observed in WAD. Coefficient of variation for pouch girth was between 6.70%–16.83% with standard deviation ranging from 3.50 to 9.32. Mean heart girth in Red Sokoto goats had 63.92±0.60cm and 62.14±0.72cm, Sahel goats had 60.58±0.87cm and 59.87±0.79cm while West African Dwarf goat had 60.80±0.90cm and 54.73±0.67cm for does and bucks respectively. Associated coefficient of variation and standard deviation were between 6.61%–10.47% and 4.22-6.37 respectively. Body length in Sahel does was 61.14±0.97cm and 53.76±0.75cm for bucks, WAD does have a mean body length of



51.84±0.97cm and bucks measured 49.23±0.87cm while Red Sokoto does measured 50.93±1.21cm and 46.83±1.99cm for bucks. Associated coefficient of variation and standard deviation ranges between 9.81%–18.10% and 5.28-8.71 respectively. Mean neck circumference showed WAD goats had the highest values (25.87±0.53cm and 26.16±0.65cm), followed by Sahel goats (25.33±0.44cm and 25.97±0.35cm), then Red Sokoto goats (24.73±0.35cm and 24.69±0.47cm) for does and bucks respectively. Coefficient of variation was from 9.43%–17.63% and standard deviation lies between 2.44 - 4.61. Tail length in Sahel does was 13.48±0.26cm and 13.43±0.31cm in bucks, 12.27±0.27cm in does and 12.79±0.25cm for Red Sokoto bucks, 10.95±0.29cm and 11.56±0.25cm in West African Dwarf does and bucks respectively. Coefficient of variation for tail length in goats studied ranged between 13.49%–18.78%. The shoulder width measurement showed the Sahel had the highest values (23.24±0.44cm and 22.58±0.42cm), while the least values (13.31±0.30 and 13.54±0.40) for doe and buck respectively were observed in Red Sokoto. Averagely, WAD showed an intermediate distribution (14.12±0.48cm and 16.23±0.49cm) respectively. Associated coefficient of variation and standard deviation ranges between 13.26–24.21 and 2.12-3.57 respectively.

Mean liveweight in goat breeds above 2 years are presented in Table 3. Sahel goats had a mean weight of 23.60±0.69Kg and 25.12±0.73Kg in doe and buck respectively is superior to Red Sokoto and WAD goats. Within the Red Sokoto population, mean weight was 18.95±0.47Kg and 19.21±0.47Kg; and 19.13Kg±0.29 and 19.35±0.24Kg in West African Dwarf does and bucks respectively. Coefficient of variation ranged from 8.80 – 20.74 suggesting the influence of major genes on live weight of goat breeds studied and standard deviation of 1.70 to 5.05. Sahel goats had the highest horn lengths of 10.53±0.38cm and 10.76±0.36cm, followed by Red Sokoto goats (7.47±0.29cm and 7.29±0.25cm) and least in WAD goats (7.25±0.30cm and 6.81±0.28cm) for doe and buck respectively. Coefficient of variation and standard deviation were between 23.50 – 29.36% and 1.77 - 2.72 respectively. Mean ear length in Sahel had the highest values (13.71±0.33cm and 13.70±0.29cm), while the least values (11.22±0.32cm and 10.64±0.24cm) for doe and buck respectively in WAD breed. Red Sokoto does have a mean ear length 12.33±0.33cm and 11.88±0.21cm in bucks. Coefficient of variation ranged from 12.43%–20.06% and standard deviation was from 1.48 to 2.35. The mean wither height in Sahel goats had 70.51±0.67cm and 72.02±0.58cm, followed by 58.17±0.85cm and 58.56±0.81cm in Red Sokoto goats and 46.62±0.87cm and 45.42±0.78cm in WAD goats for doe and buck respectively. Coefficient of variation and standard deviation were 5.74%–13.19% and 4.14-6.15 respectively. The mean pouch girth in WAD was the highest (69.94±0.92cm and 69.47±0.95cm), Sahel had 66.96±0.48cm and 67.27±0.77cm and Red Sokoto had the lowest (63.46±0.65cm and 63.23±0.57cm) for doe and buck respectively. Coefficient of variation was 5.11%–9.46% and standard deviation ranged from 3.43 to 6.75. The mean heart girth in Red Sokoto goats had the highest (68.93±0.64cm and 68.57±0.54cm), followed by Sahel goats (64.61±0.48cm and 65.69±0.33cm) and WAD goats with 61.15±0.65cm and 60.64±0.58cm for doe and buck respectively. Associated coefficient of variation and standard deviation ranges between 3.58%–7.57% and 2.35-4.53 respectively. Body length for Sahel goats was 58.53±0.82cm and 56.28±0.99cm, 43.76±0.86cm and 43.69±0.66cm in West African Dwarf goats and 43.76±0.86cm and 43.69±0.66cm in Red Sokoto goats for doe and buck respectively. Associated coefficient of variation and standard deviation ranges between 9.86%–23.35% and 4.70-10.51. Mean neck circumference was 27.70±0.37cm and 27.15±0.35cm in Red Sokoto goats, 27.16±0.61cm and 26.52±0.55cm in Sahel goats and WAD goats had 24.91±0.51cm and 24.66±0.38cm respectively in does and bucks. Coefficient of variation was from 9.06% to 15.70% and standard deviation ranged from 2.44 - 4.61 for this trait. Similar and high tail



lengths for Sahel ( $12.69 \pm 0.27$  cm and  $12.90 \pm 0.31$  cm) and Red Sokoto goats ( $13.15 \pm 0.30$  cm and  $12.45 \pm 0.24$  cm) were observed while WAD goats had  $11.81 \pm 0.22$  cm and  $11.93 \pm 0.20$  cm for does and bucks respectively. Coefficient of variation was between 11.97% and 16.84%; standard deviation ranged from 1.71 to 2.13. The shoulder width measurement in Red Sokoto, Sahel and West African Dwarf breed in this study. Sahel goats had  $24.97 \pm 0.37$  cm and  $24.71 \pm 0.44$  cm, followed by WAD ( $19.65 \pm 0.58$  and  $19.93 \pm 0.70$ ) and Red Sokoto goats with  $14.98 \pm 0.38$  cm and  $15.40 \pm 0.41$  cm for doe and buck respectively. Associated coefficient of variation was 10.61%–24.96% and standard deviation ranged from 2.65 to 4.97.

Mean live weight of 13.73 Kg, 13.56 Kg, 10.44 Kg and 12.86 Kg, 14.94 Kg and 10.44 Kg for does and bucks for the three breeds respectively for age less than one year are within the range of 12-18 Kg reported by Katongole *et al.* (1996) for indigenous goats in Kenya. Mean live weight of 13.60 Kg, 17.13 Kg, 15.45 Kg and 13.51 Kg, 16.84 Kg and 17.65 Kg for does and bucks between the age one through two years are lower than values of 20-40 kg reported by several authors (Akpa *et al.*, 1998, Otoikhan *et al.*, 2008 and Salako *et al.*, 2006) in different breeds of goat. Mean live weight of 18.95 Kg, 23.60 Kg, 19.13 Kg and 19.21, 25.12 Kg and 19.35 Kg for does and bucks above two years are closer to the range reported by several authors (Akpa *et al.*, 1998, Osuhor *et al.*, 2002 and Otoikhan *et al.*, 2008), with Sahel breed fell within the specified range of 20-40 Kg. The average body weight at different ages fell within the range of 16-30 Kg specified by Hassan and Ciroma (1990) for both does and bucks from age 1-5 years in different breeds of goats. The associated coefficient of variation which range between 13.98 and 29.40 across the breeds suggests absence of selection criterion and is also a pointer to the presence of major genes since live weight is a quantitative trait and under the influence of many genes of small effect.

The body confirmation traits were comparable to those reported by Ibrahim (2010) for Red Sokoto goats. These measurements were less variable than previous measurements on Red Sokoto goats (Akpa *et al.* 1998) including for both lactating and dry does. The variability of body weight suggests possibility of improving this trait through careful selection. The sex dependent variation recorded among the breeds was also observed by Salako *et al.*, (2006) who commented that the does differ from the bucks in that they have slower growth rate with earlier maturing carcass. Hassan and Ciroma (1990) also reported that bucks were heavier than does which lend credence to the findings of this study on the average.

Sex had a robust variation for liveweight and body measurements. The sexual dimorphism might be controlled hormonally in male animals until testes are well developed (Akpa *et al.*, 1998). This report is in consonance with the report of Devendra and Burns, (1983) who reported that interactions between sex and strains significantly affect bodyweights and linear traits. This however contrast the report by Otoikhan *et al.* (2008) that sex has no influence on weight gain of Red sokoto and West African Dwarf goat since non-significant difference was found on the overall body weight of the animals. The discrepancy in the reports could be ascribed to breed, age and environmental differences. Umoren and Orok (1994) observed that male animals generally have higher linear body measurement than females. Adu *et al.* (1978) made similar observations in the four indigenous sheep breeds of Nigeria, Salako and Ngere (2002) in WAD. This agrees with patterns observed in this study where inspection of ordinary mean figures showed that males appear superior to females. The relationships between BW and morphological traits appeared to be higher in bucks than in does.

The average height at wither for bucks and does in this study (58.91 and 57.23 cm) is within the range reported by Hassan and Ciroma (1990) whose observations ranged between 56-67



cm. The average body length (48.25 and 47.92 cm), horn length (7.88 and 7.89cm), heart girth (59.42 and 58.67 cm), pauch girth (60.05 and 60.68 cm), neck circumference (24.36 and 25.23 cm), ear length (11.64 and 11.57 cm), tail length (11.89 and 12.07 cm) and scrotal circumference (0.00 and 17.04 cm) for does and bucks are within the range reported by Salako (2002) in different goat breeds. Katongola *et al.*(1996) for indigenous goats in Kenya, Samakula *et al.* (2010) for Mubemde and South East African goat breed, and Abubakar (2012) in Red Sokoto goats. Also, the report of this finding is in agreement with the observation of Salako and Ngere (2002) on body parameters in WAD. However, Ibrahim (2010) reported a shorter body length in Red Sokoto goats which was contrary to the observed longer body length recorded in this study. This could probably be due to the different age classification methods used in this study. In normal animals, increase in age leads to increase in size; this may be the cause of variation in body length. The observed body measurements are within the range reported by Hassan and Ciroma (1990). The low coefficient of variation among the morphometric traits reflects poor environmental sensitivity in all the studied breeds.

#### Conclusion

Present results showed that age has strong influence with huge variability in bodyweight. This implies that improvement of bodyweight among different breeds should take cognizance of age as a covariate for fast response.

#### References

- Adu, I.F., and Ngere, L.O. (1979). The indigenous Sheep of Nigeria. *World Review of Animal Production*. 15(3); 52 – 62.
- Akpa, G.N. (1999). Evaluation of Dairy Characteristics and lactation curves of small holder Red Sokoto (Maradi) Goats. Ph.D Thesis.Ahmadu Bello University, Zaria pp 64-68.
- Akpa, G.N., Ambali, A.L. and Suleiman, I.O. (2013).Body Conformation, Testicular and Semen Characteristics as Influenced by Age, Hair Type and Body Condition of Red Sokoto Goat. *N Y Sci J* 6(7):44-58]. (ISSN:1554-0200).
- Akpa, G.N., Duru, S. and Amos, I.T. (1998). Influence of strain and Sex on estimation of within age-group body weight of Nigerian Maradi goats from hair linear body measurements. *Tropical Agriculture (Trinidad)* 74:462-467
- Banerjee, A.K., Animat, G., and Ermias, E. (2000).Selection and Breeding Strategies for increased production of goats in Ethiopia. In: Merkel, R.G., and Goetsch, A.L, (ed). The opportunities and challenges of enhancing goat production in east Africa, PP. 70 – 79. Dehub University, Awassa, Ethiopia 10 – 12 November 2000.
- Bunge, R., Thomas, D.L., and Stookey, J.M. (1990) Factors Affecting the Productivity of Ram bottle ewes mated to ram Lambs. *J. Anim. Sci.* (68). 2253 – 2262.
- Dettmers, A. (1983). Performance of Hair Sheep in Nigerian. In Fitzhugh, H. E. and Bradford, G.E. (eds). *Hair Sheep of West Africa and the Americas*. Boulder, Col., West View Press pp 201 – 218.
- Devendra, C., and Burns, M. (1983).Goat Production in the Tropics. 2<sup>nd</sup> ed. Farham Royal, Commonwealth Agricultural Bureau, pp 1 – 60.
- Hall, S.J.G. (1991). Body Dimensions of Nigerian Cattle, Sheep and Goats. *Anim. Prod.* 53:61-69
- Hassan, A. and A. Ciroma. 1990. Bodyweight measurements relationship in Nigerian Red Sokoto goats. Department of Animal Science, Usmanu Danfodiyo University, Sokoto, Nigeria <http://www.fao.org/wairdocs/ilri/x5520b/x5520b1d.htm>.



- Katongole J B D, Sebolai B and Madimabe M J 1996 Morphological characteristics of the Tswana goat. In S H B Lebbie and E Kagwini (eds). Proceedings of the Third Biennial Conference on the African Small Ruminant Research Network UICC, Kampala, Uganda –9 December 1994:43-47.
- Mamabolo, M.J. and Webb, E.C. (2005). Goat production Survey – Fundamental Aspects to Model Goat Production Systems in Southern Africa. Case Study of Agricultural Commission 2005.
- Mohammad, M.T., M. Rafeeq, M.A. Bajwa, M.A. Awan, F. Abbas, A. Waheed, F.A. Bukhari and P. Akhta. 2012. Prediction of body weight from body measurements using regression tree (RT) method for indigenous sheep breeds in Balochistan. Pakistan. J. Anim. Plant Sci., 22(1): 20-24.
- Ojedapo, L.O., Adedeji, T.A., Olayeni, T.B. Adedeji, O.S., Abdullah, A.R., Ojebiyi, O.O., (2007). Influence of age and sex on body weight and some body linear measurements of extensively reared wad goats in derived savannah zone of Nigeria. *Journal of Animal Veterinar Advances*. 6(1), 114- 117.
- Osuhor, C.U., J.P. Alawa and G.N. Akpa, 2002. Manure production by goats grazing native pasture in Nigeria. *Trop. Grassland*, 36: 123-125.
- Otoikhian, C.S.O., A.M. Otoikhian, O.P. Akporhwarho and C. Isidahoman. 2008. Correlation of body weight and some body measurement parameters in Quda sheep under extensive management system. *Afr. J. Gen. Agric*. 4(3): 129-133.
- Salako, A.E. and Ngere, L.O. (2002). Application of Multifactorial Discriminate analysis in the morphometric structural differentiation of West African Dwarf (WAD) and Yankassa Sheep in the South West. *Nigerian Journal of Animal Production* 29 (2): 163 – 167.
- SAS (1990). Statistical Analysis System (SAS). Institute Inc. (1990). SAS/STAT User's guide version 6, 4<sup>th</sup> edition Vol. 2 SAS Inst-Inc. carry, N.C 846.
- Semakula, J., Mutetikka, D., Kugonza, R. D. and Mpaiwe D. (2010). Variability in Body Morphometric Measurements and Their Application in Predicting Live Body Weight of Mubende and Small East African Goat Breeds in Uganda. *Middle-East Journal of Scientific Research* 5 (2): 98-105,
- Umoren, U.E. and Orok, E.J. (1994). Voluntary intake of selected indigenous browse species by west african dwarf goats. *Journal of Applied Chemistry and Agricultural Research* Vol. 1 (1) 1994: pp. 30-34.

Table 1: Least squares means (X), standard deviation (SD), and coefficient of Variation (CV) of Morphological traits among Red Sokoto, Sahel and WAD goat (< 1 year)

| RED SOKOTO | SAHEL | WAD |
|------------|-------|-----|
|------------|-------|-----|



| Parameter |    | Doe   | Buck  | Total  | Doe   | Buck  | Total | Doe   | Buck  | Total |
|-----------|----|-------|-------|--------|-------|-------|-------|-------|-------|-------|
| BW        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 13.73 | 12.86 | 13.30  | 13.56 | 14.94 | 14.25 | 10.44 | 10.62 | 10.53 |
|           | SD | 3.62  | 3.00  | 3.33   | 3.90  | 3.70  | 3.89  | 1.53  | 1.49  | 1.50  |
|           | CV | 26.38 | 23.36 | 25.12  | 29.40 | 24.78 | 27.30 | 14.65 | 13.98 | 14.27 |
| EL        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 10.22 | 9.86  | 10.04  | 11.52 | 11.06 | 11.29 | 9.56  | 9.43  | 9.49  |
|           | SD | 1.17  | 1.07  | 1.13   | 1.79  | 1.88  | 1.83  | 1.08  | 1.20  | 1.13  |
|           | CV | 11.46 | 10.81 | 11.24  | 15.53 | 16.96 | 16.28 | 11.27 | 12.70 | 11.95 |
| SW        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 13.34 | 13.83 | 13.59  | 18.18 | 18.59 | 18.39 | 12.75 | 12.99 | 12.87 |
|           | SD | 3.94  | 3.40  | 3.68   | 3.02  | 2.57  | 2.79  | 3.02  | 3.40  | 3.20  |
|           | CV | 29.57 | 24.65 | 27.06  | 16.60 | 13.80 | 15.20 | 23.70 | 26.21 | 24.90 |
| NC        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 22.69 | 23.43 | 23.06  | 21.02 | 23.31 | 22.17 | 23.47 | 24.39 | 23.93 |
|           | SD | 5.10  | 4.30  | 4.71   | 3.67  | 4.53  | 4.26  | 4.16  | 4.16  | 4.16  |
|           | CV | 22.48 | 18.35 | 20.41  | 17.47 | 19.43 | 19.22 | 17.73 | 17.04 | 17.40 |
| BL        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 41.92 | 41.97 | 41.95  | 35.36 | 39.56 | 37.46 | 42.04 | 41.88 | 41.96 |
|           | SD | 4.70  | 3.99  | 4.33   | 7.16  | 9.25  | 8.50  | 5.85  | 5.13  | 5.48  |
|           | CV | 11.20 | 9.50  | 10.33  | 20.25 | 23.39 | 22.68 | 13.92 | 12.27 | 13.06 |
| WH        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 52.06 | 51.63 | 51.85  | 42.47 | 45.80 | 44.13 | 47.93 | 48.07 | 48.00 |
|           | SD | 6.13  | 5.69  | 5.89   | 10.04 | 9.44  | 9.84  | 7.98  | 7.53  | 5.33  |
|           | CV | 11.78 | 11.03 | 11.36  | 23.64 | 20.62 | 22.30 | 16.65 | 15.67 | 16.08 |
| HG        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 53.33 | 57.53 | 55.43  | 49.90 | 48.77 | 49.33 | 50.15 | 49.71 | 49.93 |
|           | SD | 9.24  | 8.29  | 8.98   | 3.46  | 4.68  | 4.14  | 5.51  | 5.18  | 5.33  |
|           | CV | 17.33 | 14.40 | 16.21  | 6.74  | 9.60  | 8.39  | 10.98 | 10.43 | 10.67 |
| PG        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 55.37 | 59.61 | 57.49  | 52.20 | 50.63 | 51.41 | 51.42 | 50.99 | 51.19 |
|           | SD | 9.32  | 8.40  | 9.08   | 3.50  | 4.68  | 4.18  | 5.48  | 5.05  | 5.25  |
|           | CV | 16.83 | 14.09 | 15.79  | 6.70  | 9.24  | 8.14  | 10.65 | 9.91  | 10.25 |
| TL        | N  | 50    | 50    | 100    | 50    | 50    | 100   | 50    | 50    | 100   |
|           | X  | 12.08 | 11.71 | 11.807 | 11.62 | 11.97 | 11.79 | 10.29 | 10.27 | 10.28 |
|           | SD | 1.84  | 2.13  | 1.99   | 2.27  | 2.28  | 2.27  | 1.33  | 1.46  | 1.39  |
|           | CV | 15.24 | 18.17 | 16.71  | 19.52 | 19.06 | 19.25 | 12.90 | 14.24 | 13.52 |

Horn Length (HL), Ear Length (EL) Shoulder Width (SW) Neck Circumference (NC) Body Length (BL) Withers Height (WH) Heart Girth (HG), Pouch Girth (PG), tail length (TL), scrotal circumference (SC) and Body Weight (BW)

Table 2: Least squares means (X), standard deviation (SD), and coefficient of variation (CV) of Morphological traits in Red Sokoto, Sahel and WAD goat (1-2years of Age)

|           | RED SOKOTO |      |       | SAHEL |      |       | WAD |      |       |
|-----------|------------|------|-------|-------|------|-------|-----|------|-------|
| Parameter | Doe        | Buck | Total | Doe   | Buck | Total | Doe | Buck | Total |



|    |    |       |       |       |       |       |       |       |       |       |
|----|----|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| BW | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 13.60 | 13.51 | 13.56 | 17.13 | 16.86 | 16.99 | 15.45 | 17.65 | 16.55 |
|    | SD | 3.25  | 3.03  | 3.12  | 3.01  | 3.48  | 3.24  | 3.79  | 3.14  | 3.63  |
|    | CV | 23.85 | 22.40 | 23.03 | 17.56 | 20.61 | 19.04 | 24.51 | 17.81 | 21.96 |
| EL | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 12.24 | 11.88 | 12.06 | 13.82 | 12.81 | 13.32 | 11.25 | 10.52 | 10.89 |
|    | SD | 2.15  | 2.24  | 2.19  | 2.81  | 2.67  | 2.77  | 1.54  | 1.63  | 1.62  |
|    | CV | 17.58 | 18.83 | 18.17 | 20.30 | 20.80 | 20.79 | 13.70 | 15.47 | 14.87 |
| SW | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 13.31 | 13.54 | 13.42 | 23.24 | 22.58 | 22.91 | 14.12 | 16.23 | 15.18 |
|    | SD | 2.12  | 2.86  | 2.51  | 3.10  | 2.99  | 3.05  | 3.42  | 3.44  | 3.57  |
|    | CV | 15.91 | 21.11 | 18.66 | 13.32 | 13.26 | 13.30 | 24.21 | 21.19 | 23.54 |
| NC | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 25.87 | 26.16 | 26.01 | 25.33 | 25.97 | 25.65 | 24.73 | 24.69 | 24.71 |
|    | SD | 3.72  | 4.61  | 4.17  | 3.14  | 2.44  | 2.82  | 2.49  | 3.29  | 2.90  |
|    | CV | 14.37 | 17.63 | 16.03 | 12.38 | 9.43  | 10.98 | 10.07 | 13.34 | 11.76 |
| BL | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 50.93 | 46.83 | 48.88 | 61.14 | 53.76 | 54.77 | 51.84 | 49.23 | 50.54 |
|    | SD | 8.54  | 8.47  | 8.71  | 6.56  | 5.28  | 7.03  | 6.86  | 6.13  | 6.60  |
|    | CV | 16.76 | 18.10 | 17.82 | 10.73 | 9.81  | 12.84 | 13.24 | 12.44 | 13.07 |
| WH | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 55.47 | 53.60 | 54.53 | 60.58 | 60.12 | 60.63 | 52.20 | 49.37 | 50.79 |
|    | SD | 5.42  | 6.44  | 5.99  | 6.16  | 6.22  | 6.37  | 7.40  | 8.51  | 8.06  |
|    | CV | 9.78  | 12.01 | 10.99 | 10.17 | 10.33 | 10.52 | 14.17 | 17.24 | 15.87 |
| HG | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 63.92 | 62.14 | 63.03 | 60.58 | 59.87 | 60.23 | 60.80 | 54.73 | 57.77 |
|    | SD | 4.22  | 5.07  | 4.73  | 6.16  | 5.56  | 5.85  | 6.37  | 4.75  | 6.37  |
|    | CV | 6.61  | 8.16  | 7.50  | 10.17 | 9.29  | 9.72  | 10.47 | 8.68  | 1.103 |
| PG | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 66.13 | 64.49 | 65.31 | 62.98 | 62.06 | 62.51 | 62.54 | 56.01 | 59.28 |
|    | SD | 4.11  | 5.29  | 4.78  | 6.18  | 6.74  | 6.45  | 6.99  | 4.82  | 6.82  |
|    | CV | 6.21  | 8.20  | 7.32  | 9.82  | 10.86 | 10.32 | 11.19 | 8.61  | 11.51 |
| TL | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 12.27 | 12.79 | 12.53 | 13.48 | 13.43 | 13.45 | 10.95 | 11.56 | 11.25 |
|    | SD | 1.91  | 1.76  | 1.85  | 1.81  | 2.20  | 2.01  | 2.05  | 1.74  | 1.92  |
|    | CV | 15.55 | 13.80 | 14.74 | 13.49 | 16.41 | 14.94 | 18.78 | 15.08 | 17.07 |

Table 3 : Least squares means (X), standard deviation (SD), standard error (SE), and coefficient of variation (CV) of morphological traits among Red Sokoto, Sahel and WAD goat of Age > 2 years

| Parameter |   | RED SOKOTO |      |       | SAHEL |      |       | WAD |      |       |
|-----------|---|------------|------|-------|-------|------|-------|-----|------|-------|
|           |   | Doe        | Buck | Total | Doe   | Buck | Total | Doe | Buck | Total |
| BW        | N | 50         | 50   | 100   | 50    | 50   | 100   | 50  | 50   | 100   |



**NIGERIAN SOCIETY FOR ANIMAL PRODUCTION**  
**44<sup>th</sup> Annual Conference – ABUJA 2019**  
**Book of Proceedings**



|    |    |       |       |       |       |       |       |       |       |       |
|----|----|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| HL | X  | 18.95 | 19.21 | 19.80 | 23.60 | 25.12 | 24.36 | 19.13 | 19.35 | 19.24 |
|    | SD | 3.30  | 3.33  | 3.30  | 4.85  | 5.18  | 5.05  | 2.06  | 1.70  | 1.88  |
|    | CV | 17.39 | 17.34 | 17.29 | 20.55 | 20.64 | 20.74 | 10.76 | 8.80  | 9.78  |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| EL | X  | 7.47  | 7.29  | 7.38  | 10.53 | 10.76 | 10.64 | 7.25  | 6.81  | 7.03  |
|    | SD | 2.06  | 1.77  | 1.91  | 2.72  | 2.53  | 2.61  | 2.13  | 1.97  | 2.05  |
|    | CV | 27.54 | 24.22 | 25.87 | 25.82 | 23.50 | 24.56 | 29.36 | 28.94 | 29.21 |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| SW | X  | 12.33 | 11.88 | 12.11 | 13.71 | 13.70 | 13.71 | 11.22 | 10.64 | 10.93 |
|    | SD | 2.34  | 1.48  | 1.96  | 2.35  | 2.05  | 2.19  | 2.25  | 1.72  | 2.01  |
|    | CV | 18.90 | 12.43 | 16.19 | 17.11 | 14.97 | 15.99 | 20.06 | 16.19 | 18.44 |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| NC | X  | 19.65 | 19.93 | 19.79 | 24.97 | 24.71 | 24.84 | 14.98 | 15.40 | 15.19 |
|    | SD | 4.08  | 4.97  | 4.53  | 2.65  | 3.10  | 2.87  | 2.72  | 2.91  | 2.81  |
|    | CV | 20.77 | 24.96 | 22.88 | 10.61 | 12.55 | 11.56 | 18.13 | 18.91 | 18.49 |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| BL | X  | 24.91 | 24.66 | 24.76 | 27.16 | 26.52 | 26.91 | 27.70 | 27.15 | 27.42 |
|    | SD | 3.58  | 2.67  | 3.15  | 4.26  | 3.87  | 4.06  | 2.59  | 2.46  | 2.53  |
|    | CV | 14.38 | 10.83 | 12.69 | 15.70 | 14.54 | 15.09 | 9.36  | 9.06  | 9.22  |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| WH | X  | 45.01 | 45.93 | 45.47 | 58.53 | 56.28 | 57.40 | 43.76 | 43.69 | 43.72 |
|    | SD | 10.51 | 9.75  | 10.10 | 5.77  | 5.58  | 5.76  | 6.07  | 4.70  | 5.40  |
|    | CV | 23.35 | 21.23 | 22.20 | 9.86  | 9.92  | 10.03 | 13.88 | 10.76 | 12.35 |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| HG | X  | 58.17 | 58.56 | 58.37 | 70.51 | 72.02 | 71.26 | 46.62 | 45.42 | 46.02 |
|    | SD | 5.98  | 5.71  | 5.82  | 4.76  | 4.14  | 4.50  | 6.15  | 5.54  | 5.85  |
|    | CV | 10.28 | 9.75  | 9.97  | 6.75  | 5.74  | 6.31  | 13.19 | 12.20 | 12.72 |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| PG | X  | 61.15 | 60.64 | 60.89 | 64.61 | 65.69 | 65.15 | 68.93 | 68.57 | 68.75 |
|    | SD | 4.63  | 4.09  | 4.35  | 3.39  | 2.35  | 2.95  | 4.53  | 3.79  | 4.16  |
|    | CV | 7.57  | 6.74  | 7.15  | 5.25  | 3.58  | 4.53  | 6.58  | 5.52  | 6.05  |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
| TL | X  | 63.46 | 63.23 | 63.35 | 66.96 | 67.27 | 67.12 | 69.94 | 69.47 | 69.71 |
|    | SD | 4.62  | 4.05  | 4.32  | 3.43  | 5.46  | 4.54  | 6.50  | 6.75  | 6.60  |
|    | CV | 7.28  | 6.41  | 6.83  | 5.11  | 8.11  | 6.76  | 9.29  | 9.72  | 9.46  |
|    | N  | 50    | 50    | 100   | 50    | 50    | 100   | 50    | 50    | 100   |
|    | X  | 13.15 | 12.45 | 12.80 | 12.69 | 12.90 | 12.80 | 11.81 | 11.93 | 11.87 |
|    | SD | 2.13  | 1.71  | 1.96  | 1.92  | 2.17  | 2.04  | 1.57  | 1.43  | 1.49  |
|    | CV | 16.20 | 13.77 | 15.28 | 15.12 | 16.84 | 15.96 | 13.30 | 11.97 | 12.59 |

Horn Length (HL), Ear Length (EL) Shoulder Width (SW) Neck Circumference (NC) Body Length (BL) Withers Height (WH) Heart Girth (HG), Pouch Girth (PG), tail length (TL), scrotal circumference (SC) and Body Weight (BW)