

A STUDY OF COMMON QUAIL LICE INFESTATION IN SOKOTO METROPOLIS

¹ADAMU, Y. A.; ²ALAYANDE, M. O.; ⁴BELLO, A.; ⁴ONU, J. E.; ⁵UMARU, M. A.; ³SADIQ, Y. A.

¹Department of Veterinary Medicine; ²Department of Veterinary Parasitology and Entomology; ³Faculty of Veterinary Medicine; ⁴Department of Veterinary Anatomy; ⁵Department of Theriogenology and Animal production; Usmanu Danfodiyo University, Sokoto*Corresponding Author E-mail: musolade2000@yahoo.com

ABSTRACT

Two hundred Japanese quails from different locations in Sokoto metropolis were collected and examined between October and November for ectoparasites. All the birds were dusted with Rambo[®] (Permethrin 0.60%) powder for lice recovery. The recovered lice were then cleared with lactophenol overnight, before been examined under the light microscope (x 40). Of the 200 quails examined, only 4 (2%) were infested with lice. The lice recovered were *Menacanthus sp* 2%, *Lipeurus sp* 2% and *Gonoides sp* 0.5%. No other ectoparasite was found. Lice infestation constitutes the common ectoparasitism of Japanese quails in Sokoto metropolis and is a very low, with little or no observable harm.

INTRODUCTION

Poultry is an acceptable form of animal protein to most people throughout the world, with the exception of strict vegetarians and vegans (Smith, 2000). Animal protein consumption per head is greater in developed countries than in developing countries such as Nigeria with a population of over 140 million people, over the years has not been able to meet the food and agricultural organization (FAO) recommended minimum protein intake requirement of 65gm per person per day (Musa *et al.*, 2008). The number of poultry in the world is large and domestic chickens contribute 90% to poultry populations (Smith, 2000). So many factors such as inadequate supply of day old chicks, diseases, high cost of drugs and biological and the time taken to raise the birds all these led to search for an alternative cheaper source of poultry production and subsequent introduction of quail birds in Nigeria. Japanese quails also referred to as Coturnix quails (*Coturnix coturnix japonica*) is an old world bird found in the East Asia. They are migratory species of birds which dwell on grasslands and cultivated fields (Musa *et al.*, 2008). Japanese quails were widely domesticated for their meat and egg in that country. This led to sharp increase in the population of Japanese quails. In Nigeria, Japanese quails were introduced at National Veterinary Research Institute, Vom (NVRI) Plateau State, with the support of Directorate for Food, Road and Rural Infrastructure (DFRRI). Diseases from endoparasites and ectoparasites origins may however limit the production of quail. Ectoparasites cause a lot of damage and

economic losses in poultry production. In view of this, there is need to investigate the ectoparasites affecting Japanese quails in the study area and their effect on the production.

MATERIALS AND METHODS

Study area and Laboratory Identification

The study was carried out in Sokoto, Sokoto State, located in the extreme north western part of the country, covering between longitudes 5°14'E and latitudes 13°04'N. A total of 200 quail birds collected from different residential locations of the metropolis were examined. Each bird was examined carefully, taking cognizance of each region, the head, neck, wings, and breast/keel muscle and around the vent. Each bird was then placed on a white background (about 60x60cm white cardboard), then dust-ruffled with about 2gm of Rambo[®] (Cypermethrin) powder to recover the lice from the birds as described by Clayton and Drown (2001). Plain thumb forceps were then used to pick the ectoparasites. The recovered lice were then stored in 70% alcohol for identification after clearing in Lactophenol for about 12 hours. The identification of lice was as described by Ansari (1955), Emerson (1956) and Matsudaira and Kaneko (1969). Both stero- and binocular microscopes were used for the identification, at magnification of X40.

RESULTS

Out of the 200 birds examined, only four were positive representing 2% (Table 1). The only ectoparasite found was lice. Four different species of lice were identified (Table 2).

DISCUSSION

This study conducted showed very low prevalence of ectoparasitism on Japanese quails in Sokoto (2%). The ectoparasites that were obtained on the quails were lice of the genera *Menacanthus* sp, *Lipeurus* sp and *Gonoides* sp with prevalence of 2%, 2%, and 0.5% respectively. This low prevalence can be as a result of the management systems due to the fact that quails are kept in cages or deep litter system with improved biosecurity measures. This low prevalence was similar to that obtained on wild quails in Turkey by Nursel (2010). In addition to other lice; *Cuelotogastercinereus* was obtained, while in this study, *Lipeurus* sp. was identified. No significant lesion was observed to be associated with any of the species obtained in this research. Doster *et al.*, (1980), also recorded similar result on quails in south-eastern United States. The presence of the lice on the Japanese quails is due to the fact that mallophagans' spend

their entire lifecycle on the host (Keymer, 1982) irritation and discomfort arises only when there is very high infestation. Though lousiness can be a problem, but effects are only observed when the number on the host is very high. Some of the observable problems include; irritation, anorexia, pale mucous membrane, loss in production and reduced egg production. It is important that for maximum production, adequate preventive measures should be taken into consideration so as not to allow the infestation of lice exceed the level at which the impact of lice will manifest. This is only achieved by proper management system. Currently, this study showed that there is low ectoparasitic infestation on Japanese quails in Sokoto. Farmers are advised to embark on Japanese quail production and ensure more biosecurity/control measures. There is need to improve the management system so as to prevent the occurrence of other parasites.

REFERENCES

- Ansari; M.A.R. (1955) synoptic table for the determination of mallophaga infesting the domestic fowl (*Gallus gallus domestica*). *Indian Journal of Entomology* 17:245-270
- Clayton, D.H. and Drown, D.M. (2001) critical evaluation of five methods of quantifying chewing lice (Insecta: Phthiraptera). *Journal of parasitology* 87(6) 1291 – 1300.
- Doster, G.L. Wilson N, Kellog, F.E (1980) Ectoparasite collected from Bibwile quail in the south eastern united state. *Journal of Wildlife diseases* 16(4) 515-520.
- Emerson; K.C (1965) Mallophaga (Chewing lice) occurring on domestic chicken journal of Kansas Entomological Society 29(2): 63-73.
- Keymer, I.F (1982) parasitic diseases, Diseases of cage and aviary birds 2nd ed lea febiger, Philadelphia.
- Matsudaira, Y. and Kaneko, K. (1969) notes on biting lice (Mallophaga) collected from domestic fowl, *Japan santory zoology* 20; 219-229
- Musa, U. Haruna, E.S. Lobin. L.H (2008) Quail production in the topics. 1st Edition 63pp
- Nursel, A. (2010) The presence of chewing lice (Insecta: Phthiraptera) sp. On wild Quails. *Journal of Animal and Veterinary Advances* 9(9): 1377-1379
- Smoth. I.S. (2000) *Poultry Problem and Prospects in the Tropics* (2nd Edition). Alena Publishers Enugu. Enugu State, Nigeria

Table 1- Prevalence of Ectoparasite on Japanese Quails

No of samples	No of positive samples	Prevalence %
200	4	2

Table 2- Prevalence of Lice species (lice) on Japanese Quails

Ectoparasite	No of samples	No of positive samples	Prevalence %
<i>Menacanthus</i> sp.	200	4	2
<i>Lipeurus</i> sp.	200	4	2
<i>Gonoides</i> sp.	200	1	0.5

Plate1 *Gonoides* sp, X4



Plate 2: *Menacanthus* sp. X4





Plate 3 *Lipeurus* sp. (X 4 magnifications)