EFFECT OF FEEDING WHOLE MILLET ON THE PERFORMANCE OF EGG PRODUCTION OF LAYER BIRDS

M.L. ABUBAKAR

Department of Science Laboratory Technology, Federal Polytechnic Kaura Namoda, Zamfara State, Nigeria.

ABSTRACT

This research was conducted to determine the effect of feeding whole millet on the diet of Isa brown layers breed. Four hundred and twenty birds aged 43 weeks were randomly assigned to the following dietary treatments: Treatment 1 contained ground maize with 18% C.P (Control). Treatment 2, 3 and 4 contained ground millet with 16, 18 and 20% C.P. Treatment 5, 6 and 7 contained whole millet with 16, 18 and 20% C.P feed intake did not differ significantly between the treatments. Egg productions were generally better with millet based diets. Egg production was significantly higher ($P<0.05$) for birds fed ground millet with 18% C.P compared to those fed the maize based diets. It was concluded that birds fed with the millet based diets have better performance as indicated in the study results.

INTRODUCTION

The term poultry refers to a rather wide variety of birds of several species. Thus, it applies to chickens, turkeys, ducks, geese, guinea fowls, pigeons, ostriches, quails, and other game birds. Zoologically the fowl belongs to the family *phasianidae* and the genus Gallus. It is simply called Gallus domesticus. The ancestors of the domestic fowl originated from south East Asia and have been subjected to extensive breeding for size, colour pattern, conformation and egg laying ability (Payne, 1990). In Nigeria, daily protein intake is estimated at 53.8kg per day, with only 8.4g being of animal origin. This low intake of protein leads to high rate of infant mortality, low disease resistance, poor growth and development, mental retardation, kwashiorkor etc. all these could be prevented by consuming adequate quantities of high quality protein (Awosanmi, 1999). Poultry play a significant supportive role to various food productive systems by providing wastes and by products used in other agricultural systems. For example poultry droppings are used as manure that replenishes soil nutrients (Atteh, 2004). In order to meet the animal protein requirement, poultry have been singled out as animals of first choice. This is due to their high and quick turnover. Furthermore, they are easy to manage, highly productive, with short generation interval; and they adapt to a wide range of ecological conditions (Awosanmi, 1999).This research was aimed at determining the effect of feeding millet either whole or ground as the major source of energy in the diets of layers.

MATERIALS AND METHODS

Experimental Location:

This study was conducted at the Sokoto Veterinary Clinic, Ministry of Agriculture and Natural Resources, Aliyu Jodi Road, Sokoto. Sokoto is located in the Sudan Savannah Zone in the extreme north western part of Nigeria, and lies between latitudes 12^0N and 13^0N and longitudes 40 and 60E (Mamman *et al.*, 2000) Sokoto has low humidity and high solar radiation. Maximum temperature of 42°C has been reported in May and 13°C has been reported in January during the cold dry season (harmattan)..

Experimental Procedure:

A total of four hundred and twenty ISA Brown layers of 43 weeks of age were used in the experiment. The birds were randomly allocated to seven treatments. Each treatment contained ten replicates with six birds per replicate. Seven diets (representing treatments) were compounded as follows: Treatment 1 contained ground maize with 18% crude protein. Treatment 2 contained ground millet with 16% crude protein. Treatment 3 contained ground millet with 18% crude protein. Treatment 4 contained ground millet with 20% crude protein. Treatment 5 contained whole millet with 16% crude protein. Treatment 6 contained whole millet with 18% crude protein, and Treatment 7 contained whole millet with 20% crude protein. The gross and chemical compositions of the experimental diets are shown in tables 1 and 2 respectively. The experimental diets and water were given ad-libitum. Feed intake, egg production and mortality were monitored daily. The data collected was subjected to analysis of variance (AVOVA) using the STAT VIEW (SAS, 2002). Statistical package.

Table 1: Gross composition of the experimental diets

Ingredients (%)	T_1 Maize (18%CP)	T_2 and T_5* Millet (16%CP)	T_3 and T_6** Millet (18%CP)	T_4 and T_7*** Millet (20%CP)
Maize	30.7	0.0	0.0	0.0
Millet	0.0	37.1	31.6	26.3
GNC	21.4	13.9	20.4	26.7
Wheat offal	36.8	37.8	36.9	36.0
Limestone	9.00	9.00	9.00	9.00
Bone meal	1.40	1.40	1.40	1.40
Premix	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25
Methionine	0.11	0.13	0.12	0.10
Lysine	0.07	0.16	0.08	0.00
Total	**100**	**100**	**100**	**100**

*T2 ground millet and T5 whole millet, **T3 ground millet T6 whole millet, ***T4 ground millet and T7 whole millet.

Table 2: Calculated chemical composition of the experimental diets.

Energy (kcal/kg)	2501	2501	2501	2501
CP (%)	18	16	18	20
Lysine (%)	0.8	0.8	0.8	0.8
Methionine (%)	0.4	0.4	0.4	0.4
Calcium (%)	3.6	3.6	3.6	3.6
Phosphorous	0.4	0.4	0.4	0.4
Crude fibre (%)	4.8	6.9	6.7	6.5
Other extract	4.0	4.0	4.0	4.0

RESULTS AND DISCUSSION

The results showed that generally egg production was better with the millet based diets. however, the differences were significant only between T4 (whole millet, 20%CP) and T1 (ground maize, 18%CP). Feed intake did not differ significantly between the treatments. Mortality was significantly lower for T6 compared to T3. These results therefore indicate that neither substituting maize with ground or whole millet nor varying the protein levels in the millet containing diets had any effect on feed intake. It is known that feed intake in poultry is governed by dietary energy concentration.

Conclusion

The results of this study indicate that feeding either maize or millet (whole or ground) containing 16 – 20% CP did not significantly affect feed intake. Egg production was better with the millet based diets, with significant differences between those fed ground millet with 20% CP and those on ground maize with 18% CP.

REFERENCES

Atteh J.O. (2004). *Theory and Practice of Poultry Production*, Adhec Printers, Ilorin Nigeria.

Awosanmi V.O. (1999). *Nigeria needs to recover from its present state of poultry production.* International Journal of Animal Science, 2(1):21-26.

Mamman, A.B., Ojebanji, J.O. and Peters, SW. (2000). (Eds.). *Nigeria A people united, Afuture assured (survey of states).* Vol.2. Gabumo Publishing Co-Ltd. Calabar, Nigeria.

Payne, W.J.A. (1990). *Introduction to animal Husbandry in the tropics,* 4th edition, Longman publishers, India.

SAS (2002).*Startview Statistical Package (English Version). Statistical analysis system, SAS Inc.* New York.

Table 3. Feed intake and mortality

Parameter	T1 Ground Maize	T2 Ground Millet	T3 Ground Millet	T4 Whole Millet	T5 Whole Millet	T6 Whole Millet	T7 Whole Millet	SEM

	(18%CP)	(16%CP)	(18%CP)	(20%CP)	(16%CP)	(18%CP)	(20%CP)	
Feed intake (g/b/d)	100.91	104.56	101.31	105.94	102.95	104.91	105.40	1.96
Egg production (g/b/d)	43.96^{a}	48.75^{ab}	51.88^{ab}	52.50^{b}	51.67^{ab}	47.71^{ab}	47.71^{ab}	2.94
Mortality (%)	3.33^{ab}	1.66^{ab}	6.66^{a}	3.33^{ab}	4.99^{ab}	0.00^{b}	3.70^{ab}	2.11

Standard Error of Means on the same row with different superscript are significantly different (P<0.05). SEM :- standard error of means.